



Septate Uterus. Resect or not? That is Not the Only Question

Tayfun Cok¹

¹ Department of Obstetrics and Gynecology, School of Medicine, Adana Research and Education Centre, Baskent University, Adana, Turkey.

Rev Bras Ginecol Obstet 2022;44(9):899–900.

Address for correspondence Tayfun Cok, MD, Associate Professor of Obstetrics and Gynecology, Department of Obstetrics and Gynecology, Adana Research and Education Centre, School of Medicine, Baskent University, 01250, Adana, Turkey (e-mail: tayfuncok@yahoo.com).

The septate uterus is the most common congenital anomaly of the uterus. The prevalence of septate uterus varies between 2.3 and 15.4% according to the population studied, with the percentages being higher in women with recurrent miscarriages combined with infertility.¹ A septate uterus increases adverse pregnancy outcomes such as miscarriage, preterm birth, and malpresentation of the fetus, although its direct association with infertility is likely but still controversial.^{2,3} This issue has also provoked a vivid debate on hysteroscopic resection of the septate uterus for treatment.⁴ Especially for the management of the partial uterine septum, the main point which affects the resection decision lies within the problem of diagnosis. There has been a continuous change in the diagnostic criteria of the partial septate uterus along with the new guidelines, since 1988, despite increasing diagnostic tools.^{5–9} Even in the same group of patients, the prevalence of septate uterus has been reported to vary from 5 to 31%, according to the criteria that have been used.¹⁰ One patient flying from Europe to the USA may lose her septum over the Atlantic Ocean. A patient with a 12 mm septal indentation, who used to have a septate uterus according to the American Fertility Society (1988),⁵ lost her septum in 2016, according to the American Society for Reproductive Medicine (ASRM) 2016⁷ guidelines and gained it back in the ASRM 2021 updated guidelines.⁹ The European Society of Human Reproduction and Embryology (ESHRE) and the European Society for Gynaecological Endoscopy's (ESGE) shared guidelines (2016)⁷ boosted the number of patients with a diagnosis of the septate uterus, whereas ASRM (2016)⁷ underestimates the number of patients with partial septate uterus and leaves some patients in a gray zone. Some of this underestimation is due to the 90° angle criteria, which seem to be arbitrary and are not based on strong evidence.¹⁰ In the only randomized controlled The Randomised Uterine

Septum Trial (TRUST) of uterine septum resection, 90% of the patients had a partial septate uterus, and three different criteria were used throughout the study. However, most importantly, the difference in the diagnosis reflects in the preference of the treatment regarding resection, and the interpretation of the outcomes of septum resection according to various diagnostic criteria lacks guidance in clinical practice.¹¹ After the TRUST results, we have the option to discuss expectant management with patients. In those patients who choose expectant management, we can find out cut-off values for septal indentation length and the angle for poor obstetric outcomes, if there is any relation.

Conflicts of Interests

The authors have no conflict of interests to declare.

References

- 1 Chan YY, Jayaprakasan K, Zamora J, Thornton JG, Raine-Fenning N, Coomarasamy A. The prevalence of congenital uterine anomalies in unselected and high-risk populations: a systematic review. *Hum Reprod Update*. 2011;17(06):761–771. Doi: 10.1093/humupd/dmr028
- 2 Carrera M, Pérez Millan F, Alcázar JL, et al. Effect of hysteroscopic metroplasty on reproductive outcomes in women with septate uterus: systematic review and meta-analysis. *J Minim Invasive Gynecol*. 2021;***: [ahead of print]. Doi: 10.1016/j.jmig.2021.10.001
- 3 Panagiotopoulos M, Tseke P, Michala L. Obstetric complications in women with congenital uterine anomalies according to the 2013 European Society of Human Reproduction and Embryology and the European Society for Gynaecological Endoscopy Classification: a systematic review and meta-analysis. *Obstet Gynecol*. 2022;139(01):138–148. Doi: 10.1097/AOG.0000000000004627
- 4 Alvero R, Burney RO, Khorshid A, et al. Surgical treatment of uterine septum to improve reproductive outcomes - resect or not? *Fertil Steril*. 2021;116(02):298–305. Doi: 10.1016/j.fertnstert.2021.06.009

received
January 2, 2022
accepted
January 13, 2022
published online
August 29, 2022

DOI <https://doi.org/10.1055/s-0042-1754377>.
ISSN 0100-7203.

© 2022. Federação Brasileira de Ginecologia e Obstetrícia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

- 5 The American Fertility Society classifications of adnexal adhesions, distal tubal occlusion, tubal occlusion secondary to tubal ligation, tubal pregnancies, müllerian anomalies and intrauterine adhesions. *Fertil Steril.* 1988;49(06):944–955. Doi: 10.1016/s0015-0282(16)59942-7
- 6 Grimbizis GF, Gordts S, Di Spiezio Sardo A, et al. The ESHRE/ESGE consensus on the classification of female genital tract congenital anomalies. *Hum Reprod.* 2013;28(08):2032–2044. Doi: 10.1093/humrep/det098
- 7 Grimbizis GF, Di Spiezio Sardo A, Saravelos SH, et al. The Thessaloniki ESHRE/ESGE consensus on diagnosis of female genital anomalies. *Gynecol Surg.* 2016;13:1–16. Doi: 10.1007/s10397-015-0909-1
- 8 Ludwin A, Martins WP, Nastri CO, et al. Congenital Uterine Malformation by Experts (CUME): better criteria for distinguishing between normal/arcuate and septate uterus? *Ultrasound Obstet Gynecol.* 2018;51(01):101–109. Doi: 10.1002/uog.18923
- 9 Pfeifer SM, Attaran M, Goldstein J, et al. ASRM müllerian anomalies classification 2021. *Fertil Steril.* 2021;116(05):1238–1252. Doi: 10.1016/j.fertnstert.2021.09.025
- 10 Ludwin A, Ludwin I, Coelho Neto MA, et al. Septate uterus according to ESHRE/ESGE, ASRM and CUME definitions: association with infertility and miscarriage, cost and warnings for women and healthcare systems. *Ultrasound Obstet Gynecol.* 2019;54(06):800–814. Doi: 10.1002/uog.20291
- 11 Rikken JFW, Kowalik CR, Emanuel MH, et al. Septum resection versus expectant management in women with a septate uterus: an international multicentre open-label randomized controlled trial. *Hum Reprod.* 2021;36(05):1260–1267. Doi: 10.1093/humrep/deab037