From India—International Aortic Summit: There Is Nothing Impossible for Those Who Will Try!

Mohammed Idhrees, MS, MCh¹ Mohamad Bashir, MD, PhD, MRCS^{1,2,3} Bashi V. Velayudhan, MBBS, MS, MCh¹

¹ Institute of Cardiac and Aortic Disorders, SRM Institutes for Medical Science (SIMS Hospitals), Chennai, Tamil Nadu, India

³Department of Vascular and Endovascular Surgery, Velindre University National Health Service Trust, Health Education, and Improvement Wales (HEIW), Cardiff, United Kingdom

Aorta (Stamford) 2022;10:219-223.

Address for correspondence Mohammed Idhrees, MS, MCh, Institute of Cardiac and Aortic Disorders, SRM Institutes for Medical Science (SIMS Hospitals), Chennai, Tamil Nadu 600026, India (e-mail: a.m.idhrees@gmail.com).

The aorta is a dynamic living structure, central to all vital organs with its aortovascular branches. At the center of the affair, the aorta is afflicted with aneurysm disease, a complex condition that warrants attention and a multidisciplinary huddle to understand and ameliorate its dire effects. The International Aortic Summit in Chennai, India, is not just a conference, but a confluence of multidisciplinary specialties and teams joined around a rounded table of innovation and collaboration.

The 9th International Aortic Summit 2020 (IAS2020) was a witness to such confluence and was not hindered by the untoward effects of the pandemic. We streamed live globally bringing esteemed colleagues to deliver an eye and reflective practice on their experiences and expertise. Viewed globally by over 42 countries, IAS2020 was organized into structured sessions and platforms to cover all profiles of aortic disease and surgery.

Aortic surgery practice in India was a major focus during the Summit. Discussants touched the outcome and available resources. The aim was to incentivize practicing surgeons and young aspiring colleagues to form a solid practice and surgical approach structured, devised, and based on clinical evidence and research.

Sessions dedicated to pioneers in aortic surgery were a mere reflection of sheer and driven practices. Those were imperative to impose the way the future is evolving and to incite young surgeons to follow pursuit.

Keynote speakers including Drs Joseph Coselli and Joseph Bavaria delivered their shrewd expert opinions on aortic surgery over their decades of experience; the key points and

received June 29, 2022 accepted after revision August 8, 2022 DOI https://doi.org/ 10.1055/s-0042-1756672. ISSN 2325-4637. sessions are listed in **- Tables 1** and **2**. There were discussions on a wide range of topics related to thoracic and abdominal aortic aneurysms, aortic dissections, genetics pertaining to the aorta, interventional aortic radiology, and novel endovascular aortovascular innovations. The summit was a relishing opportunity to the current trend of aortic surgery in India and advances made across the years, best told by B.V.V., the Summit Director and propeller of innovation in aortic surgery management in India.

Following on to IAS2020, we developed a strong collaboration with AORTA and reflected on the key topics selected from the core of IAS2020. The selected topics were the crux of the two consecutive days of the summit and formed this focused issue. The articles collated were debates on pertinent topics which remain controversial and require profound attention. Practical sessions highlighted during the summit were invigorated and supported by contributors including innovative surgical approaches that emerged over the past years. The endovascular side of the program was thoroughly covered adding a touchstone and face value to furthering aortic surgery in current and future eras. The thoracoabdominal part was again a station that intrigued the audience during the Summit and after. As such, articles were collated to support this stem and to a prime foundation for future events.

With the success of the 9th IAS, we are promising our readers, subscribers, and followers that the 10th IAS will see through live operating workshops performed by esteemed colleagues' debates to induce guidance and appropriate decision-making, and case-based discussions to highlight surgical perspectivity (**-Fig. 1**).

² Department of Neurovascular Research Laboratory, Faculty of Life Sciences and Education, University of South Wales, Wales, United Kingdom

^{© 2022.} The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/)

Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA

Table 1 Salient features of keynote lectures

Prof. Joseph Coselli, Houston, TX Thoracic Aortic Aneurysm Surgery: Don't Quit:
Cardiac surgical devices and prostheses have undergone multiple innovations and developments in the last few decades.
To work with our colleagues closely-cardiologists, radiologists, and vascular surgeons.
Enumerated the developments in aortic arch surgery over the years.
Prof. Joseph Bavaria, Philadelphia, PA Type A Dissection: From the 30/30 club disaster to an upcoming Innovative decade
Most important concept in future is in regard to "distal aorta":
Index proximal operations will be driven and conceived by the availability of new technology endografts.
Proximalization of the conduct of aortic arch operations will continue.
Index operations "tactics" will be driven to reduce total cardiopulmonary bypass time.
Achieve better accumulation of outcomes data on the global stage.

Table 2 The key points of all the sessions over 2 days

Day 1		
Aortic Surgery in Asia–I:		
Dr. Kay-Hyun Park, South Korea Extensive thoracic aortic replacement (ascending to distal descending thoracic) without FET via a single median sternotomy	 Extensive thoracic aorta replacement is feasible via median sternotomy and transpericardial approach Frozen elephant trunk should be reserved for desperate cases who cannot undergo two-stage operation at a short interval 	
Dr. Worawong Slisatkorn, Thailand Total arch replacement in octogenarian	 Total arch replacement can be performed in octogenarians with acceptable outcomes Team expertise and close perioperative care are mandatory to improve the operative result 	
Dr. Saeid Hosseini, Iran Outcomes of reoperation after acute Type A aortic dissection	 Challenging issues include Whether to repair or replace the aortic root? The effect of the thrombosed false lumen on reoperation, partial thrombosis an independent risk factor The extent of distal operation in the index surgery 	
Dr. Thodur Vasudevan, New Zealand Total body floss: a concept in difficult endo- vascular repair and a novel case	 Unusual method of through and through wire access Multiple tortuosities Out of the box solutions Wire from the apex to femoral vessels Substernal access 	
Dr. George Joseph, India Total arch fenestrated endovascular repair af- ter surgical replacement of ascending aorta	 Residual Type A dissection can be treated by endovascular total arch fenestrated repair Short, kinked ascending aortic surgical grafts pose a challenge to arch TEVAR Surgeons should attempt to replace as much of the ascending aorta as possible during the initial surgery and provide a landing zone for future arch TEVAR 	
Dr. Murali Krishnawami, India Thoracoabdominal aneurysm—total endovas- cular repair	 FEVAR needs careful planning and meticulous execution Needs multiple accesses and multiple catheters, wires, and sheaths Can be performed by experienced centers with excellent results 	
Dr. Arunkumar, India The current role of transesophageal echocar- diography in acute aortic syndrome	 Transesophageal echocardiogram plays a significant role inside the operative room A good adjuvant tool for open aortic repair and endovascular procedures 	

Table 2 (Continued)

Day 1		
Valve in Acute Aortic Dissection:		
Dr. Malakh Shrestha, Germany Spare the aortic valve—I do it my way!	 If aortic root is dissected but the aortic valve is normal Normal sized aortic sinus—sandwich technique Aortic root dilated—David's procedure when possible Short- /long-term results of the David-I procedure in Marfan's patient are excellent Long-term survival of hospital survivors after David-I procedure in acute dissection is comparable to elective settings 	
Dr. Roberto Di Bartolomeo, Italy Aortic valve sparing with Valsalva graft	 Valve sparing aortic root replacement has excellent long-term results Bologna experience: Long-term survival at 1 year is 96.7%, 5 year is 93.9%, and 10 year is 89.5% 	
Dr. Laurent de Kerchove, Belgium When and how I repair the aortic valve in chronic and acute aortic regurgitation	 Aortic valve repair for pure aortic insufficiency necessitates a good understanding of the underlying mechanism and a systematic repair algorithm Optimal durability is achieved by optimization of coaptation and by annulus (± root) stabilization 	
Dr. Ruggero DePaulis, Italy Biological or mechanical conduits in aortic root dissection?	 Both choices are certainly defendable Life expectancy and the possibility of valve in valve certainly is a push for larger use of bio conduits Proper tailoring for each patient remains the main strategy 	
Dr. Michael Borger, Germany Management of iatrogenic Type A aortic dissection	 Iatrogenic aortic dissection is a rare but dangerous complication of cardiac surgery and cardiac catheterization Despite elevated risk, selected patients should be operated on emergently Early mortality is higher for the condition than patients presenting with spontaneous Type A dissection, particularly in postcardiac surgery patients 	
Aortovascular Science:		
Dr. Wael Awad, United Kingdom Risk prediction model in aortic aneurysm sur- gery—unmet clinical equipoise	 Thoracic aortic aneurysm (TAA) is a dangerous, deadly, and silent disease that is notoriously difficult to detect prior to complications Early diagnosis and management remain a critical component for limiting mortality from TAA prior to dissection Greater understanding of the genetics of these pts and their specific genetic mutations can provide personalized aortic care, tailoring surgical recommendations for each patient 	
Dr. Benjamin Adams, United Kingdom Does adding a root replacement in Type A aortic dissection repair provide better outcomes?	 Addition of ARR in TAAD setting does not increase mortality or postoperative complications ARR should be considered at the time of the emergency repair, especially in young patients with a degree of root dilatation and/or aortic valve regurgitation 	
Dr. John Elefteriades, United States Update on genetic dictionary	 A new era of molecular identification of individuals at risk for thoracic aortic aneurysm Personalized strategic management of these individuals 	
Dr. Ourania Preventza, United States Extent of repair for Type I. How long and why?	 Distal extent preferably hemiarch, unless a total arch replacement is necessary Total arch replacement is performed when the entry tear is within the greater curvature, severe compression of the true lumen When stented, a 10-cm antegrade delivery is preferred 	
Dr. Arminder Jassar, United States The changing surgical approaches for Type A dissection	 Ascending aortic replacement will suffice for many, but not all, patients with acute Type A dissection While the goal of the operation is to have an alive patient, it is also reasonable to plan for the future For patients with acute malperfusion, alternative strategies or adjunct techniques should be considered to ensure reperfusion 	

(Continued)

Table 2 (Continued)

Day 2		
Dr. Bashi Velayudhan, India Acute aortic syndrome (AAS) in India—What do we know?	 Possibility of a lower incidence of the acute aortic syndrome as compared to the western world (due to diabetes mellitus) Majority of patients are in the younger age group Last 15 years—more centers to handle AAS Distal aorta in Marfan's syndrome—still a challenge 	
Dr. Shiv Choudhary, India Inexpensive way to manage the arch in acute Type A dissection	 Lack of expensive devices should not preclude lifesaving surgery in aortic dissection It is possible to operate with successful results in a resource constrained setup 	
Dr. Zile Meharwal, India Raising the bar Hemodynamics in the small aortic root	 Small aortic root poses a challenge for surgeons Patient prosthesis mismatch (PPM) is common with small aortic root All measures should be taken to avoid PPM 	
Dr. Mohammed Idhrees, India Decision-making in acute aortic dissection	 Decision-making in the management of acute aortic dissection is crucial When Plan A is not smooth, always choose Plan B Select the best and simplest option which suits the team 	
Dr. Karthikeyan, India Gated CTA of aorta–optimal imaging of aortic disease	 Imaging with CT plays a central role in diagnosis to allow expedited management Helps for clinical risk assessment and establishing a definitive diagnosis CT scans to be performed with aim of motion-free images Dedicated injection protocol should be used 	
TEVAR/Hybrid Aortic:		
Dr. Heinz Jakob, Germany FET innovation	E-vita open neo has a family of graft variationsThe graft suits all kinds of aortic pathologies	
Dr. Nimesh Desai, United States Sequential branched arch TEVAR	 Index arch operations will be driven and conceived by the new availability of new technology endografts 	
Dr. Martin Czerny, Germany Midterm results of branched endovascular aortic arch repair	 Total endovascular aortic arch is a safe and reproducible technique Primarily for nonsurgical candidates Pathology determines the mode of treatment Creation of aortic centers with the entire armamentarium will aid in doing the right things in the right patients Will reduce the need for combined vascular/endovascular procedures 	
Dr. Martin Grabenwoger, Austria Management of Type 1 endoleak following TEVAR	 Treatment strategy for Type I endoleak after TEVAR has to be decided on a case-by-case basis Decision endovascular-hybrid-surgery (FET) dependent on- Anatomy of the aortic arch Risk profile of the patient (suitable for surgery?) Is it possible to create a proximal landing zone for an endovascular extension by aortic arch rerouting procedures? Open surgery by the FET technique can be performed with good results when Type I endoleak cannot be treated by endovascular techniques? 	
Dr. Edward Chen, United States Redo-aortic arch surgery	 Redoaortic arch surgeries are high-risk and technically demanding procedures performed with acceptable morbidity and mortality Careful operative planning and execution are paramount for optimal clinical outcome Operative outcomes were impacted by several factors including cardiac function and surgical complexity, but not by prior aortic procedures 	
Dr. Maximilian Kreibich, Germany dSINE for FET and TEVAR	 dSINE may develop at any time after FET procedure and the risk of dSINE development after FET procedure is substantial No independent predictors for the development of dSINE were identified, but Thoraflex has a stiffer distal end as compared with E-Vita Reinterventions for dSINE were associated with a very good clinical outcome 	
Dr. Cherri Abraham, United States Treatment of PAU and IMH in the thoracic aorta	 Penetrating aortic ulcer and intramural hematoma are both clinically complex and part of a clinical spectrum of acute aortic syndromes Treatment strategies include reduction in aortic wall stress and tailoring the surgical approach to the patient and lesion 	

Table 2 (Continued)

Day 2		
Downstream Aorta:		
Dr. Anthony Estrera, United States Open techniques in TAAAR in aneurysm repair	 Individualized standard of approach for aneurysm patient using FLAP (fragility/life expectancy/anatomy/pathology) Vital importance of the team effort Open and endovascular techniques are complimentary to each other and not competitive with one another 	
Dr. Joseph Coselli, United States Open techniques in TAAAR in chronic dissection	 Patients with chronic dissection tend to undergo extensive thora- coabdominal repair A variety of operative strategies are targeted to aortic dissection When compared with patients without dissection, those with chronic dissection are inclined to get extensive repair 	
Dr. Leonard Girardi, United States Emergency thoracoabdominal aortic aneu- rysm scenario management	 Thoracoabdominal aneurysm emergencies are high-risk undertaking Endovascular and open repair techniques performed with similar operative risk in experience centers Open repair is associated with fewer early- /late- reinterventions 	
Dr. Roberto Chiesa, Italy Open thoracoabdominal following endovas- cular intervention	 Close follow-up is essential after TEVAR Open TAAAR conversion is technically challenging Surgery can be performed with acceptable results in centers with an "aortic team" Increased mortality in cases of infection and retrograde dissection 	
Dr. Germano Melissano, Italy Spinal cord ischemia in open and endovascular repair	 Spinal cord ischemia prevention requires optimizing all aspects of the procedures CSF drainage is a valid adjunct; however, it comes with several potential serious complications The problem is still not resolved, and more research is needed 	

Abbreviations: ARR, aortic root replacement; CSF, cerebrospinal fluid; CTA, computed tomography angiography; dSINE, distal stent graft induced new entry; FET, frozen elephant trunk; FEVAR, fenestrated aneurysm repair; IMH, intramural hematoma; PAU, penetrating atherosclerotic ulcer; TAAD, Type A aortic dissection; TAAAR, thoracoabdominal aortic aneurysm repair; TEVAR, thoracic endovascular aortic repair;



Funding

None.

Conflict of interest

The authors declare no conflict of interest related to this article.

Acknowledgments

We are grateful to AORTA and the esteemed Editorial Team for the great collaboration in hosting a Special Issue focused on the 9th International Aortic Summit.

Fig. 1 The 10th International Aortic Summit.