

Postoperative Discomfort Among Laparotomy Patients from a Selected Hospital at Mangaluru: An Observational Study

S. Latha¹ Neetha Kamath² Gincy Joseph³

¹ Department of Medical-Surgical Nursing, Nitte (Deemed to be University), Nitte Usha Institute of Nursing Sciences (NUINS), Mangaluru, Karnataka, India

² Dept of Community Health Nursing, Nitte (Deemed to be University), Nitte Usha Institute of Nursing Sciences (NUINS), Mangaluru, Karnataka, India

³Emergency Nursing and Life Support Course, RGUHS-JeevaRaksha Trust, Bengaluru, Karnataka, India

J Health Allied Sci^{NU} 2023;13:509–517.

Address for correspondence Neetha Kamath, PhD, Nitte (Deemed to be University), Nitte Usha Institute of Nursing Sciences, Department of Community Health Nursing, Mangaluru, Karnataka, India (e-mail: neethakamath@nitte.edu.in).

Abstract Keywords ► humans ► postoperative ► laparotomy ► laparoscopy ► surveys and questionnaires ► pain	 Background Factors causing postoperative discomfort after laparotomy are numerous and must be explored in depth. The postoperative distress may significantly affect the patient's state of well-being. Hence, the present study aims to assess the factors contributing to discomfort after laparotomy. Aim The aim was to assess and compare the discomfort between open and laparoscopic abdominal surgery. Objectives of the Study 1. To analyze the causes of patient's discomfort after abdominal surgery. 2. To compare the discomfort between open and laparoscopic abdominal surgery. 3. To determine the association between pain and selected variables among open and laparoscopic abdominal surgery. Materials and Methods An exploratory approach with a prospective observational design was adopted for this study. Using the purposive sampling technique, 100 patients were selected to open and 100 to laparoscopic abdominal surgery groups. Data were collected using demographic and clinical proforma and standardized postoperative discomfort inventory 6 and 24 hours after the surgery. The assessment focused on the study variables such as symptoms after the surgery and the factors contributing to the postoperative discomfort. Statistical Analysis The collected data were analyzed using descriptive and inferential statistics using the SPSS software version 20. Results Among 200 participants, most of the open (68%) and laparoscopic abdominal surgery is open and 68% in laparoscopic surgery) staved 5 and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In some store) and 10 hours in postoperative (U. In sopen and 68% in laparoscopic surgery) stav
► pain	open and 68% in laparoscopic surgery) stayed 5 and 10 hours in postoperative ICU. In

article published online January 24, 2023 DOI https://doi.org/ 10.1055/s-0042-1760234. ISSN 2582-4287. © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India open and laparoscopic abdominal surgeries, pain is the primary (100%) cause of postoperative discomfort after 6 hours of surgery. Movement restriction is also the reason for significant discomfort both in open (98%) and laparoscopic (100%) abdominal surgeries.

In contrast, pain (99% and 100%), movement restriction (92% and 95%), and abdominal distention (61% and 34%) were the major problems contributing to the patient's discomfort after 24 hours of both open and laparoscopic surgeries too. There was a significant difference in causes of patient discomfort between open and laparoscopic abdominal surgery after 6 hours (P < 0.05), except for discomfort related to IV drip (P = 0.852), constipation (P = 0.2), and chills (P = 0.6).

Conclusion Even though pain is the major distressing factor both in open and laparoscopic surgeries, the current study highlights a few other factors that affect postoperative recovery. Nurses' attention to such distressing factors will fasten patients' recovery and quality of life after abdominal surgery.

Introduction

Recent advances in the medical field have proven surgery as the treatment of choice for many diseases. Surgical procedures in the abdomen are common due to the increasing incidence of problems such as hernia, appendicitis, cholelithiasis, and peptic ulcer disease.Surgery induces a lot of physical and psychological stress on the individual. Traditionally, open surgical techniques are used for the treatment of various diseases. However, laparoscopic surgery is a better option than most open surgeries. Patients undergoing laparoscopic surgery resume activities of daily living, bowel movement, and oral intake earlier than those after open surgery. They had a shorter operative duration and hospitalization stay (8.5 ± 1.9 vs. 11.2 ± 1.8 d, p < 0.05).¹ Laparoscopic colectomy was associated with a shorter average hospital stay and fewer complications, and the length of hospitalization was significantly shorter in the laparoscopic group $(p=0.001)^2$. The risk of complications for patients who underwent laparoscopic colectomy is 2.4% compared to open colectomy, which was 42.9%. Keller et al also reported that the duration of hospital stay was significantly shorter in laparoscopic compared to open abdominal surgery (mean 5.78 vs. 7.80 days, p < 0.0001). The laparoscopic group had significantly lower readmission (5.82 vs. 7.68%, p < 0.0001), complication (32.60 vs. 42.28%, p < 0.0001), and mortality rates (0.52 vs. 1.28%, p < 0.0001)³.

During the postoperative period, patients experience much distress that adversely affects their well-being. Discomfort in the postoperative period is common and may hinder patients' recovery⁴ Patients experience much distress after surgery other than pain. The distress in the postoperative period may affect the patient's well-being and, thus, the postoperative recovery.⁵ Postoperative nausea, vomiting, and pain are most commonly experienced after surgery⁶ Postoperative pain is one of the leading causes of patient discomfort, which reduces patient satisfaction with surgery and postoperative care. In addition to pain, postoperative nausea and vomiting (PONV) are frequent causes of distress and may even extend the stay of patients in the postoperative units. Similarly, movement restriction and dry mouth cause discomfort. Patient comfort is an essential concern in patients receiving surgery; however, the seriousness of discomfort during recovery is unknown.⁷ Therefore, all symptoms causing discomfort should be considered to improve the well-being of patients after abdominal surgery.

As pain is one of the most distressing symptoms, much attention is paid to managing postoperative pain. However, little attention has been focused on other factors contributing to discomfort after the surgery. Besides pain, other symptoms can affect patients' physical well-being during the postoperative period. Alleviating some symptoms and using medical devices only when needed may help reduce the discomfort in the postoperative period and improve the well-being of patients after surgery. With the shift of emphasis from traditional hospitalization to ambulatory surgery and enhanced recovery after surgery, patient safety and comfort are essential to clients' overall well-being.⁸ Nursing practice interventions that prevent the discomfort associated with surgery and help patients effectively adapt to the postoperative situation are cost-effective and can be independently performed by nurses who observe the patient's condition and closely monitor the patient from time to time.⁹

Aim of the Study

The present study aims to identify the causes of patient's discomfort after abdominal surgery and compare the discomfort after open and laparoscopic abdominal surgery.

Materials and Methods

Ethical Consideration

 The Institutional Ethics committee approved the study (NUINS/CON/NU/IEC/2019-20/1481). Written permission was obtained from the authorities, and informed consent was taken from each participant after explaining the patient information sheet in their local language. The researcher also obtained written permission to use the questionnaire for assessing the discomfort after surgerythe postoperative discomfort inventory from the author.⁴

Design and Settings

- This exploratory survey was designed as an observational and cross-sectional study. It was carried out in the selected tertiary care hospital with 1200 beds in Mangaluru. The data were collected from surgery wards, urology wards, gynecology wards, and postoperative ICUs from January 2020 to April 2020. A total of 200 patients were selected after open abdominal and laparoscopic surgeries using the purposive sampling technique based on the prevalence of the previous study. The inclusion criteria for the sampling were patients aged 25 to 65 years admitted to the postoperative ICU in the first 2 hours after surgery and available for assessment at 6 hours and 24 hours after surgery. Patients who needed respiratory support for more than 2 hours and had preexisting severe musculoskeletal problems, and confused patients due to delirium, dementia, psychoses, were excluded from the study. The tools used for the study include demographic proforma, clinical proforma, and a postoperative discomfort inventory. The demographic proforma consisted of seven items, and the clinical proforma had five items. A postoperative discomfort inventory is a standardized tool prepared and validated by Joseph E Banos and consists of 11 items such as pain, nausea, vomiting, movement restriction, sleepiness, dry mouth, insomnia, abdominal distention, constipation, intravenous line-related problems, and chills. A score of one was given if the discomfort was present and zero if there were no discomfort. The degree of discomfort is rated further on a 10-point scale, with the least score of 1 and a maximum of 10, denoting the degree of distress experienced by the patient.
- Eleven experts established the content validity of the instruments. Pretesting of the tools was conducted, and the reliability of the postoperative discomfort inventory was established with Cronbach's alpha value of 0.8. The pilot study was conducted. The tools were administered in the local language Kannada. The baseline demographic proforma and clinical data were assessed. The postoperative discomfort questionnaire was administered to identify the symptoms causing discomfort 6 hours and 24 hours after the surgery. If the distress/discomfort was present, a score of 1 was allotted, and the absence of discomfort was denoted with 0. Those who had discomfort were asked to rate the discomfort further on a 10point scale-the postoperative discomfort inventory with the least score of 1 and a maximum of 10, depending on the degree of distress experienced. The time taken for each participant is approximately 15 minutes twice a day at 6-hourly and 24-hourly assessments.

Results

The collected data were analyzed using descriptive and inferential statistics using the SPSS software version 20. Mann–Whitey U test was used to compare the causes of discomfort at 6 hours and 24 hours after surgery. The association was analyzed using the Chi-square and likelihood ratio. The hypotheses were tested at a 5% significance level, and the p-value < 0.05 was considered significant.

Most participants from open (68%) and laparoscopic abdominal surgeries (42%) were 25 to 35 years of age. The majority, 85% and 52% of the participants, were females. Most of the study population comprised 34% with secondary education in the open surgery group, and 40% of participants had primary and secondary education in the laparoscopic group.

Thirteen percent were hypertensive, and 54% were diabetic. COPD was not elicited among the study subjects. A few participants were smokers (2% from open and 8% from laparoscopic surgery) and alcoholics (5% from the open group and 2% from laparoscopic) among the study groups. The majority, 56%, and 68% stayed 5 to 10 hours in the postoperative ICU.

Frequency Distribution of Clinical Characteristics

In open and laparoscopic abdominal surgeries, most participants (52%) were 150 to 160 cm in height, and 51% and 47% of participants weighed 51 to 60 kg. The respiratory rate was in the normal range of 17 to 24 beats/minute (among 60% of open and 53% of laparoscopic abdominal surgeries). More than half of the subjects had regular bowel movements (52% in open surgery and 61% after laparoscopic surgery). The bladder function was also regular among most participants (69% in the open and 75% in laparoscopic abdominal surgeries).

Assessment of Factors Causing Discomfort after Open and Laparoscopic Abdominal Surgeries

- Table 1 shows that both in open and laparoscopic abdominal surgeries, pain is the most common (100%) cause of postoperative discomfort after 6 hours of surgery. Movement restriction is also the reason for significant discomfort both in open (98%) and laparoscopic (100%) abdominal surgeries. Intravenous line-related problems (81% and 79%) and abdominal distension (77% and 51%) also contribute to postoperative discomfort 6 hours after surgery.

In contrast, pain (99% and 100%), movement restriction (92% and 95%), and abdominal distention (61% and 34%) were again found to be the major problems contributing to patient's discomfort after 24 hours of both open and laparoscopic surgeries too. Intravenous line-related problems show an almost 40% reduction in open abdominal surgeries and a 10% reduction in laparoscopic surgeries compared with 6 hours after surgery.

ltems	6 hours after surgery		24 hours after surgery	
	Open (<i>n</i> = 100)	Laparoscopic (n = 100)	Open (<i>n</i> = 100)	Laparoscopic (n = 100)
	Percentage (%)	Percentage (%)	Percentage (%)	Percentage (%)
Pain				
Yes	100	100	99	99
No	00	00	01	01
Nausea		•	•	•
Yes	51	37	24	19
No	49	63	76	81
Vomiting		•		•
Yes	42	31	16	16
No	58	69	84	84
Movement res	striction			
Yes	98	100	92	95
No	02	0	08	05
Sleepiness				
Yes	59	49	30	28
No	41	51	70	72
Dry mouth				
Yes	56	36	30	09
No	44	64	70	91
Insomnia				
Yes	68	53	40	23
No	32	47	60	77
Abdominal dis	stention			
Yes	77	51	61	34
No	23	49	39	66
Constipation				
Yes	46	38	36	24
No	54	62	64	76
IV-related pro	blems			
Yes	81	79	48	67
No	19	21	52	33
Chills				
Yes	53	58	14	03
No	47	42	86	97

Table 1 Percentage distribution of factors causing discomfort after open and laparoscopic abdominal surgeries

- Table 2 states that distress due to pain after 6 hours of postoperative period among the open abdominal surgery was significantly different from the laparoscopic surgery (p = 0.006). Significant differences were also observed among the other factors such as nausea (p = 0.016),vomiting (p = 0.039),movement restriction (p = 0.029), sleepiness (p = 0.009), dry mouth (p = 0.000), insomnia (p = 0.000) and abdominal distention (median (IQR) (p = 0.000). However there was no change in distress caused by constipation

(p = 0.204), IV-related problems (p = 0.817), and chills (p = 0.611) as these *p*-values were > 0.05.

• Table 3 emphasizes that pain among open abdominal surgeries after 6 hours was higher as the mean difference was 0.80 (95% CI =0.26–1.34), but that is statistically highly significant with a *t*-value = 2.91, p =0.004.

-Table 4 depicts that pain after 24 hours of surgery became almost similar for both groups as the median was 4 and the *p*-value = 0.14. However, discomfort caused by

ltems	Median (IQR)		Mann-Whitney	Z value	<i>p</i> -Value
	Open (<i>n</i> = 100)	Laparoscopic (n = 100)	U test value		
Pain	8 (5-9)	6(5-8)	3899.50	-2.74	0.006*
Nausea	1(0-4)	0(0-3)	4103.50	-2.42	0.016*
Vomiting	0(0-4)	0(0-2)	4271.50	-2.07	0.039*
Movement restriction	4(3-6)	4(3-5)	4123.50	-2.18	0.029*
Sleepiness	2.50 (0-4)	0(0-3)	3993.00	-2.60	0.009*
Dry mouth	1(0-3)	0(0-1)	3612.00	-3.70	0.000*
Insomnia	3(0-5)	2(0-3)	3278.50	-4.37	0.000*
Abdominal distention	2(1-3)	1(0-2)	3216.50	-4.50	0.000*
Constipation	0(0-3)	0(0-2)	4535.00	-1.27	0.204
IV related problems	2(1-3)	2(2-3)	4908.50	-0.23	0.817
Chills	1(0-3)	1(0-2)	4803.00	-0.51	0.611

 Table 2
 Comparison of discomfort after 6 hours of surgery using Mann–Whitney U test

*-significant.

Table 3 Comparison of distress caused by pain among open and Laparoscopic surgery after 6 hours using an independent t-test

Items	$Mean\pmSD$		Mean	t-Value	p-Value	95% CI of the difference (lower to upper)	
	Open (<i>n</i> = 100)	Laparoscopic (n = 100)	Difference				
Distress caused by pain	7.08 ± 2.04	6.28 ± 1.84	0.80	2.91	0.004	0.26 to 1.34	

Table 4	Comparison of	postoperative	discomfort after	24 hours of surgery	/ using Mann–'	Whitey U test
---------	---------------	---------------	------------------	---------------------	----------------	---------------

Patient discomfort	Median (IQR)		Mann–Whitney	Z-Value	<i>p</i> -Value
	Open (<i>n</i> = 100)	Laparoscopic (n = 100)	U test value		
Pain	4 (3-5.75)	4 (3-5)	4418.00	-1.48	0.138
Nausea	0 (0-0)	0 (0-0)	4743.00	-0.87	0.382
Vomiting	0 (0-0)	0 (0-0)	4932.50	-0.26	0.796
Movement restriction	3 (2-3)	2 (2-3)	4068.00	-2.36	0.018
Sleepiness	0(0-1)	0(0-1)	4971.00	-0.09	0.929
Dry mouth	0(0-1)	0(0-0)	3868.50	-4.00	0.000
Insomnia	0(0-2)	0(0-0)	4169.00	-2.47	0.014
Abdominal distention	1(0-2)	0(0-0)	3451.00	-4.12	0.000
Constipation	0(0-1.75)	0(0-0)	4266.00	-2.22	0.027
IV related problems	0(0-2)	1(0-2)	4325.50	-1.74	0.082
Chills	0(0-0)	0(0-0)	4453.50	-2.76	0.006

movement restriction continues to be slightly higher among open abdominal surgeries as median (IQR = 3 [2-3], p = 0.018. Similar effects continued for dry mouth (median [IQR = 0 [0-1], $p \le 0.001$), insomnia (median IQR = 0 [0-2],

P = 0.014) and abdominal distention (median IQR = 1 [0-2], P = 0.000). In contrast, discomfort caused by constipation (p = 0.027) and chills (p = 0.006) become statistically significant after 24 hours of surgery.

Table 5 Comparison of distress caused by pain among open and Laparoscopic surgery groups after 24 hours using an independentt-test

ltems	$Mean \pm SD$		Mean	t-Value	p-Value	95% CI of the	
	Open (<i>n</i> = 100)	Laparoscopic (n = 100)	difference			difference (lower to upper)	
Distress caused by pain	4.53 ± 2.01	3.92±1.01	0.61	2.71	0.007	0.17 to 1.05	

Findings from **- Table 5** explain that the pain among open abdominal surgeries after 24 hours of surgery was higher as the mean difference was 0.61 (95%Cl = 0.17–1.05), with a (*t*-value = 2.71, *p* =0.007); hence a statistically significant difference was found between the groups.

Association between Pain Level with Selected Variables among the Open and Laparoscopic Abdominal Surgery.

The above **- Table 6** depicts that the 'p' values are < 0.05 for the variables hypertension (0.049) and diabetes mellitus (0.017). Hence, there is an association between pain level

in open abdominal surgeries with hypertension and diabetes mellitus.

Similarly, the association between the level of pain with selected variables among the laparoscopic abdominal surgery was also analyzed using chi-square and likelihood ratio, which revealed that the *p*-value for the variable diabetes mellitus is (0.023); as depicted in **– Table 7** hence, there is an association between the level of pain and diabetes mellitus.

Discussion

The present study showed that the major causes of discomfort after surgery was pain, movement restriction, and

Table 6 Association between pain with selected variables using chi-square test and likelihood ratio among open abdominal surgery

Demographic	Level of	pain			Likelihood ratio	<i>p</i> -Value
characteristics (n = 100)	None	Mild (1-3)	Moderate (4-6)	Severe (7-10)		
Age						
25-35	2	28	32	6	9.024	0.435
36-45	0	4	3	0]	
46-55	0	5	3	1	1	
56-65	0	5	6	5	1	
Gender	•					•
Male	0	3	10	2	4.95	0.176
Female	2	39	34	10	1	
Educational status	•					
No formal education	0	4	3	1	15.47	0.217
Primary	2	12	11	5	1	
Secondary	0	19	11	4	1	
Pre-university	0	6	15	2	1	
Graduation and above	0	1	4	0		
Hypertension	0	7	2	4	7.88	0.049
Diabetes Mellitus	2	29	18	5	10.18	0.017
Smoking	0	1	1	0	0.61	0.894
Alcoholism	0	1	4	0	3.44	0.328
Time in postoperative ICU				•	•	
< 5	0	5	11	2	9.16	0.423

Table 6 (Continued)

Demographic	Level of p	oain	Likelihood ratio	<i>p</i> -Value		
characteristics (n = 100)	None	Mild (1-3)	Moderate (4-6)	Severe (7-10)		
5-10	2	27	23	4		
10-15	0	5	5	2		
>15	0	5	5	4		

Table 7	Association	between	the pain	with	selected	variables	using	chi-square	and	likelihood	ratio	among	the	laparosc	opic
abdomina	al surgery														

Demographic	Level of	pain			Likelihood ratio	<i>p</i> -Value
characteristics (n = 100)	None	Mild (1-3)	Moderate (4-6)	Severe (7-10)	1	
Age	•		•			•
25-35	1	23	15	3	8.23	0.511
36-45	1	16	6	2		
46-55	1	9	2	0		
56-65	0	10	10	1		
Gender		•	,			·
Male	0	27	18	3	4.55	0.208
Female	3	31	15	3		
Educational status					•	
No formal education	0	2	1	0	14.86	0.249
Primary	1	21	15	3		
Secondary	1	21	16	2		
Pre University	1	13	0	1		
Graduation & above	0	1	1	0		
Hypertension				•		•
Yes	1	10	2	0	5.04	0.169
No	2	48	31	6		
DM						
Yes	0	4	9	0	9.49	0.023*
No	3	54	24	6		
Smoking						
Yes	0	4	4	0	2.27	0.519
No	3	54	29	6		
Alcoholism						
Yes	0	1	1	0	0.54	0.910
No	3	57	32	6		
Length of stay in postoperat	ive ICU					
< 5	1	6	1	0	8.22	0.513
5-10	2	36	26	4		
10-15	0	7	4	1		
>15	0	9	2	1		

*Significant.

abdominal distention. Similar findings were reported by other researchers (11,12,13,14,15,1,17). The rate of patients with maximal pain scores > 3 was 70% for all patients, 77% for the laparoscopic surgical procedure, and 68% for an open surgical procedure.¹⁰ A prospective observational study on the causes of distress following abdominal surgery showed that the foremost causes of distress at 24 hours were pain (82%), movement restriction (79%) and dry mouth (70%). No significant differences were observed among clients who had undergone open or laparoscopic surgical procedures¹¹ A similar survey on the causes of decreased physical welfare in the postoperative phase among patients, nurses, and physicians discovered that nurses scored pain (8.48) a nasogastric tube (7.13), and nausea (7.10) at the top whereas patients rated pain (5.41), and movement restriction (4.62) as the maximum.¹² Gasparini et al also revealed that the main reason for the patients to avoid phase II procedures was the concern of more severe postoperative discomfort.¹³

In a survey on postoperative distress (other than pain) among 431 clients, 93 (22%) expressed distress. The chief discomfort was the urinary catheter (43%), followed by the throat and gastric tube hurting the nostril (11% each).¹⁴

This agrees with earlier reports that pain and non-pain symptoms contributed 0.084 and 0.074 to the overall discomfort level in the postoperative period. Dry mouth was the most common non-pain symptom in the postoperative period.¹⁵ A cross-sectional study on patient discomfort after abdominal surgery also showed that the perception of discomfort between patients undergoing open surgery and those undergoing laparoscopic surgery; differed significantly. The most common types of discomfort after open surgery were pain (7.1 ± 1.8) , movement restriction (7 ± 1.75) , and dry mouth (6.6 \pm 2.6), whereas, after laparoscopic surgery, the most common types were dry mouth (5.85 ± 2.8) , abdominal bloating (5.3 \pm 2.5), and pain (5 \pm 2.2). There was a relationship between the variable dry mouth and the open surgery group (r=0.40).¹⁶ Further study can be conducted with an intervention to relieve the discomfort after the surgery.

Postoperative pain was not related to gender in the present research. However, MH et al found that women experienced more severe complaints than men after surgery. Also, postoperative pain, nausea, and dry mouth/thirst were symptoms with the highest effective load.¹⁷ The difference could be attributed to a large number of females (85% vs. 52%) in the present study whose perception of symptoms differs from males.¹⁸

Also, 54% of the high-risk group and 13% of the low-risk group had nausea twice or more in the early postoperative period (p < 0.0001).¹⁹ The comparison of postoperative nausea and vomiting after abdominal surgery revealed that the mean score of nausea was 0.68 ± 1.61 versus the vomiting mean score of 0.05 ± 0.22 .⁹ Kenan et al also reported that 20.5% of children experienced discomfort due to nausea and vomiting after dental rehabilitation.²⁰

The risk factors of nausea derived from the study were female gender (p = 0.01) and history of nausea and vomiting

(p = 0.03). Our study found no significant difference in nausea 24 hours after surgery. This could be due to the decreasing side effects of drugs after 24 hours compared to 6 hours. The current research was limited to a small population in a specific hospital area . The study revealed all the causes of discomfort and distress after surgery. Apart from pain all other aspects causing discomfort are usually neglected among postoperative patients. The study revealed a need to focus on all areas of discomfort apart from pain to enhance recovery after laparotomy with fewer complications.²¹ Further study can be conducted with an intervention and a control group to relieve the discomfort.

Conclusion

Pain is the primary cause of discomfort after both open and laparoscopic abdominal surgeries. Restriction of movement and distention of the abdomen also add to discomfort in the postoperative period. While comparing the degree of discomfort, the study findings depict that the patients after laparoscopic surgery were comfortable and at ease. Hence, laparoscopic surgeries can be recommended for patients whenever possible to enhance the recovery from surgery.

Ethical Approval

The study is approved by the Institutional review board approval (NUINS/CON/NU/IEC/2019-20/1481).

Conflict of Interest None declared.

Acknowledgments

The authors are grateful to the authorities of the Nitte Deemed to be University for the financial support for the study. The authors also thank Prof. Josep-E Banos, Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain, for kindly providing us with the postoperative discomfort symptom questionnaire.

Institution to which this study is associated with Nitte Usha Institute of Nursing Sciences, A constituent College of Nitte (Deemed to be University).

References

- 1 Hu MG, Ou-yang CG, Zhao GD, Xu DB, Liu R. Outcomes of open versus laparoscopic procedure for synchronous radical resection of liver metastatic colorectal cancer: a comparative study. Surg Laparosc Endosc Percutan Tech 2012;22(04):364–369
- 2 Kapritsou M, Korkolis DP, Konstantinou EA. Open or laparoscopic surgery for colorectal cancer: a retrospective comparative study. Gastroenterol Nurs 2013;36(01):37–41
- 3 Keller DS, Delaney CP, Hashemi L, Haas EM. A national evaluation of clinical and economic outcomes in open versus laparoscopic colorectal surgery. Surg Endosc 2016;30(10):4220–4228
- 4 Andión Ó, Cañellas M, Baños JE. The postoperative discomfort inventory: a psychometric analysis. J Nurs Educ Pract 2016;6(05)
- 5 Brusciano L, Gambardella C, Terracciano G, et al. Postoperative discomfort and pain in the management of hemorrhoidal disease:

laser hemorrhoidoplasty, a minimal invasive treatment of symptomatic hemorrhoids. Updates Surg 2020;72(03):851–857

- 6 Elsaid RM, Namrouti AS, Samara AM, Sadaqa W, Zyoud SH. Assessment of pain and postoperative nausea and vomiting and their association in the early postoperative period: an observational study from Palestine. BMC Surg 2021;21(01):177
- 7 Fan YL, Qian JL, Ma EL, Stricker PA, Zuo YX. Incidence and risk factors of postoperative severe discomfort after elective surgery under general anesthesia: a prospective observational study. J Perianesth Nurs 2021;36(03):253–261
- 8 Ye H, Chen R, Lian X, et al. Risk factors associated with postoperative pain and discomfort in oculoplastic surgery with general anesthesia: a prospective study. J Pain Res 2018;11:407–415
- 9 Seok Y, Suh EE, Yu SY, Park J, Park H, Lee E. Effectiveness of integrated education to reduce postoperative nausea, vomiting, and dizziness after abdominal surgery under general anesthesia. Int J Environ Res Public Health 2021;18(11):6124
- 10 Allvin R, Rawal N, Johanzon E, Bäckström R. Open versus laparoscopic surgery: Does the surgical technique influence pain outcome? Results from an international registry. Pain Res Treat 2016; 2016:4087325
- 11 Chanif C, Petpichetchian W, Wimo W. Acute postoperative pain of Indonesian patients after abdominal surgery. Nurse Media Journal of Nursing 2012;2(02):409–420
- 12 Andión O, Cañellas M, Baños JEO APhysical well-being in postoperative period: a survey in patients, nurses and physicians. J Clin Nurs 2014;23(9-10):1421–1429

- 13 Gasparini G, Torroni A, Di Nardo F, et al. OSAS surgery and postoperative discomfort: phase I surgery versus phase II surgery. BioMed Res Int 2015;2015:439847
- 14 Zegerman A, Ezri T, Weinbroum AA. Postoperative discomfort (other than pain) - a neglected feature of postanesthesia patient care. J Clin Monit Comput 2008;22(04):279–284
- 15 Tian C, Yu Y, Mao J, Davidson PM. Perceived discomfort, pain and nonpain symptoms in a postanesthesia care unit: an observational study. J Perianesth Nurs 2019;34(05):1032–1039
- 16 Robleda G, Baños JE. Health care professionals' assessment of patient discomfort after abdominal surgery. J Perianesth Nurs 2021;36(05):553–558
- 17 Huppe M, Kemter A, Schmidtke C, Klotz KF. Postoperative complaints: gender differences in expectations, prevalence, and appraisal. Anaesthetist 2013;62(07):528–536
- 18 Amirshahi M, Behnamfar N, Badakhsh M, et al. Prevalence of postoperative nausea and vomiting: A systematic review and meta-analysis. Saudi J Anaesth 2020;14(01):48–56
- 19 Sawatzky JA, Rivet M, Ariano RE, Hiebert B, Arora RC. Postoperative nausea and vomiting in the cardiac surgery population: who is at risk? Heart Lung 2014;43(06):550–554
- 20 Cantekin K, Yildirim MD, Delikan E, Çetin S. Postoperative discomfort of dental rehabilitation under general anesthesia. Pak J Med Sci 2014;30(04):784–788
- 21 Kim TS, Choi SE, Won RS. Undesirable symptoms of patients in the postanesthesia recovery situations. Korean J Anesthesiol 2001;40 (05):619–624