

© () = S

CT-Guided Percutaneous Thrombin Injection of Posttraumatic Aortic Branch Pseudoaneurysm

Aron Michael Devane¹ Christine Marie Gilligan Schammel²

¹Department of Radiology, Prisma Health Upstate, Greenville, South Carolina, United States

²Pathology Associates, Greenville, South Carolina, United States

J Clin Interv Radiol ISVIR 2024;8:49–52.

Abstract

Keywords

- aortic branch pseudoaneurysm
- lumbar artery pseudoaneurysm
- CT-guided thrombin injection

A pseudoaneurysm of an aortic branch artery is a potentially life-threatening uncommon occurrence, which may result in retroperitoneal hemorrhage. Imaging-guided percutaneous thrombin injection has a high technical success and effectiveness rate for treating pseudoaneurysms of the femoral, iliac, and popliteal arteries. We present a case of a retroperitoneal hemorrhage in a patient with a periaortic branch pseudoaneurysm, potentially from an avulsed lumbar artery. Anatomy prevented fixation with a covered stent; the comorbidities eliminated safe surgical repair, and endovascular embolization was not an option due to the avulsion. Computed tomography (CT) fluoroscopy-guided thrombin injection provided a safe, successful embolization with no postprocedural complications.

Greenville, SC 29605, United States

(e-mail: Christine.schammel@prismahealth.org).

Introduction

Posttraumatic aortic branch pseudoaneurysms (PSAs) are uncommon¹ and are typically associated with high-energy, blunt force or penetrating trauma with injury to intraabdominal structures, resulting in a potentially life-threatening retroperitoneal hemorrhage¹ The risk of rupture requires prompt treatment as no reliable predictive symptoms exist.²

The recommended treatment for PSA is catheter angiography with endovascular embolization or endovascular stent graft exclusion. The most common embolic materials are coils; other embolic materials, such as polyvinyl alcohol (PVA), acrylic microspheres, gelatin sponge, and n-BCA are less commonly used.² Imaging-guided percutaneous thrombin injection has been widely accepted³ as a treatment of femoral artery PSAs, with a suggestion for the broader use of thrombin for PSAs across other vessels.⁴

article published online September 3, 2023 DOI https://doi.org/ 10.1055/s-0043-1772493. ISSN 2457-0214. Here, we present a case of a retroperitoneal hemorrhage in an elderly patient presenting with a spinal fracture after a ground-level fall successfully treated with a computed tomography (CT) fluoroscopy-guided thrombin injection. Following institutional review board (IRB) approval and upon receiving publication consent, this report was completed.

Address for correspondence Christine Marie Gilligan Schammel, PhD,

Department of Pathology, USCSOMG, 8 Memorial Medical Ct.,

Case Report

An 81-year-old woman with a history of breast cancer presented to the emergency department with abdominal and back pain after a fall from a standing position. CT examination demonstrated a right periaortic PSA at the T12/L1 level ($2.1 \times 2.1 \times 2.7$ cm; **- Fig. 1**) and a compression fracture of the L1 vertebral body. No apparent communication of the PSA with any branch of the abdominal aorta

^{© 2023.} Indian Society of Vascular and Interventional Radiology. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-ncnd/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



Fig. 1 Contrast-enhanced computed tomography (CT) axial plane demonstrating a pseudoaneurysm (*white arrow*) adjacent to the abdominal aorta.



Fig. 3 Axial computed tomography (CT) angiography (arterial phase) demonstrating resolution of contrast enhancement of the pseudoa-neurysm after thrombin injection (*white arrow*).

tions were observed. At 3.5 months postinjection, the patient was asymptomatic, and a CTA illustrated a successful embolization.

was noted, suggesting an avulsed origin of a lumbar artery or the right subcostal artery given the T12/L1 location, eliminating the potential for catheter angiography and coil embolization. The adjacent aortic branches, such as the celiac axis, made fixation with a covered stent technically challenging and, as the spinal artery arises from this location, induced an increased risk of paraplegia. The location of the PSA and the patient's comorbidities made surgical repair high risk.

Utilizing CT fluoroscopy guidance, a 10-cm 19-gauge needle (Argon Medical, Plano, TX) was advanced into the PSA with a prone right paravertebral approach (**-Fig. 2**). Pulsatile arterial blood emanated from the needle spontaneously. Reconstituted thrombin (0.5 mL; 500 units; Pfizer, New York, NY) was administered with cessation of blood from the needle. No blood could be manually aspirated with a 5-mL syringe after a 10- to 15-second wait period. The needle was removed. CT angiography (CTA) demonstrated a lack of contrast opacification of the PSA in the arterial (**-Fig. 3**) and the delayed phase of intravenous contrast enhancement. No procedural-related complica-



Fig. 2 Computed tomography (CT) fluoroscopy image demonstrating the right paravertebral approach of a 19-gauge needle into the pseudoaneurysm.

Discussion

While aortic branch PSAs are typically associated with blunt abdominal trauma with a spinal injury,¹ in an elderly patient, this should be considered a source of hemorrhage, even from a ground-level fall. On imaging, aortic branch PSAs present as an enhanced density close to the psoas major muscle, the lumen enhancing similar to that of the aorta.^{1,5} A literature review of 26 cases (>Table 1) noted that all reports but two used embolization. Our report notes the limitations of the anatomy and lack of PSA communication with any abdominal aortic branch, suggesting an avulsed origin. This makes it impossible to catheterize the visceral aortic branch PSA endovascularly, and coil embolization was not feasible. The use of other embolic agents, such as glue or Onyx, would have required visualization, which was not a possibility in our case. Also, these embolic agents typically require larger volumes, risking reflux; glue would have been an unsafe option due to the possibility of attaching the needle to the PSA, risking a tear. While direct percutaneous injection of thrombin has been the preferred treatment for postcatheterization femoral artery PSAs,^{2,3} it has had limited use in aortic branch PSA, except when embolization is not an option.^{2,5} Given these data, thrombin appeared to be the safest option for this patient. A small amount of thrombin was administered initially to prevent reflux into the abdominal aorta, which could have resulted in visceral and lower extremity thrombus. Preparations were made for repeated small injections with repeated contrast-enhanced CT imaging if the first dose was ineffective; fortunately, this was unnecessary.

Given the favorable outcomes at other sites and that aortic PSAs are life-threatening, thrombin is a viable option for this site.

Compliance with Ethical Standards

All procedures performed in studies involving human participants were in accordance with the ethical standards of

Study	Age	Cause	Presentation	Resolution
Ikubo et al ⁶	62	Blunt trauma	Abdominal pain/hypotension	Embolization
Siablis et al ⁷	45	Fall from 6 m	"Agonizing pain"	Embolization
Chan and Korivi ⁸	24	Gunshot	Low back/flank pain	Embolization
Kessel et al ⁹	21	Penetrating abdominal trauma	Missed during laparotomy	Embolization
Lee et al ¹⁰	47	Fracture of lumbar spine	Hypotensive shock	Embolization
Domenicucci et al ¹¹	23	Thoracolumbar vertebral dislocation	Anemia/hemorrhage	Embolization
Ramsay and Marshall ^{12,a}	23	Renal biopsy		Thrombin
Sharma et al ^{5,a}	47	Spontaneous	Abdominal pain	Thrombin
Tomescot et al ¹³	79	Osteomyelitis	Acute back pain	Embolization
Oh et al ¹⁴	55	Lumbar posterolateral fusion	Retroperitoneal Hemorrhage	Embolization
Counihan et al ¹⁵	18	Knife stab	Bleeding	Embolization
Suresh et al ¹⁶	60	Postpercutaneous coronary intervention	Back ache/hypotensive	Embolization
Tsekouras et al ¹⁷	64	Inferior vena cava filter	Abdominal pain	Embolization
Djuimo et al ¹⁸	21	Nephrostomy removal	Bleeding on removal	Embolization
Giordano et al ¹⁹	73	Cryoablation/vertebroplasty lumbar met	Abdominal pain	Embolization
Vashisht et al ⁴	32	Lumbar/spine fracture from fall	Hypotensive shock	Symptom management
Salam and Khandwala ²⁰	71	Renal biopsy	Hemodynamic instability	Embolization
Panagiotopoulos et al ²¹	39	Lumbar diskectomy	Abdominal pain	Embolization
Latka et al ²²	58	Postlumbar fusion	Abdominal pain	Embolization
Méndez et al ²³	72	Postlumbar fusion	Monoplegia abdominal pain	Embolization
Eissa et al ²⁴	30	Fall from 2 m/postlumbar fusion	Back pain	Embolization
Ruffilli et al ²⁵	53	Postlumbar fusion	Back pain	Embolization
Keerthivasan et al ²⁶	80	Postlumbar fusion	Leg pain	Embolization
Kim et al ²⁷	73	Postlumbar fusion	Swelling of leg	Stent graft
Nam et al ²⁸	62	Postvertebral augmentation	Back pain	Embolization
Nam et al ²⁸	88	Postvertebral augmentation	Back pain	No intervention

Table I Survey of reported lumbar aftery pseudoaneurys

Note: The table provides a survey of reported lumbar artery pseudoaneurysms, as this was the most likely location in our patient. ^aThe only report of thrombin injections for an aortic branch (lumbar artery) pseudoaneurysm.

the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

This study has obtained IRB approval from (Prisma Health), and the need for informed consent was waived. Dr. Devane is a paid speaker for Johnson and Johnson and TriSalus Life Sciences and is a consultant with Boston Scientific and Guerbet.

Funding None.

Conflict of Interest None declared.

References

- 1 Shigematsu Y, Kudoh K, Nakasone Y, Fujisaki T, Uemura S, Yamashita Y. Nontraumatic rupture of lumbar artery causing an intravertebral body pseudoaneurysm: treatment by transcatheter embolization. Cardiovasc Intervent Radiol 2006;29(05):870–874
- 2 Thomson B, Patel V, Moser S, Diamantopoulos A. Successful Treatment of a left gastric artery pseudoaneurysm by image guided percutaneous thrombin injection. Vasc Endovascular Surg 2023;57(03):306–310
- 3 Ehieli WL, Bozdogan E, Janas G, et al. Imaging-guided percutaneous thrombin injection for the treatment of iatrogenic femoral artery pseudoaneurysms. Abdom Radiol (NY) 2019;44(03): 1120–1126
- 4 Vashisht S, Bachhal V, Kumar V, Kumar D. Lumbar artery pseudoaneurysm following blunt trauma: A case report and literature review. Trauma Case Rep 2019;23:100232

- 5 Sharma D, Elmore JRE, Nadal LLS, Sheldon DGI, Franklin DPI. Treatment of spontaneous lumbar artery pseudoaneurysm by computed tomography: guided thrombin injection. Vasc Endovascular Surg 2008;42(03):284–288
- 6 Ikubo A, Komura M, Matoba N, et al. Lumbar artery pseudoaneurysm: an unusual cause of a retroperitoneal hematoma: report of a case. Surg Today 1993;23(07):635–638
- 7 Siablis D, Panagopoulos C, Karamessini M, et al. Delayed diagnosis of a false aneurysm after lumbar arterial injury: treatment with endovascular embolization: a case report. Spine 2003;28(04): E71–E73
- 8 Chan KT, Korivi N. Lumbar artery pseudoaneurysm in traumatic spinal cord injury: a case report. Arch Phys Med Rehabil 2003;84 (03):455–457
- 9 Kessel BJ, Habib FA, Thompson B, Bajayo DE, Shatz D. Lumbar artery pseudoaneurysm: an unusual complication of penetrating abdominal trauma. Eur J Trauma 2004;30:187–190
- 10 Lee JS, Kim CW, Suh KT. Lumbar artery injury combined with a transverse process fracture of the lumbar spine presenting with hypovolemic shock after a fall: a case report. J Korean Orthop Assoc 2008;43:400–403
- 11 Domenicucci M, Ramieri A, Lenzi J, Fontana E, Martini S. Pseudoaneurysm of a lumbar artery after flexion-distraction injury of the thoraco-lumbar spine and surgical realignment: rupture treated by endovascular embolization. Spine 2008;33(03):E81–E84
- 12 Ramsay DW, Marshall M. Lumbar artery pseudoaneurysm following renal biopsy: treatment with ultrasound-guided thrombin injection. Australas Radiol 2002;46(02):201–203
- 13 Tomescot A, Dallaudiere B, Zurlinden O, Manelfe J. Lumbar artery pseudoaneurysm as a late complication of osteomyelitis with vertebral body destruction. J Vasc Surg 2013;58(04):1084–1087
- 14 Oh YM, Choi HY, Eun JP. Delayed retroperitoneal hemorrhage due to lumbar artery pseudoaneurysm after lumbar posterolateral fusion. J Korean Neurosurg Soc 2013;54(04):344–346
- 15 Counihan M, Pontell ME, Selvan B, Trebelev A, Nunez A. Delayed presentation of a lumbar artery pseudoaneurysm resulting from isolated penetrating trauma. J Surg Case Rep 2015;2015(07):rjv083
- 16 Suresh A, Seshagiri Rao D, Rammurti S, Srinivas B, Sivaprasad A. Coil closure of lumbar artery pseudoaneurysm: an unusual complication associated with anomalous left circumflex artery primary percutaneous coronary intervention. Chronic Dis Transl Med 2015;1(01):55–58

- 17 Tsekouras N, Whalen RC, Comerota AJ. Lumbar artery pseudoaneurysm in a patient with inferior vena cava filter and history of strenuous physical exercise. J Vasc Surg 2015;61(03):796–799
- 18 Djuimo M, Aubé M, Beland M, Jeldres C, Carmel M, Benko A. Lumbar artery pseudoaneurysm: a complication of percutaneous nephrostomy. Urol Case Rep 2017;13:66–68
- 19 Giordano AV, Arrigoni F, Bruno F, et al. Interventional radiology management of a ruptured lumbar artery pseudoaneurysm after cryoablation and vertebroplasty of a lumbar metastasis. Cardiovasc Intervent Radiol 2017;40(05):776–779
- 20 Salam B, Khandwala K. Lumbar artery pseudoaneurysm following renal biopsy. Cureus 2018;10(05):e2634
- 21 Panagiotopoulos K, Gazzeri R, Bruni A, Agrillo U. Pseudoaneurysm of a segmental lumbar artery following a full-endoscopic transforaminal lumbar discectomy: a rare approach-related complication. Acta Neurochir (Wien) 2019;161(05):907–910
- 22 Latka K, Zurawel R, Maj B, Olbrycht T, Chowaniec J, Latka D. latrogenic lumbar artery pseudoaneurysm after lumbar transpedicular fixation: case report. SAGE Open Med Case Rep 2019;7: X19835344
- 23 Méndez JC, Fandino E, Bermúdez-Coronel I, Prieto MA, Blázquez J. Endovascular occlusion of iatrogenic lumbar artery pseudoaneurysm using liquid embolic agent: case report. Trauma Case Rep 2020;29:100352
- 24 Eissa AT, Shahbaz A, Alhelal F, et al. Postoperative paralumbar pseudoaneurysm: a case report. J Neurol Surg Rep 2020;81(02): e33-e37
- 25 Ruffilli A, Barile F, Fiore M, Pasini S, Facchini G, Faldini C. Occurrence of L4 lumbar artery pseudoaneurysm after posterior reduction and L4-L5 fusion for grade I anterolisthesis: a case report. JBJS Case Connect 2020;10(04):00117
- 26 Keerthivasan P, Anupama NV, Kanna RM, Shetty AP, Rajasekaran S. Lumbar artery pseudoaneurysm: a rare case of delayed onset incomplete cauda equina syndrome following transforaminal lumbar interbody fusion. Eur Spine J 2020;29(Suppl 2):156–161
- 27 Kim C, Hwang D, Yun WS. Endovascular repair of an ilio-iliac arteriovenous fistula with pseudoaneurysm after lumbar disc surgery: a case report. Vasc Spec Int 2021;37:30
- 28 Nam D, Johnson MH, Mojibian H, Hebert RM, Cornman-Homonoff J. Development of paravertebral pseudoaneurysms following vertebral augmentation: a report of two cases. Clin Imaging 2022;90:39–43