





# Fracture-Related Infection – What Does the Literature Tell Us?

## *Infección asociada a fractura ¿Qué nos dice la literatura?*

Tomás Pineda<sup>1</sup>  Andrés Fuentealba<sup>2</sup>  Álvaro Zamorano<sup>3,4</sup> 

<sup>1</sup> El Carmen Hospital, Santiago, Chile

<sup>2</sup> Emergency and Public Assistance Hospital, Santiago, Chile

<sup>3</sup> University of Chile Clinical Hospital, Santiago, Chile

<sup>4</sup> Department of Traumatology and Orthopedics, Mutual de Seguridad, Santiago, Chile

Address for correspondence Tomás Pineda, MD, El Carmen Hospital, Avenida Camino a Rinconada 1201, Maipú, Santiago, Chile (e-mail: tpinedarojas@gmail.com).

Rev Chil Ortop Traumatol 2024;65(1):e23–e33.

### Abstract

Fracture-related infection (FRI) is one of the most frequent and challenging complications of orthopedic trauma; however, its importance has been underestimated. Historically, there has been a lack of standardization in its management. Nevertheless, the available scientific evidence has increased in recent years, given multiple clinical guidelines and expert consensus. This review aims to provide an update for orthopedic trauma surgeons to standardize diagnostic and treatment criteria based on recent scientific evidence.

### Keywords

- ▶ fracture
- ▶ infection
- ▶ diagnosis
- ▶ treatment

### Resumen

La infección asociada a fracturas (IAF) es una de las complicaciones más frecuentes y desafiantes del trauma ortopédico, sin embargo, su importancia ha sido subestimada existiendo históricamente una falta de estandarización en su manejo. En los últimos años la evidencia científica disponible ha ido en aumento, y a consecuencia de ello múltiples guías clínicas y consensos de expertos han sido publicados.

### Palabras clave

- ▶ infección
- ▶ fractura
- ▶ diagnóstico
- ▶ tratamiento

El objetivo de este trabajo es proporcionar una actualización, dirigida principalmente a especialistas en Ortopedia y Traumatología, buscado estandarizar criterios diagnósticos y de tratamiento basado en evidencia científica reciente.

## Introduction

Fracture-related infection (FRI) is one of the most significant complications of orthopedic trauma. It has substantial differences compared to other infectious conditions. The presence of osteosynthesis material to provide the required stability for bone consolidation makes FRI a challenge from its diagnosis to treatment.

Despite advances in prevention measures in recent decades, the incidence of this complication remains high,<sup>1,2</sup> and reported successful treatment rates range from only 70 to 90%.<sup>3,4</sup> FRI results in critical sequelae and represents a considerable global socioeconomic problem.<sup>5,6</sup> However, the real impact of this complication is difficult to estimate due to the lack of a clear definition and diagnostic criteria standardization. In a recent systematic review, Metsemakers

received

June 14, 2022

accepted

November 27, 2023

DOI <https://doi.org/10.1055/s-0043-177750>.

ISSN 0716-4548.

© 2024. Sociedad Chilena de Ortopedia y Traumatología. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

et al. demonstrated significant variability in the criteria for FRI definition. Metsemakers et al. reported that only 2% of the total randomized controlled studies published to date used a validated definition for FRI diagnosis, 27% used an arbitrary definition created by the authors themselves, and the remaining works did not present any definition.<sup>7</sup>

The lack of rigor in FRI diagnosis has made it difficult to evaluate any result and properly compare different treatments when giving recommendations with a high evidence level. As such, recent years saw an increase in the number of scientific publications on this topic, highlighting several consensus and expert recommendations.<sup>8-12</sup>

### Definition

In 2018, a group of experts from different parts of the world published, based on current evidence, a proposed definition of FRI using a methodology similar to that described by Cats-Baril et al. for periprosthetic infection (PPI).<sup>8,13</sup> Their goal was to provide a tool allowing clinical study standardization and improving the available evidence quality.

This consensus unanimously decided there should be a single definition of FRI without subdivisions regarding time (acute or chronic infection), anatomical location, or infection depth (superficial or deep). This decision has two reasons. First, a potential subdivision would make such a definition unnecessarily complex and difficult to use in daily practice. Second, the classifications described in the literature to date are based mainly on progression time and define deadlines arbitrarily.<sup>14,15</sup>

Infections are acute or chronic, potentially requiring different treatment strategies. However, this should not affect how physicians define FRI.<sup>8</sup>

On the other hand, experts recognize superficial infections may not have any relation with the fracture or the implant. Nonetheless, they understand that the definition of

the superficial nature of an infection can only occur retrospectively and cannot guide the treatment.<sup>8</sup> Therefore, for definition (and data collection), surgeons must determine the presence of infection, not its extent, location, or classification.<sup>8,16</sup>

### Diagnosis

Two systematic reviews analyzing clinical, laboratory, and imaging features from FRI based on which criteria to determine its diagnosis.<sup>7,17</sup>

For Metsemakers et al., some of these criteria may provide definitive proof of infection, while other less-specific criteria can suggest the diagnosis and occur in patients with no infection. This realization resulted in a set of confirmatory and suggestive criteria<sup>8</sup> (►Table 1).

Recently, Onsea et al. evaluated the performance of these diagnostic criteria in a multicenter retrospective cohort study, demonstrating excellent outcomes. These authors detected a sensitivity and specificity of 97.5% and 100%, respectively, for the presence of a confirmatory criterion and a 95% specificity for the suggestive clinical criteria (fever, exudation, local heat, and redness), reaffirming the importance of the physical examination for FRI diagnosis.<sup>18</sup>

It is worth mentioning that, in multiple aspects, solid scientific evidence is scarce; therefore, many criteria are based on expert opinion.<sup>8,17</sup> Scientific evidence on histopathological diagnosis, for instance, is limited.<sup>19</sup> In contrast to the PPI definition, the recently published consensus expert panel did not include the presence of an inflammatory cell infiltrate on histopathological examination (i.e., polymorphonuclear cell count) because there is no clear scientific evidence on a cut-off value for reliable diagnosis determination. Today, there is still no standardized and reproducible protocol for the evaluation of histopathological samples obtained during surgery for FRI.<sup>8,20</sup>

**Table 1** Diagnostic Criteria

Confirmatory	Suggestive
1. Operative fistula or dehiscence (osteosynthesis material or bone exposure)	1. Clinical: Pain (non-weight bearing, increasing over time, or recent onset), local heat or redness, local swelling, local increase in temperature, fever.
2. Purulent wound discharge or presence of pus during surgery	2. Radiological: Osteolysis (at the fracture site or near the implant), implant loosening, sequestration, lack of consolidation progress, periosteum formation (not in fracture focus or a consolidated fracture)
3. Pathogen identification in at least two deep samples (implant-bone interface) cultured separately (also consider samples obtained by sonication)	3. Pathogen identification in a single deep culture sample
4. Detection of organisms in deep tissues using special staining techniques (tuberculosis, fungi, etc.)	4. Laboratory: Elevation of serum inflammatory markers (white blood cell count, erythrocyte sedimentation rate [ESR], C-reactive protein [PCR] levels)
	5. Persistent, increasing, or new discharge beyond the first postoperative days with no valid alternative explanation
	6. New-onset joint effusion in patients with fracture (remember bone and joint infections can present as adjacent septic arthritis [prosthesis penetrating the joint capsule during application or intra-articular fractures])

On the other hand, a recent systematic review and meta-analysis assessed the diagnostic value of different serum inflammatory markers (C-reactive protein levels, white blood cell count, and erythrocyte sedimentation rate) in suspected chronic or late-onset FRI. This meta-analysis demonstrated a limited diagnostic value for these tests, concluding that they are insufficient to confirm or rule out a chronic or late-onset FRI, therefore considering them suggestive criteria alone.<sup>21</sup>

Imaging studies have three basic indications for greater certainty in FRI diagnosis, including extension, the presence of sequestration or abscesses, and the degree of fracture consolidation and implant stability.<sup>11</sup> Depending on local preference and availability, these techniques range from conventional radiography, computed tomography (CT), or magnetic resonance imaging (MRI) to more complex techniques with lower availability, such as three-phase bone scintigraphy (TPBS), positron emission tomography (PET), and radiolabeled white blood cell scintigraphy (WBCS).

For conventional radiography, a widely available and low-cost test, there are no studies of good methodological quality to evaluate its diagnostic usefulness; however, it provides basic information regarding fracture consolidation and implant stability.

CT has 47% sensitivity and 60% specificity; as such, it is a useful test for determining sequestration or bone cavitation. CT also allows observing implant loosening, osteolysis signs, and nonunion.<sup>8,22,23</sup>

MRI helps in the evaluation of soft tissues. In addition, it is very sensitive in detecting morphological bone changes, allowing the evaluation of the extent of bone and soft tissue involvement, and the presence of sequestration, sinus tracts, and/or subcortical abscesses. However, the differentiation between infection and inflammation-related changes and normal tissue healing may be difficult, and the effect generated by metal implants can affect image quality. MRI sensitivity and specificity reported for FRI range from 82% to 100% and 43% to 60%, respectively.<sup>22–24</sup>

Nuclear medicine studies described in the literature include TPBS, PET, and WBCS. In recent years, hybrid images (single photon emission CT [SPECT]/CT, PET/CT, PET/MRI) became available, improving the results.<sup>22</sup>

TPBS has a high sensitivity (89 to 100%) but a very low specificity (0 to 10%); as a result, it is not recommended for the FRI treatment.<sup>24–26</sup>

WBCS plus SPECT have high sensitivity (79 to 100%) and specificity (89 to 97%).<sup>22,27,28</sup> As a benefit, recent surgeries do not influence them.<sup>28</sup>

Although PET presents a slightly lower sensitivity (65 to 94%) and specificity (76 to 100%), it should not be used for FRI diagnosis within the first month after surgery.<sup>29</sup>

Despite the previously described usefulness of imaging studies, there is not enough scientific evidence to determine whether any of them can be categorical for FRI diagnosis, and this is why they are currently considered suggestive criteria.<sup>8,11</sup>

Determination of the causative pathogen, the cornerstone of treatment, is usually achieved through intraoperative culture. Sample collection must be careful since false-positive or false-negative results can lead to erroneous treatment decisions, compromising the outcomes. Today, FRI diagnosis uses sampling protocols validated for PPI management and a series of recommendations:<sup>8,11,12</sup>

– Avoid antibiotic therapy for at least two weeks before sample collection.

- Take at least five deep tissue or fluid samples, ideally from the interface between the bone and the implant.
- Cultivate all samples separately
- Take and manipulate each sample using different surgical instrumentation to avoid cross-contamination.
- Avoid taking samples from the skin or fistulous tracts.
- Avoid using swabs for sample collection

There is still controversy in the literature regarding the culturing time. Although a 7 to 14-day period is reasonable, the final decision must consider the local reality. As such, adequate coordination between the treating surgeon and the microbiology team at each center is essential to balance the risk of losing a pathogen difficult to culture versus obtaining a culture contaminated with an irrelevant pathogen.<sup>30</sup>

At the same time, the recommendation is to consider using specific cultures for mycobacteria or fungi based on local environmental and epidemiological risk factors.<sup>31</sup>

## Treatment

### Treatment Team Composition

The management of FRI patients represents a challenge for the entire medical team. Currently, the evidence favoring multidisciplinary FRI management is increasing.<sup>1,32</sup> The team composition relies on the requirements of each patient. However, the recommendation is to have at least members from the traumatology and infectious disease team. In addition, many cases require support from other specialties, such as plastic surgery, anesthesiology, internal medicine, and nutrition. It is advisable to refer the patient to a more complex center if the institution does not have the proper equipment to meet their needs.<sup>1</sup>

### Preoperative Optimization

It is essential to identify and correct nutritional and metabolic disorders in FRI patients, including malnutrition and diabetes mellitus, since these conditions can compromise the outcomes and increase the risk of complications, hospitalization time, and costs.<sup>33–35</sup> In this context, several clinical guidelines recommend preoperative optimization and even intraoperative monitoring of blood glucose levels to maintain them between 140 mg/dL and 180 mg/dL throughout the surgery.<sup>36,37</sup>

Hypovitaminosis D is a common condition among trauma patients and a presumed fundamental factor for bone consolidation and infection prevention. However, there is not enough evidence to reliably recommend its routine screening and supplementation.<sup>38,39</sup>

### Treatment Strategy

There are two main strategies regarding surgical treatment, and their choice depends on the context. The first alternative

consists of implant retention combined with debridement and antibiotic therapy. The second alternative is implant removal or replacement associated with debridement and antibiotic therapy.

The importance of the present osteosynthesis material lies in the bacterial colonization and biofilm formation ability, allowing the development of local antibiotic resistance and protection against the immune system. These factors let organisms survive antibiotic doses up to 1,000 times higher than in their planktonic state.<sup>40</sup>

Although some concepts described for PPI were a base to establish the diagnostic criteria for FRI, one of the fundamental differences between these conditions is the potential implant removal after fracture consolidation on FRI, with a high probability of infection eradication. This is why complete infection eradication is not always the initial objective, and the surgeon may opt for suppressive management while maintaining an implant as long as it provides adequate stability until bone consolidation.<sup>41</sup>

In recent decades, several published preclinical studies demonstrated the importance of fracture stability in FRI management. The first authors to describe this importance were Ritmann and Perren, who conducted a study on sheep and reported that consolidation is possible during an infection provided the implant is absolutely or relatively stable.<sup>42</sup>

Similarly, different works reaffirmed the importance of this variable not only for bone consolidation but also for infection prevention and eradication.<sup>43–45</sup> The pathophysiology of the lower susceptibility of a stabilized fracture to infection remains unclear. It is difficult to determine whether instability is the cause or consequence of the infection. However, what is currently most accepted is that both act simultaneously, with positive feedback between them.<sup>45</sup>

Considering all the above, Metsemakers et al. proposed the following therapeutic goals for FRI:<sup>41</sup>

1. Fracture consolidation
2. Infection control (eradication versus suppression until fracture consolidation)
3. Soft tissue coverage
4. Prevention of chronic osteomyelitis
5. Functional recovery

Time is a relevant factor in deciding on implant removal or retention since it directly correlates with the biofilm maturity and degree of bone consolidation. It is postulated that in the presence of an immature biofilm, cleansing, debridement, and antibiotic therapy may eradicate the infection without the need to remove the implant.<sup>46</sup>

Morgenstern et al. recently published a systematic review evaluating outcomes of implant retention and its correlation with time. These authors observed good success rates (86% to 100%) based on the absence of recurrent infection and implant retention in infections occurring for less than 3 weeks. In longer infections (with 3 to 10 weeks of duration), the success rates were lower (82 to 89%), decreasing considerably (67%) in cases of more than 10 weeks of progression, with limited literature available for this last scenario.

However, the authors concluded that despite the importance of this variable, it should not be the only factor to consider when deciding whether to retain or remove an implant.<sup>46</sup>

Other factors that must be considered when deciding between implant removal or replacement are its stability, whether the reduction obtained is acceptable or not, the patient's baseline medical condition in cases of very severe infections, and the presence of an intramedullary implant whose maintenance can make surgical cleanliness difficult. Based on this last point, there is consensus on the removal of the intramedullary implant as part of FRI treatment after fracture consolidation.<sup>12,41</sup>

Tschudin-Sutter et al. propose a management algorithm considering strict criteria for implant retention. These authors obtained a 90% success rate in a series of 122 patients followed up for 2 years. The inclusion criteria used were symptoms for less than 3 weeks, stable implant, absence of abscess or fistula, identified pathogen, and sensitivity of the pathogen to an antibiotic agent active against the biofilm.<sup>4</sup>

In conclusion, the choice of treatment strategy is a complex process. It is critical to perform a complete study of each patient and consider the multiple variables involved, including time, which is a relevant parameter but not the only factor (► Fig. 1).


### Surgical Cleansing and Debridement

Irrigation removes bacterial load and debris by dragging, resulting in macroscopically clean tissues. Today, the best quality evidence on irrigation is extrapolated from open fracture treatment. FLOW, a multicenter randomized study, showed the usefulness of irrigation with saline or lactated Ringer's solution at low pressure, not requiring detergents.<sup>47</sup> Although additional antimicrobial solutions could reduce infection rates, this topic remains controversial since *in vitro* studies demonstrated that some have a cytotoxic effect on osteoblasts.<sup>48–51</sup>

Debridement, a key element in treatment, consists of resecting all necrotic and contaminated tissues, including bone tissues. We must consider that periostized bone remains susceptible to maintenance, while non-periostized bone fragments are avascular and require removal. It is still debatable how to define a critical bone defect requiring additional surgical interventions and what to do with large non-periostized structural fragments. Nevertheless, following their removal, the next treatment strategy needs consideration.<sup>52–54</sup>

Ideally, we must fill this critical bone defect with vital tissue to stimulate neovascularization, allowing the arrival of systemic antibiotics and immune cells. The use of local antibiotics associated with polymethylmethacrylate (PMMA), as described by Masquelet, is a good alternative for these cases because it manages the defect and allows antibiotic release for a longer time.<sup>12,41,55,56</sup>

Finally, in recent years, bioactive glass has gained interest. Bioactive glass is a biocompatible synthetic material with antibacterial, osteoconductive, and angiogenic properties and promising results in managing bone defects in infections.<sup>57,58</sup>



**Fig. 1** Flow chart representing the optimal treatment course for a patient with fracture-related infection (FRI). Adapted from Metsemakers et al.<sup>12</sup>

### Antibiotic Therapy

Identifying the etiological agent is essential throughout the treatment. This explains why antibiotic therapy before intraoperative culture sample collection should be reserved only for seriously ill patients with systemic involvement or local extremity involvement when empirical treatment is imperative.<sup>9</sup>

In general terms, patients with FRI must undergo inflammatory parameter tests and blood cultures before therapy, as well as follow-up parameters depending on the antibiotic agent (complete blood count, kidney function, liver function, etc.). These tests allow the determination of baseline values and the design of a therapy monitoring curve.<sup>12</sup>

Empirical intravenous antibiotic treatment should start soon after sample collection and debridement to reduce the local bacterial load and the risk of antibiotic resistance.<sup>9,10</sup> Initially, treatment must include broad-spectrum agents per the local reality. In general, it should include a lipopeptide or a glycopeptide and an agent covering gram-negative bacilli. This initial treatment must be modified per the culture and sensitivity results as soon as possible<sup>9</sup> (► **Table 2**).

Although treatment duration remains controversial, the current trend is to reduce the time of intravenous therapy. A recent randomized controlled trial demonstrated that patients treated for up to 7 days with intravenous antibiotic therapy followed by oral therapy had the same outcomes as those with prolonged intravenous therapy.<sup>59</sup>

The curative approach of antibiotic therapy in case of implant retention is only successful when the antibiotic agent is effective against the biofilm. Rifampicin is the antibiotic of choice against most gram-positive bacteria,<sup>60-62</sup> while fluoroquinolones are recommended for gram-negative bacteria.<sup>63,64</sup> Even so, it is worth mentioning that rifampicin must always be combined with a second antibiotic agent due to the rapid development of resistance. For the same reason, its administration should start before the initial reduction in the bacterial load through surgery and antibiotic therapy, removal of all drains, and the presence of a closed, dry wound.<sup>65,66</sup> For staphylococci, quinolones, such as ciprofloxacin or levofloxacin, are the most studied and effective oral antibiotics in association with rifampicin.<sup>67</sup>

For bacteria resistant to antibiotics active against the biofilm, the eradication alternative by retaining the implant is not viable, so the surgeon must seriously consider its removal for proper treatment.<sup>68</sup>

Follow-up duration for FRI patients must extend for a minimum of 12 months after cessation of antibiotic therapy due to the risk of recurrence.<sup>9</sup>

### Negative Culture

Culture-negative FRI represents a major challenge for the multidisciplinary treating team. The estimated rate of FRI with negative cultures ranges from 1% to 16%. However, the incidence could be even higher, especially in cases of nonunion.<sup>69-73</sup>

The causes of this phenomenon are variable (low number of cultures, inadequate sample collection site, poor sample handling, fastidious organisms, etc.). In these cases, depending on each institution's availability, additional methods to increase organism detection, including implant sonication or molecular techniques, should be considered. If the causative agent is not identifiable, one must choose the treatment targeting the most likely organism.

On the other hand, if the diagnosis of FRI is not confirmed and cultures are negative, the current recommendation is to suspend empirical antibiotic treatment to observe the clinical progression, and eventually repeat tissue sampling, which avoids antibiotic toxicity and the risk of antibiotic resistance.<sup>9</sup>

### Local Antibiotics

The application of local antibiotics is a critical complement to FRI treatment, especially in the presence of a bone defect.<sup>10</sup> A recent systematic review showed a considerable reduction in the risk of infection in open fractures with local antibiotic administration, mostly with polymethylmethacrylate (PMMA) as a distribution form.<sup>55</sup>

Outside of the commercially available preparations of PMMA with antibiotics, antibiotic agent addition is a valid alternative considering the resulting structural stability. To do this, Metsemakers et al. recommended adding up to 10% of the weight of the cement bag, respecting the maximum doses of each antibiotic agent.<sup>10</sup>

Gentamicin and vancomycin are the most commonly used antibiotics in our institution. Nonetheless, there is no solid evidence today to recommend the maximum antibiotic dose for PMMA addition. In a recent article from the consensus FRI group, Metsemakers et al. described, based on the available literature, a dose of 4.8 g for gentamicin and 6 g for vancomycin, recommending 4 g of vancomycin per PMMA bag (40 g).<sup>10</sup>

Other antibiotic agents less frequently used in FRI, including clindamycin, colistin, amphotericin B, cefazolin, and ampicillin, have been described as alternatives.<sup>10</sup> Some drugs are not recommended, such as  $\beta$ -lactams (due to their limited thermal stability)<sup>74</sup> and fluoroquinolones, rifampicin, tetracyclines, and macrolides (because of their potential local detrimental effects on cell viability and osteogenic activity).<sup>74-76</sup>

In contrast, the recently published VANCO study demonstrated that topical vancomycin directly applied to the surgical area during wound closure reduces gram-positive FRI rates.<sup>77</sup> In turn, vancomycin local application has limited systemic absorption, making it a safe alternative regarding nephrotoxicity.<sup>78</sup>

Although the local administration of antibiotics is usually deemed safe,<sup>78,79</sup> we cannot neglect the potential for local and systemic toxicity.<sup>75,80,81</sup> In addition, there is no solid evidence favoring antibiotic therapy without a local delivery method, except for vancomycin.<sup>82-84</sup> No study evaluated its direct use for FRI treatment.

**Table 2** Antibiotic Agents

Organism		Antibiotic agent	Dosage	Frequency	Route
<i>Staphylococcus</i> spp.					
Methicillin-sensitive (MS)	–	Flucloxacillin	2 g	6 h	IV
	and	Rifampicin	300-450 mg	12 h	PO
		<i>Followed by oral treatment (depending on sensitivity)</i>			
	–	Rifampicin	300-450 mg	12 h	PO
		<i>and (per sensitivity)</i>			
	–	Levofloxacin	500 mg	12 h	PO
	or	Trimethoprim/Sulfamethoxazole	960 mg	8 h	PO
	or	Doxycycline	600 mg	12 h	PO
	or	Clindamycin	600 mg	8 h	PO
Methicillin-resistant (MR)	–	Vancomycin	Loading dose: 25-30 mg/kg Maintenance dose: 15 mg/kg	12 h	IV
	and	Rifampicin	300-450 mg	12 h	PO
		<i>Followed by oral treatment (depending on sensitivity)</i>			
	–	Rifampicin	300-450 mg	12 h	PO
		<i>and (per sensitivity)</i>			
	–	Levofloxacin	500 mg	12 h	PO
	or	Trimethoprim/Sulfamethoxazole	960 mg	8 h	PO
	or	Doxycycline	600 mg	12 h	PO
	or	Clindamycin	600 mg	8 h	PO
Rifampicin-resistant		<i>Intravenous treatment per sensitivity followed by antibiotic suppression until bone consolidation and implant removal</i>			
<i>Streptococcus</i> spp.		<i>(per sensitivity)</i>			
		Penicillin G	5 million units	6 h	IV
	–		or		
			4 million units	5 h	IV
	or	Ceftriaxone	2 g	24 h	IV
		<i>Followed by oral treatment per sensitivity</i>			
	–	Amoxicillin	1 g	6-8 h	PO
	or	Clindamycin	450-600 mg	8 h	PO
<i>Enterococcus</i> spp.					
Penicillin-sensitive	–	Ampicillin	2 g	6 h	IV
	and	Gentamicin	3 mg/kg	24 h	IV
		<i>Followed by oral treatment per sensitivity</i>			
	–	Amoxicillin	1 g	6-8 h	PO
Penicillin-resistant					
	–	Vancomycin	Loading dose: 25-30 mg/kg Maintenance dose: 15 mg/kg	12 h	IV
		<i>Followed by oral treatment per sensitivity</i>			
	–	Linezolid (maximum, 4 weeks)	600 mg	12 h	PO
Gram-negative organisms					
Enterobacteriaceae (e.g., <i>Escherichia coli</i> ,	–	<i>Beta-lactam per sensitivity</i>			

(Continued)

**Table 2** (Continued)

Organism		Antibiotic agent	Dosage	Frequency	Route
<i>Klebsiella</i> spp., and <i>Enterobacter</i> spp.)					
		<i>Followed by oral treatment per sensitivity</i>			
	–	Ciprofloxacin	750 mg	12 h	PO
	or	Levofloxacin	500 mg	12 h	PO
Non-fermenting (e.g., <i>Pseudomonas aeruginosa</i> , <i>Acinetobacter</i> spp.)	–	Meropenem	2 g	8 h	IV
	or	Ceftazidime	2 g	8 h	
	and	Gentamicin (short course)	5 mg/kg	24 h	IV
	or	Tobramycin (short course)	5 mg/kg	24 h	
	or	Amikacin (short course)	15 mg/kg	24 h	IV
		<i>Followed by oral treatment per sensitivity</i>			
	–	Ciprofloxacin	750 mg	12 h	PO
	or	Levofloxacin	500 mg	12 h	PO
Ciprofloxacin-resistant		<i>Intravenous treatment per sensitivity followed by antibiotic suppression until bone consolidation and implant removal</i>			
Negative culture	–	Ampicillin/sulbactam (2 weeks)	3 g	8 h	IV
		<i>Followed by oral treatment</i>			
		Rifampicin	300-450 mg	12 h	PO
	and	Levofloxacin	500 mg	12 h	PO

Abbreviations: IV, intravenous; PO, per os.

\*The antibiotic treatment must be discussed with the infectious disease team of each center for adaptation to the local reality.

Table adapted from Depypere et al.<sup>9</sup>

## Conclusions

FRI continues to be a challenging complication for the entire healthcare team. As such, multidisciplinary confrontation is essential to evaluate and manage these patients in all dimensions. Today, scientific high-level evidence for FRI management guidance is limited. Therefore, diagnosis and treatment standardization per the recently published clinical guidelines seeks to improve outcomes and the level of evidence to guide the management of these patients.

This study received no specific funding.

### Conflict of Interests

No.

### Acknowledgments

We thank Dr. Willem-Jan Metsemakers for his significant contribution to the subject over the past few years and for allowing us to adapt his figures for this publication.

## Bibliography

- Metsemakers WJ, Onsea J, Neutjens E, et al. Prevention of fracture-related infection: a multidisciplinary care package. *Int Orthop* 2017; 41(12):2457–2469
- Morgenstern M, Moriarty TF, Kuehl R, et al. International survey among orthopaedic trauma surgeons: Lack of a definition of fracture-related infection. *Injury* 2018;49(03):491–496
- Berkes M, Obremskey WT, Scannell B, Ellington JK, Hymes RA, Bosse M Southeast Fracture Consortium. Maintenance of hardware after early postoperative infection following fracture internal fixation. *J Bone Joint Surg Am* 2010;92(04):823–828
- Tschudin-Sutter S, Frei R, Dangel M, et al. Validation of a treatment algorithm for orthopaedic implant-related infections with device-retention—results from a prospective observational cohort study. *Clin Microbiol Infect* 2016;22(05):457.e1–457.e9
- Thakore RV, Greenberg SE, Shi H, et al. Surgical site infection in orthopedic trauma: A case-control study evaluating risk factors and cost. *J Clin Orthop Trauma* 2015;6(04):220–226
- Darouiche RO. Treatment of infections associated with surgical implants. *N Engl J Med* 2004;350(14):1422–1429
- Metsemakers WJ, Kortram K, Morgenstern M, et al. Definition of infection after fracture fixation: A systematic review of randomized controlled trials to evaluate current practice. *Injury* 2018;49(03): 497–504
- Metsemakers WJ, Morgenstern M, McNally MA, et al. Fracture-related infection: A consensus on definition from an international expert group. *Injury* 2018;49(03):505–510
- Depypere M, Kuehl R, Metsemakers WJ, et al; Fracture-Related Infection (FRI) Consensus Group. Recommendations for Systemic Antimicrobial Therapy in Fracture-Related Infection: A Consensus From an International Expert Group. *J Orthop Trauma* 2020;34 (01):30–41



- 10 Metsemakers WJ, Fragomen AT, Moriarty TF, et al; Fracture-Related Infection (FRI) consensus group. Evidence-Based Recommendations for Local Antimicrobial Strategies and Dead Space Management in Fracture-Related Infection. *J Orthop Trauma* 2020;34(01):18–29
- 11 Govaert GAM, Kuehl R, Atkins BL, et al; Fracture-Related Infection (FRI) Consensus Group. Diagnosing Fracture-Related Infection: Current Concepts and Recommendations. *J Orthop Trauma* 2020; 34(01):8–17
- 12 Metsemakers WJ, Morgenstern M, Senneville E, et al; Fracture-Related Infection (FRI) group. General treatment principles for fracture-related infection: recommendations from an international expert group. *Arch Orthop Trauma Surg* 2020;140(08): 1013–1027
- 13 Cats-Baril W, Gehrke T, Huff K, Kendoff D, Maltenfort M, Parvizi J. International consensus on periprosthetic joint infection: description of the consensus process. *Clin Orthop Relat Res* 2013; 471(12):4065–4075
- 14 Kleber C, Schaser KD, Trampuz A. [Complication management of infected osteosynthesis: Therapy algorithm for peri-implant infections]. *Chir Z Alle Geb Oper Medizin* 2015;86(10):925–934
- 15 Patzakis MJ, Zalavras CG. Chronic posttraumatic osteomyelitis and infected nonunion of the tibia: current management concepts. *J Am Acad Orthop Surg* 2005;13(06):417–427
- 16 Bonneville P, Bonnomet F, Philippe R, et al; SOFCOT. Early surgical site infection in adult appendicular skeleton trauma surgery: a multicenter prospective series. *Orthop Traumatol Surg Res* 2012;98(06):684–689
- 17 Bezstarosti H, Van Lieshout EMM, Voskamp LW, et al. Insights into treatment and outcome of fracture-related infection: a systematic literature review. *Arch Orthop Trauma Surg* 2019;139(01): 61–72
- 18 Onsea J, Van Lieshout EMM, Zalavras C, et al. Validation of the diagnostic criteria of the consensus definition of fracture-related infection. *Injury* 2022;53(06):1867–1879
- 19 Tiemann AH, Hofmann GO. Principles of the therapy of bone infections in adult extremities : Are there any new developments? *Strateg Trauma Limb Reconstr* 2009;4(02):57–64
- 20 Omar M, Suero EM, Lioudakis E, et al. Diagnostic performance of swab PCR as an alternative to tissue culture methods for diagnosing infections associated with fracture fixation devices. *Injury* 2016;47(07):1421–1426
- 21 van den Kieboom J, Bosch P, Plate DJJ, et al. Diagnostic accuracy of serum inflammatory markers in late fracture-related infection: a systematic review and meta-analysis. *Bone Joint J* 2018;100-B (12):1542–1550
- 22 Govaert GA, Ijpmma FF, McNally M, McNally E, Reininga IH, Glaudemans AW. Accuracy of diagnostic imaging modalities for peripheral post-traumatic osteomyelitis - a systematic review of the recent literature. *Eur J Nucl Med Mol Imaging* 2017;44(08): 1393–1407
- 23 Goebel M, Rosa F, Tatsch K, Grillhoesl A, Hofmann GO, Kirschner MH. [Diagnosis of chronic osteitis of the bones in the extremities. Relative value of F-18 FDG-PET]. *Unfallchirurg* 2007;110(10): 859–866
- 24 Kaim A, Ledermann HP, Bongartz G, Messmer P, Müller-Brand J, Steinbrich W. Chronic post-traumatic osteomyelitis of the lower extremity: comparison of magnetic resonance imaging and combined bone scintigraphy/immunoscintigraphy with radiolabelled monoclonal antigranulocyte antibodies. *Skeletal Radiol* 2000;29 (07):378–386
- 25 Ballani NS, Al-Huda FA, Khan HA, Al-Mohannadi S, Mahmood H, Al-Enezi F. The value of quantitative uptake of (99m)Tc-MDP and (99m)Tc-HMPAO white blood cells in detecting osteomyelitis in violated peripheral bones. *J Nucl Med Technol* 2007;35(02): 91–95
- 26 Meller J, Köster G, Liersch T, et al. Chronic bacterial osteomyelitis: prospective comparison of (18)F-FDG imaging with a dual-head coincidence camera and (111)In-labelled autologous leucocyte scintigraphy. *Eur J Nucl Med Mol Imaging* 2002;29(01):53–60
- 27 Glaudemans AWJM, de Vries EFJ, Vermeulen LEM, Slart RHJA, Dierckx RAJO, Signore A. A large retrospective single-centre study to define the best image acquisition protocols and interpretation criteria for white blood cell scintigraphy with <sup>99m</sup>Tc-HMPAO-labelled leucocytes in musculoskeletal infections. *Eur J Nucl Med Mol Imaging* 2013;40(11):1760–1769
- 28 Govaert GAM, Bosch P, Ijpmma FFA, et al. High diagnostic accuracy of white blood cell scintigraphy for fracture related infections: Results of a large retrospective single-center study. *Injury* 2018; 49(06):1085–1090
- 29 Lemans JVC, Hobbelink MGG, Ijpmma FFA, et al. The diagnostic accuracy of <sup>18</sup>F-FDG PET/CT in diagnosing fracture-related infections. *Eur J Nucl Med Mol Imaging* 2019;46(04):999–1008
- 30 Schwotzer N, Wahl P, Fracheboud D, Gautier E, Chuard C. Optimal culture incubation time in orthopedic device-associated infections: a retrospective analysis of prolonged 14-day incubation. *J Clin Microbiol* 2014;52(01):61–66
- 31 Firoozabadi R, Alton T, Wenke J. Novel Strategies for the Diagnosis of Posttraumatic Infections in Orthopaedic Trauma Patients. *J Am Acad Orthop Surg* 2015;23(07):443–451
- 32 Greene LR. Guide to the elimination of orthopedic surgery surgical site infections: an executive summary of the Association for Professionals in Infection Control and Epidemiology elimination guide. *Am J Infect Control* 2012;40(04):384–386
- 33 Myers WT, Leong M, Phillips LG. Optimizing the patient for surgical treatment of the wound. *Clin Plast Surg* 2007;34(04):607–620
- 34 Underwood P, Askari R, Hurwitz S, Chamarthi B, Garg R. Preoperative A1C and clinical outcomes in patients with diabetes undergoing major noncardiac surgical procedures. *Diabetes Care* 2014; 37(03):611–616
- 35 Ernst A, Wilson JM, Ahn J, Shapiro M, Schenker ML. Malnutrition and the Orthopaedic Trauma Patient: A Systematic Review of the Literature. *J Orthop Trauma* 2018;32(10):491–499
- 36 Duggan EW, Carlson K, Umpierrez GE. Perioperative Hyperglycemia Management: An Update. *Anesthesiology* 2017;126(03): 547–560
- 37 Allegranzi B, Zayed B, Bischoff P, et al; WHO Guidelines Development Group. New WHO recommendations on intraoperative and postoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infect Dis* 2016;16 (12):e288–e303
- 38 Haines N, Kempton LB, Seymour RB, et al. The effect of a single early high-dose vitamin D supplement on fracture union in patients with hypovitaminosis D: a prospective randomised trial. *Bone Joint J* 2017;99-B(11):1520–1525
- 39 Zargarán A, Zargarán D, Trompeter AJ. The role of Vitamin D in orthopaedic infection: a systematic literature review. *Bone Jt Open* 2021;2(09):721–727
- 40 Mah TF, O'Toole GA. Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* 2001;9(01):34–39
- 41 Metsemakers WJ, Kuehl R, Moriarty TF, et al. Infection after fracture fixation: Current surgical and microbiological concepts. *Injury* 2018;49(03):511–522
- 42 Rittmann WW, Perren SM, Furnas DW. CORTICAL BONE HEALING AFTER INTERNAL FIXATION AND INFECTION. *Plast Reconstr Surg* 1976;57(01):91
- 43 Worlock P, Slack R, Harvey L, Mawhinney R. The prevention of infection in open fractures: an experimental study of the effect of fracture stability. *Injury* 1994;25(01):31–38
- 44 Schmidt AH, Swiontkowski MF. Pathophysiology of infections after internal fixation of fractures. *J Am Acad Orthop Surg* 2000;8(05):285–291

- 45 Foster AL, Moriarty TF, Zalavras C, et al. The influence of bio-mechanical stability on bone healing and fracture-related infection: the legacy of Stephan Perren. *Injury* 2021;52(01):43–52
- 46 Morgenstern M, Kuehl R, Zalavras CG, et al. The influence of duration of infection on outcome of debridement and implant retention in fracture-related infection. *Bone Joint J* 2021;103-B(02):213–221
- 47 Bhandari M, Jeray KJ, Petrisor BA, et al; FLOW Investigators. A Trial of Wound Irrigation in the Initial Management of Open Fracture Wounds. *N Engl J Med* 2015;373(27):2629–2641
- 48 Penn-Barwell JG, Murray CK, Wenke JC. Comparison of the antimicrobial effect of chlorhexidine and saline for irrigating a contaminated open fracture model. *J Orthop Trauma* 2012;26(12):728–732
- 49 O'Donnell JA, Wu M, Cochrane NH, et al. Efficacy of common antiseptic solutions against clinically relevant microorganisms in biofilm. *Bone Joint J* 2021;103-B(05):908–915
- 50 Markel JF, Bou-Akl T, Dietz P, Afsari AM. The Effect of Different Irrigation Solutions on the Cytotoxicity and Recovery Potential of Human Osteoblast Cells In Vitro. *Arthroplast Today* 2021;7:120–125
- 51 Böhle S, Röhner E, Zippelius T, Jacob B, Matziolis G, Rohe S. Cytotoxic effect of sodium hypochlorite (Lavanox 0.08%) and chlorhexidine gluconate (Irrisept 0.05%) on human osteoblasts. *Eur J Orthop Surg Traumatol Orthop Traumatol* 2021
- 52 Zalavras CG. Prevention of Infection in Open Fractures. *Infect Dis Clin North Am* 2017;31(02):339–352
- 53 Schemitsch EH. Size Matters: Defining Critical in Bone Defect Size!. *J Orthop Trauma* 2017;31(Suppl 5):S20–S22
- 54 Giannoudis PV, Tosounidis TH. Acute and chronic infection: Is there a gold standard for management of the wound and bone defect? *OTA Int* 2020;3(01):e068
- 55 Morgenstern M, Vallejo A, McNally MA, et al. The effect of local antibiotic prophylaxis when treating open limb fractures: A systematic review and meta-analysis. *Bone Joint Res* 2018;7(07):447–456
- 56 ter Boo GJA, Grijpma DW, Moriarty TF, Richards RG, Eglin D. Antimicrobial delivery systems for local infection prophylaxis in orthopedic- and trauma surgery. *Biomaterials* 2015;52:113–125
- 57 Aurégan JC, Bégue T. Bioactive glass for long bone infection: a systematic review. *Injury* 2015;46(Suppl 8):S3–S7
- 58 Kojima KE, de Andrade E Silva FB, Leonhardt MC, et al. Bioactive glass S53P4 to fill-up large cavitary bone defect after acute and chronic osteomyelitis treated with antibiotic-loaded cement beads: A prospective case series with a minimum 2-year follow-up. *Injury* 2021;52(Suppl 3):S23–S28
- 59 Li HK, Rombach I, Zambellas R, et al; OVIVA Trial Collaborators. Oral versus Intravenous Antibiotics for Bone and Joint Infection. *N Engl J Med* 2019;380(05):425–436
- 60 Fassbender M, Minkwitz S, Kronbach Z, et al. Local gentamicin application does not interfere with bone healing in a rat model. *Bone* 2013;55(02):298–304
- 61 Zimmerli W, Widmer AF, Blatter M, Frei R, Ochsner PE Foreign-Body Infection (FBI) Study Group. Role of rifampin for treatment of orthopedic implant-related staphylococcal infections: a randomized controlled trial. *JAMA* 1998;279(19):1537–1541
- 62 El Helou OC, Barbari EF, Lahr BD, et al. Efficacy and safety of rifampin containing regimen for staphylococcal prosthetic joint infections treated with debridement and retention. *Eur J Clin Microbiol Infect Dis* 2010;29(08):961–967
- 63 Aboltins CA, Dowsey MM, Buising KL, et al. Gram-negative prosthetic joint infection treated with debridement, prosthesis retention and antibiotic regimens including a fluoroquinolone. *Clin Microbiol Infect* 2011;17(06):862–867
- 64 Martínez-Pastor JC, Muñoz-Mahamud E, Vilchez F, et al. Outcome of acute prosthetic joint infections due to gram-negative bacilli treated with open debridement and retention of the prosthesis. *Antimicrob Agents Chemother* 2009;53(11):4772–4777
- 65 Sendi P, Zimmerli W. Antimicrobial treatment concepts for orthopaedic device-related infection. *Clin Microbiol Infect* 2012;18(12):1176–1184
- 66 Achermann Y, Eigenmann K, Ledergerber B, et al. Factors associated with rifampin resistance in staphylococcal periprosthetic joint infections (PJI): a matched case-control study. *Infection* 2013;41(02):431–437
- 67 Senneville E, Joulie D, Legout L, et al. Outcome and predictors of treatment failure in total hip/knee prosthetic joint infections due to *Staphylococcus aureus*. *Clin Infect Dis* 2011;53(04):334–340
- 68 Zimmerli W, Trampuz A, Ochsner PE. Prosthetic-joint infections. *N Engl J Med* 2004;351(16):1645–1654
- 69 Morgenstern M, Athanasou NA, Ferguson JY, Metsemakers WJ, Atkins BL, McNally MA. The value of quantitative histology in the diagnosis of fracture-related infection. *Bone Joint J* 2018;100-B(07):966–972
- 70 Sheehy SH, Atkins BA, Bejon P, et al. The microbiology of chronic osteomyelitis: prevalence of resistance to common empirical anti-microbial regimens. *J Infect* 2010;60(05):338–343
- 71 Kuehl R, Tschudin-Sutter S, Morgenstern M, et al. Time-dependent differences in management and microbiology of orthopaedic internal fixation-associated infections: an observational prospective study with 229 patients. *Clin Microbiol Infect* 2019;25(01):76–81
- 72 Gille J, Wallstabe S, Schulz AP, Paech A, Gerlach U. Is non-union of tibial shaft fractures due to nonculturable bacterial pathogens? A clinical investigation using PCR and culture techniques. *J Orthop Surg Res* 2012;7:20
- 73 Palmer MP, Altman DT, Altman GT, et al. Can we trust intra-operative culture results in nonunions? *J Orthop Trauma* 2014;28(07):384–390
- 74 Samara E, Moriarty TF, Decosterd LA, Richards RG, Gautier E, Wahl P. Antibiotic stability over six weeks in aqueous solution at body temperature with and without heat treatment that mimics the curing of bone cement. *Bone Joint Res* 2017;6(05):296–306
- 75 Rathbone CR, Cross JD, Brown KV, Murray CK, Wenke JC. Effect of various concentrations of antibiotics on osteogenic cell viability and activity. *J Orthop Res* 2011;29(07):1070–1074
- 76 Holtom PD, Pavkovic SA, Bravos PD, Patzakis MJ, Shepherd LE, Frenkel B. Inhibitory effects of the quinolone antibiotics trovafloxacin, ciprofloxacin, and levofloxacin on osteoblastic cells in vitro. *J Orthop Res* 2000;18(05):721–727
- 77 O'Toole RV, Joshi M, Carlini AR, et al; Major Extremity Trauma Research Consortium (METRC) Effect of Intra-wound Vancomycin Powder in Operatively Treated High-risk Tibia Fractures: A Randomized Clinical Trial. *JAMA Surg* 2021;156(05):e207259. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/33760010/>
- 78 O'Toole RV, Degani Y, Carlini AR, Castillo RC, O'Hara NN, Joshi M, Mand METRC. Systemic Absorption and Nephrotoxicity Associated With Topical Vancomycin Powder for Fracture Surgery. *J Orthop Trauma* 2021;35(01):29–34
- 79 Springer BD, Lee GC, Osmon D, Haidukewych GJ, Hanssen AD, Jacofsky DJ. Systemic safety of high-dose antibiotic-loaded cement spacers after resection of an infected total knee arthroplasty. *Clin Orthop Relat Res* 2004;(427):47–51
- 80 Antoci V Jr, Adams CS, Hickok NJ, Shapiro IM, Parvizi J. Antibiotics for local delivery systems cause skeletal cell toxicity in vitro. *Clin Orthop Relat Res* 2007;462(462):200–206

- 81 van Raaij TM, Visser LE, Vulto AG, Verhaar JAN. Acute renal failure after local gentamicin treatment in an infected total knee arthroplasty. *J Arthroplasty* 2002;17(07):948–950
- 82 Tubaki VR, Rajasekaran S, Shetty AP. Effects of using intravenous antibiotic only versus local intrawound vancomycin antibiotic powder application in addition to intravenous antibiotics on postoperative infection in spine surgery in 907 patients. *Spine* 2013;38(25):2149–2155
- 83 Adogwa O, Elsamadicy AA, Sergesketter A, et al. Prophylactic use of intraoperative vancomycin powder and postoperative infection: an analysis of microbiological patterns in 1200 consecutive surgical cases. *J Neurosurg Spine* 2017;27(03):328–334
- 84 Ghobrial GM, Thakkar V, Andrews E, et al. Intraoperative vancomycin use in spinal surgery: single institution experience and microbial trends. *Spine* 2014;39(07):550–555