

Comments on "Utility of First Dorsal Metacarpal Artery Flap for Thumb Defects"

Ilker Uyar¹⁰

¹ Department of Plastic, Reconstructive and Aesthetic Surgery, Izmir Katip Celebi University Medical Faculty, Izmir, Turkey

Indian J Plast Surg 2023;56:290-291.

Address for correspondence Ilker Uyar, MD, Izmir Katip Celebi University Ataturk Training and Research Hospital, Basın Sitesi, Atatürk Eğitim ve Araştırma Hastanesi, 35360 Karabağlar/İzmir, Turkey (e-mail: ilkeruyar@windowslive.com).

Dear Editor,

I reviewed the study "Utility of First Dorsal Metacarpal Artery Flap for Thumb Defects," by K. Aggarwal and K. Singh, published online in the *Indian Journal of Plastic Surgery*.¹ The study is well designed and the findings are presented in the discussion section accompanied by the literature.

The first dorsal metacarpal artery (FDMA) flap is particularly useful for 1st and 3rd finger defects.^{2,3} The advantages of this flap are that its vascular anatomy is stable, its learning



Fig. 1 View of a patient who developed partial necrosis after venous insufficiency during follow-up.

article published online April 19, 2023 DOI https://doi.org/ 10.1055/s-0043-57269. ISSN 0970-0358. curve is short, and its donor site morbidity is low.⁴ The major disadvantage is venous insufficiency occurring in the flap.^{1,4} Venous insufficiency and necrosis are inevitable in cases where the width of the pedicle is narrow and the pedicle is compressed (**~ Fig. 1**).

If the FDMA flap is to be designed as an island, the pedicle width should be at least 5 mm to prevent venous insufficiency.² In addition, if the flap is to be transferred to the recipient area by tunneling, it should be ensured that the tunnel width is sufficient. If a skin incision is to be made without tunneling during the flap transfer phase, the incised skin should not be resutured on the pedicle. It should be preferable to place a graft on the incision line (**-Fig. 2**).



Fig. 2 The pedicle width should be at least 5 mm to prevent venous insufficiency.

© 2023. Association of Plastic Surgeons of India. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/ licenses/by-nc-nd/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

In this study by K. Aggarwal and K. Singh, it was understood that the skin on the pedicle was incised at the flap inset stage in some patients and sutured again after flap transfer. After the incision, tissue contraction and edema occurred. When the edematous contracted tissue is resutured, it compresses the pedicle below, and after compression, circulation disorder occurs in the flap. This may be the cause of partial necrosis in some patients in the study. Therefore, pedicle width of at least 5 mm and the use of a sufficiently wide tunneling technique can prevent venous problems during flap transfer.² If a skin incision is to be made without using a tunnel, the skin should not be resutured, but the pedicle should be grafted.

Authorship

The conception and design of the study, or acquisition of data, or analysis and interpretation of data: Ilker Uyar Drafting the article or revising it critically for important intellectual content: Ilker Uyar Final approval of the version to be submitted: Ilker Uyar

Informed Consent Informed consent was obtained.

Conflict of Interest None declared.

References

- 1 Aggarwal K, Singh K. Utility of first dorsal metacarpal artery flap for thumb defects. Indian J Plast Surg 2022;55(04):368–375
- 2 Uyar I, Basol TB. Use of the first dorsal metacarpal artery flap in finger defects – case series. Selcuk Med J 2022;38(04): 180–185
- 3 Yannascoli SM, Thibaudeau S, Levin LS. Management of soft tissue defects of the hand. J Hand Surg Am 2015;40(06):1237–1244, quiz 1245
- 4 Al-Baza TH, Gadb SS, Keshkc TF, Alyc MS. Evaluation of dorsal metacarpal artery perforator flaps in the reconstruction of hand soft-tissue defects. Menoufia Med J 2019;32(01):1256–1261