

Seizure-Related Injuries among Patients with Epilepsy at the Douala General Hospital, Cameroon: A Retrospective Cohort Study

Cyrille Nkouonlack^{1®} Adrien Yongho Mounchili² Daniel Massi Gams^{1,3®} James Fenibe Mbinta⁴ Alain Mefire Chichom⁵ Yacouba Njankouo Mapoure^{3,6}

¹Department of Internal Medicine and Paediatrics, Faculty of Health Sciences, University of Buea, Buea, Cameroon

² Department of Internal Medicine, Douala General Hospital, Douala, Cameroon

³Douala General Hospital, Douala, Cameroon

⁴ School of Health, Wellington Faculty of Health, Victoria University of Wellington, Wellington, New Zealand

⁵Department of Surgery, Faculty of Health Sciences, University of Buea, Buea, Cameroon

⁶Department of Clinical Sciences, Faculty of Medicine and Pharmaceutical Sciences, University of Douala, Douala, Cameroon

Int J Ep 2023;9:13-18.

Address for correspondence Cyrille Nkouonlack, MD, Department of Internal Medicine and Paediatrics, Faculty of Health Sciences, University of Buea, P.O. Box 560, Buea, South West Region, Cameroon (e-mail: nkouonlack@gmail.com).

Abstract Background Seizures can lead to injuries in patients with epilepsy (PWE). This study seeks to determine the 1-year incidence, types, and associated factors of seizure-related injuries in PWE compared with that of patients without epilepsy.

Methods A retrospective cohort study was carried out at the Douala General Hospital, Douala, Cameroon. Patients with active epilepsy within the past 5 years were compared to patients without epilepsy after matching them by age and gender during a 12-month period. Data was collected from patient records using a question-naire and analyzed using the SPSS version 20 software. The chi-square test was used to compare the proportions of injuries at a significant level of p < 0.05 and a confidence interval (CI) of 95%.

Results A total of 155 participants were recruited in each group. Note that 51.6% were males, and the mean age was 25.46 ± 16.35 years. The 1-year incidence of all injuries was 26.5% in PWE against 16.8% among patients without epilepsy, respectively (p = 0.038; odds ratio = 1.8, 95\% CI: 1.03-3.10). Soft tissue injury was higher in PWE (82.9%, p < 0.05) compared to those without epilepsy (50%), whereas patients without epilepsy had more cases of burns compared to PWE (p < 0.05). PWE and having frequent seizures with falls and those on more than one antiseizure medication had an increased risk of injuries (p < 0.05).

Keywords

- epileptic seizures
- physical injuries
- antiseizure medication

Conclusion PWE have a double risk of having mild soft tissue injuries attributed to epileptic seizures. The risk is increased in PWE having seizures with falls and those taking more than one antiseizure medication.

article published online August 6, 2024 DOI https://doi.org/ 10.1055/s-0044-1788265. ISSN 2213-6320.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/ licenses/by-nc-nd/4.0/)

^{© 2024.} Indian Epilepsy Society. All rights reserved.

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Introduction

Epilepsy is a common disease in Cameroon with the highest prevalence worldwide reported in the Mbam valley, though prevalence studies on the whole country have not yet been done.^{1–3} Most patients with epilepsy (PWE) in the country seek medical attention more than a year after the onset of seizures, and they do so directly in referral centres.⁴ The management of epilepsy is still a major challenge in Cameroon, very difficult mainly due to lack of human and material resources, limited access to antiseizure medications, and the stigma associated with epilepsy.^{5–7} Seizures are unforeseen events that can cause physical injury, especially when associated with loss of consciousness, and occurring in high-risk environments. Several studies in Cameroon have focused on the psychological impact and the stigma associated with epilepsy, however, there is scarcity of data on physical injuries that patients living with epilepsy endure. This study aimed at determining the 1-year incidence of physical injuries attributable to seizures in PWE attending the Douala General Hospital, Douala, Cameroon and compared it with that of patients without epilepsy.

Materials and Methods

Study Design

We carried out a retrospective cohort study in the Douala General Hospital in the Neurological Unit of the Internal Medicine Department from January 1 to December 1, 2017.

Study Setting

The Douala General Hospital is a third-level reference hospital in the country and the Central Africa zone and is also a university teaching hospital. The hospital offers facilities like neurophysiologic studies (electroencephalogram and electromyogram), imaging studies (computed tomography scan, magnetic resonance imaging, ultrasonography, etc.), and laboratory analysis with international standards among other facilities. Patient files and records are kept in alphabetical order and classified according to the attending neurologist.

Study Population

The population of interest was all patients diagnosed with epilepsy at the Douala General Hospital and have had at least one seizure over the past 5 years, that is, from 2010 to 2016. However, those with active neurological, psychiatric, or other comorbid conditions (diabetes mellitus, hypertension, stroke, etc.) were excluded. Using an exhaustive sampling method, files of PWE were selected from the records and incomplete files were excluded. This constituted the exposure group, and these were matched according to age (± 1 year) and gender with another group of patients without epilepsy selected randomly from the general population as the unexposed group.

Sociodemographic and clinical data on epilepsy were collected from the files of PWE, and the patients were interviewed at follow-up visits in the outpatient department, or by telephone for the occurrence of physical injuries, due to seizures, over the study period.

Data collection of patients without epilepsy was done in public places or institutions like schools or markets. Verbal and written consent were obtained before questionnaire administration.

The questionnaire included sociodemographic questions (age, gender, occupation, and educational level), clinical questions on epilepsy (age at onset of epilepsy, age at diagnosis, types and frequency of seizures, seizures with fall, antiseizure medications used), and information on physical injury in the past 12 months. We defined some terms as follow:

- PWE: people with clinical and/or electrophysiologic diagnosis of epilepsy and confirmed by a neurologist.
- Patients without epilepsy: people in the general population who have never experienced a seizure in the past.
- Injury: harm or hurt; damage inflicted on the body by an external force.
- Epilepsy type: This was done according to the 2017 International League Against Epilepsy classification of epilepsies as follows; generalized epilepsy, focal epilepsy, combined generalized and focal epilepsy, and unknown.⁸

Data Analysis

The incidence of physical injury was determined by dividing the number of epileptic patients who sustained at least one seizure-related injury within a period of 12 months with the total number of epileptic patients at the start of the study. The chi-square test was used to compare the proportions of physical injury in people with and without epilepsy, at a significance level of p < 0.05 and a confidence interval (CI) of 95%. SPSS version 20.0 was used for statistical analysis. A patient could have more than one type of injury during the study period (e.g., both a limb fracture and a burn), but each type of injury was counted only once per patient during the 1-year period. Injuries that were not seizure-related in PWE was not considered as an injury. The factors that could predict injury were found using binary logistic regression with their respective odds ratios (ORs) at 95% CI.

Ethical Considerations

This study was approved by the Institutional Review Board of the Faculty of Health Sciences, University of Buea, Buea, Cameroon (Ref: 2018/144/UB/SG/IRB/FHS of 31.01.2018) and administrative approval and authorization to carry out the research in the Douala General Hospital was obtained from the Director General of the hospital (Ref: No. 009/AR/MINSANTE/HGD/DM/01/18).

Results

A total of 155 participants were enrolled out of the 239 files reviewed, giving a response rate of 89%. Eighty of the PWE (51.6%) were male. The male:female ratio was 1.1:1. The mean age of PWE was 25.46 ± 16.5 years, with a range from 2 to 86 years. More than half (n = 78, 50.3%) of the study participants were less than 20 years old, and only 11% (n = 17) of them were above 50. The mean age at seizure onset was 19.52 ± 16.37 years and the mean age at diagnosis was 21.31 ± 16.33 years. More than half (n = 81, 52.3%) of PWE had had no seizures for the past 1 year and 38.1% (n = 59) had 1 to 12 seizures. This gives a cumulative percentage of 90.3%, meaning only less than 10% of PWE had more than one seizure a month. Generalized epilepsies were the most common type of epilepsy among the participants (n = 111, 71.6%) of all seizures (p < 0.001). The other types include focal epilepsies and unknown type. A total of 106 (68.4%) PWE had seizures that led to falls, and of these, 91.5% (n = 97) had generalized epilepsy (p < 0.001). The antiseizure medications used were valproate in 76 PWE (49.0%), carbamazepine in 37 (23.9%), phenobarbital in 17 (11.0%), clonazepam in 9 (5.8%), clobazam in 8 (5.2%), and lamotrigine in 2 (1.3%). However, 2.7% of PWE (n=5) were on traditional remedy and 15.4% (n = 28) were not on any antiseizure medication. Twenty epileptics (12.9%) were on more than one antiseizure medication.

The 1-year incidence of overall physical injury (having at least one injury) was 26.5% (n = 41) among those with epilepsy and 16.8% (n = 16) among those without epilepsy (p = 0.038, OR = 1.8, 95% CI: 1.03–3.10). Soft tissue injury was the only type of injury that was significantly high (p < 0.001)

in PWE compared to those without epilepsy (**-Table 1**). Tongue bites accounted for this high level. Burns (7.3% vs. 26.9%) were higher in patients without epilepsy at a significant level (p = 0.038); however, all cases of burns in PWE were due to flames. Head injury was also high in PWE, but the value was not applicable statistically because we had no case among patients without epilepsy.

There were more cases of road traffic accidents (p = 0.02) in patients without epilepsy (n = 11, 7.1%) than in PWE (n = 2, 1.3%). There was no case of drowning in PWE. Injuries occurred mostly at home in PWE, as opposed to injuries occurring in the streets in patients without epilepsy as shown in **- Table 2**.

There was no significant difference between PWE and patients without epilepsy regarding medical attention and hospitalization following injury (\succ Table 3).

As shown in **-Table 4**, after binary logistic regression analysis, having a fall during seizures and using more than one antiseizure medication were associated with physical injury.

Discussion

This hospital-based retrospective study aimed at determining the incidence of physical injury attributable to seizures in

Table I Comparison of types of injuries between PWE and patients without ep	epilepsy
------------------------------------------------------------------------------------	----------

Injury	PWE N (%)	Non-PWE N (%)	OR (95% CI)	p-Value
Burn	3 (7.3)	7 (26.9)	0.2 (0.05–0.92)	0.038
Scald Flame Chemical Thermal	0 3 (100) 0 0	2 (28.6) 0 1 (14.3) 4 (57.1)		0.298 ^a 0.002 ^a 0.490 ^a 0.091 ^a
Soft tissue injury	34 (82.9)	13 (50)	4.9 (1.59–14.88)	0.004
Abrasion Laceration Hematoma Tongue bite	11 (32.4) 8 (23.5) 2 (5.9) 13 (38.2)	5 (38.5) 7 (53.8) 1 (7.7) 0		0.689 0.046 0.841 0.009 ^a
Head injury	13 (31.7)	0		0.001
Abrasion Laceration Intracranial hematoma Contusion	9 (69.2) 2 (15.4) 0 2 (15.4)	0 0 0 0		
Dental injury	2 (4.9)	1 (3.8)	1.2 (0.11–14.89)	0.842
Tooth loss Jaw fracture	2 (100) 0	0 1 (100)		0.089 ^a 0.089 ^a
Fracture	1 (2.4)	3 (11.5)	0.2 (0.02–1.95)	0.126
Upper limb Lower limb Vertebrae Pelvis	0 1 (100) 0 0	2 (66.7) 1 (33.3) 0 0		0.230ª 0.0230
Dislocation	1 (2.4)	1 (2.4)	1	1
Drowning	0	2 (7.7)	0.9 (0.83–1.03)	0.07 ^a
Multiple injury	8 (19.5)	2 (7.7)	2.9 (0.57–14.94)	0.186

Abbreviations: CI, confidence interval; OR, odds ratio; PWE, patient with epilepsy.

Place of injury	PWE, N (%)		OR (95% CI)	p-Value
	Yes	No		
House School Street During sleep Workplace	30 (73.2) 5 (12.2) 2 (4.9) 2 (4.9) 2 (4.9)	6 (23.1) 1 (3.8) 15 (57.7) 0 1 (11.5	9.1 (2.9–28.6) 3.5 (0.4–31.6) 0.04 (0.01–0.19) 0.4 (0.06–2.53) 1.0 (0.89–1.04)	< 0.001 0.243 < 0.001 0.253 0.312
Recreational activities	0	1 (3.8)		0.206

Table 2 Place of occurrence of injuries among patients with and without epilepsy

Abbreviations: CI, confidence interval; OR, odds ratio; PWE, patient with epilepsy.

Table 3 Medical attention and hospitalization following injury

Variables	PWE N (%)	Non-PWE N (%)	<i>p</i> -Value
Medical attention	19 (46.3)	16 (61.5)	0.225
Hospitalization	7 (17.1)	5 (19)	0.822

Abbreviation: PWE, patient with epilepsy.

Table 4 Factors associated with physical injury in people with epilepsy

Factors	Physical injury in PWE		OR (95% CI)	p-Value
	Yes	No		
Type of epilepsy Generalized (motor) Focal Generalized (nonmotor) Unknown Combined generalized and focal	36 0 0 1 4	75 28 7 1 3	0.4 (0.08–1.69) 0.0 0.0 0.8 (0.03–17.51)	0.196 0.998 0.999 0.858
Number of seizures/years 0 1–12 13–24 25–36 > 36 Seizures with fall Use of > 1 ASM	4 30 5 1 1 106 20	76 30 4 2 2 49 135	0.1 (0.01-1.42) 2 (0.17-23.25) 2.5 (0.16-38.6) 1 (0.04-29.81) 8.6 (2.50-29.41) 2.6 (1.00-6.92)	0.09 0.580 0.512 1.0 0.001 0.049
Duration of illness from onset of seizure 0-2 3-5 6-10 > 11	30 14 10 5	12 33 39 12	0.1 (0.28–3.32) 1.0 (0.30–3.44) 0.6 (0.18–2.16)	0.960 0.997 0.448
Duration of illness from time of diagnosis 0-2 3-5 6-10 Compliance to ASMs Aura	15 16 10 13 11	31 46 37 28 30	1.8 (0.71–4.55) 1.3 (0.52–3.17) 0 0	0.220 0.583 0.998 0.999

Abbreviations: ASM, antiseizure medication; CI, confidence interval; OR, odds ratio; PWE, patient with epilepsy.

patients living with epilepsy. Our findings suggest that PWE were 1.8 times more likely to sustain a physical injury in 12 months, than those without epilepsy. Soft tissue injury was the only type of injury that was significantly high in PWE, and tongue bites accounted for this high level. The other types of injuries in PWE were head injury, burns, dental injury, fracture, and dislocation.

These results are similar to those of van den Broek et al in a multicentric study done in eight European countries, who reported that PWE were 1.5 times more at risk of injury and soft tissue injuries was the most common cause of injury.⁹ This observation is also in keeping with many other studies all over the world like in Turkey,¹⁰ Canada,¹¹ Iran,¹² India,¹³ and Ethiopia.¹⁴

Another important finding is that injuries in PWE occurred mostly at home, compared to patients without epilepsy, where injuries occurred on the streets. This finding is similar to that of van den Broek et al. However, this is in contrast with other reports where injuries in PWE occurred in equal proportions indoors, on the streets, and at the workplace. We believe that the reason for this discrepancy may be due to the fact that isolation due to stigma might cause the PWE to spend less time on the street and prefer to stay indoors.

Despite many reports in favor of a higher risk of injury in PWE, Téllez-Zenteno et al in 2008 in Canada found no significant difference in the rate of injury in PWE and the general population.¹⁵ It is possible that a difference in the methodology could explain the varying findings.

First, we carried out a hospital-based study while theirs was a population-based study where institutionalized patients, who could be at a substantial risk of injury, were excluded. Second, only injuries that interferes with normal daily activities were considered in their study whereas all injuries including those that did not need medical attention were included in ours.

We found that PWE were less likely to sustain a burn. This observation is in keeping with Josty et al who reported that there is a reduction in the incidence of burns due to epilepsy.¹⁶ However, the only cause of burn in our study was due to flames which is in contrast to Josty et al's report. In our setting, people use mostly fire for cooking. So if a seizure occurs while cooking there will be more chances that the patient sustains a burn either directly from the fire or boiling liquids from the pot. On the contrary, Josty et al's report was conducted in a developed country where technology is advanced and people mostly use gas stoves and plates at a considerable height that would less likely cause a burn in case the patient loses consciousness or fall because of a seizure.

We found that there was no case of drowning over the study period. It is possible that because of stigmatization, and isolation, PWE are not involved in social activities like swimming, and also very few people use bathtubs in our context.

Previous studies have reported that PWE had more hospitalization than those without epilepsy.^{9,15} This observation is in contrast to our findings, as we observed no increased rate of hospitalization or medical attention in PWE compared to those without epilepsy. The higher rate of hospitalizations in previous reports could be explained by the differences in methodologies, the other authors included PWE with injuries requiring medical attention or injuries causing disability, whereas we included all patients with injuries irrespective of whether they seek medical attention or not.

In a hospital-based retrospective controlled study conducted in Iran by Asadi-Pooya et al in 2012,¹² the authors reported that having seizures with falls was a significant risk factor of injury in PWE. This observation is similar to our findings, where a significant proportion of seizures with falls were associated with physical injuries. We also observed that the use of more than one antiseizure medication was associated with the occurrence of physical injuries. Lawn et al, in 2004, conducted a study on 247 Rochester residents and found that a greater number of antiseizure medications used was a significant risk factor of injury in epileptic patients.¹⁷ This result also correlates with our funding.

We found that the frequency of seizures and the type of seizure did not impact the incidence of physical injuries. This observation is in contrast to previous reports. This might be because the majority of our patients had less than one seizure per month, and though generalized seizures were more common, it occurred mostly indoors, and mostly in the night during sleep. The retrospective design of our study implies that there could be recall bias and some information might be incomplete. Some major physical injuries might be missed because of the design. The fact that this study was done in a tertiary hospital in an urban setting means the results might not be applicable to all the epilepsy population in our context; however, the center receives patients from all over the country and the subregion. The small sample size could also be a limitation to our study. We could extrapolate that our hospital is dealing with moderate to high epileptogenic cases and many low epileptogenic cases never reach the medical facility, as is seen in many developing countries.

Conclusion

In conclusion, here we have shown that the incidence of physical injuries in PWE at the Douala General Hospital is 25.5%, and is significantly higher than in patients without epilepsy. Soft tissue injuries and mild head injuries like abrasions and lacerations are significantly higher in PWE, while burns and road traffic accidents were less common in PWE. The risk of physical injury is increased in patients having seizures with falls and in those taking more than one antiseizure medication. Our results suggest the need that PWE, especially those on more than one medication and those having seizures with frequent falls should be educated on preventive measures on physical injuries. Despite its limitations in design, this is the first study on physical injuries in PWE in Cameroon and will serve as a baseline for further studies.

Conflict of Interest

C.-M. reported grants or contracts from NIH and Fogarty International; Leadership or fiduciary role in IATSIC and ASAP. G.M.D. reported support for attending meetings and/or travel from World Federation of Neurology (World Congress of Neurology 2019 travel grant) and International Brain Research Organization (Dakar Teaching Tools Workshop 2018 travel and accommodation); Leadership or fiduciary role in African Academy of Neurology (Secretary General term 2019 - 2021) and Epilepsy Alliance Africa (Vice President for West African Region). All other authors reported no conflict of interest.

Acknowledgments

We are grateful for the personal assistance of the personnel in charge of the archives of the Internal Medicine Department of the Douala General Hospital.

References

- 1 Epilepsy in the WHO African Region: Bridging the Gap. WHO Regional Office for Africa.2015
- 2 Dongmo L. Épilepsie au Sud-Cameroun: enquête préliminaire dans le village Bilomo. Bull Soc Path Exot 2000;93:266–267
- ³ Kuate C, Tchaleu B, Motah M, Nguefack S, Doumbe J. Situation de l'épilepsie au Cameroun. African Middle East Epilepsy J 2013;2(03):4–7
- 4 Mbonda NM, Kuate C, Nguefack S, et al. Itineraire Therapeutique Des Patients Epileptiques A Yaounde: A propos de 149 observations. Clin Mother Child Health 2008;5(02):893–898
- 5 Njamnshi AK, Yepnjio FN, Bissek ACZK, et al. A survey of public knowledge, attitudes, and practices with respect to epilepsy in Badissa village, centre region of Cameroon. Epilepsy Behav 2009; 16(02):254–259
- 6 Njamnshi AK, Bissek ACZK, Yepnjio FN, et al. A community survey of knowledge, perceptions, and practice with respect to epilepsy among traditional healers in the Batibo Health District, Cameroon. Epilepsy Behav 2010;17(01):95–102
- 7 Preux P-M, Tiemagni F, Fodzo L, et al. Antiepileptic therapies in the Mifi Province in Cameroon. Epilepsia 2000;41(04):432–439
- 8 Scheffer IE, Berkovic S, Capovilla G, et al. ILAE classification of the epilepsies: position paper of the ILAE Commission for Classification and Terminology. Epilepsia 2017;58(04):512–521

- 9 van den Broek M, Beghi ERESt-1 Group. Accidents in patients with epilepsy: types, circumstances, and complications: a European cohort study. Epilepsia 2004;45(06):667–672
- 10 Mollaoğlu M, Bolayir E. Injuries in patients with epilepsy and some factors associated with injury. Noro Psikiyatri Arsivi 2013; 50(03):269–273
- 11 Kwon CS, Liu M, Quan H, et al. The incidence of injuries in persons with and without epilepsy–a population-based study. Epilepsia 2010;51(11):2247–2253
- 12 Asadi-Pooya AA, Nikseresht A, Yaghoubi E, Nei M. Physical injuries in patients with epilepsy and their associated risk factors. Seizure 2012;21(03):165–168
- 13 Sapna CS, Sheeba SL, John J, Thomas SV. Accidents and injuries in people with epilepsy attending a tertiary care center in India. Epileptic Disord 2008;10(04):276–281
- 14 Bifftu BB, Tadesse Tiruneh B, Mekonnen Kelkay M, et al. Seizurerelated injuries among people with epilepsy at the outpatient department of the University of Gondar Hospital, Northwest Ethiopia: cross-sectional institutional-based study. Neurol Res Int 2017;2017:4970691
- 15 Téllez-Zenteno JF, Hunter G, Wiebe S. Injuries in people with selfreported epilepsy: a population-based study. Epilepsia 2008;49 (06):954–961
- 16 Josty IC, Narayanan V, Dickson WA. Burns in patients with epilepsy: changes in epidemiology and implications for burn treatment and prevention. Epilepsia 2000;41(04):453–456
- 17 Lawn ND, Bamlet WR, Radhakrishnan K, O'Brien PC, So EL. Injuries due to seizures in persons with epilepsy: a population-based study. Neurology 2004;63(09):1565–1570