



# Surgical Rhizarthrosis Treatment: Trapezius Resection Arthroplasty Associated with Tendon Interposition versus the Kuhns Technique

## *Tratamento cirúrgico da rizartrose: Artroplastia de ressecção do trapézio associada a interposição tendínea versus técnica de Kuhns*

Guilherme Henrique Teixeira Reis<sup>1</sup> Willker Galvão de Carvalho<sup>1</sup> Mauricio Foster Rodrigues<sup>1</sup>  
Jorge Raduan Neto<sup>1</sup> Aldo Okamura<sup>1</sup> João Carlos Belloti<sup>1</sup>

<sup>1</sup>Department of Hand Surgery, Hospital Alvorada, São Paulo, SP, Brazil

Address for correspondence: Guilherme Henrique Teixeira Reis, M.D., Passos, Minas Gerais, Brasil (e-mail: guilhermereis\_1@hotmail.com).

Rev Bras Ortop 2024;59(4):e572–e579.

### Abstract

**Objective** This study aimed to evaluate and compare the clinical and functional outcomes of two surgical procedures performed in patients with severe grade III and IV rhizarthrosis.

**Methods** We evaluated 39 patients who underwent two surgical techniques for rhizarthrosis treatment: trapeziectomy using the Kuhns technique or tendon interposition, with a minimum follow-up period of 6 months. The primary outcome assessment used the specific Trapeziometacarpal Arthrosis Symptoms and Disability (TASD) questionnaire, and the secondary outcome evaluation employed the shortened version of the Disabilities of the Arm, Shoulder, and Hand (QuickDASH) questionnaire and the Visual Analog Scale (VAS).

**Results** There was no statistically significant difference between groups in the TASD, QuickDASH, and VAS results, and both techniques demonstrated good functional and pain outcomes. No complication required a new surgical approach. We found a positive correlation between TASD and QuickDASH questionnaire scores, suggesting they are both effective in assessing functionality and disability in subjects with rhizarthrosis.

**Conclusion** Trapeziectomy using the Kuhns technique and tendon interposition proved effective in the surgical treatment of rhizarthrosis. There was no significant difference between the techniques concerning functional outcomes.

### Keywords

- ▶ orthopedic procedures
- ▶ osteoarthritis
- ▶ trapezium bone

Work carried out at the Hospital Alvorada, São Paulo, SP, Brazil.

received  
October 23, 2023  
accepted  
April 5, 2024

DOI <https://doi.org/10.1055/s-0044-1788289>  
ISSN 0102-3616.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

## Resumo

**Objetivo** O objetivo deste estudo é avaliar e comparar os resultados clínicos e funcionais de dois procedimentos cirúrgicos realizados em pacientes com rizartrose grave graus III e IV.

**Métodos** Avaliamos 39 pacientes submetidos a 2 técnicas cirúrgicas para o tratamento da rizartrose: trapeziectomia pela técnica de Kuhns ou com interposição tendínea, com mínimo de 6 meses de seguimento. O desfecho primário foi avaliado pelo questionário específico Sintomas e Incapacidade da Artrose Trapeziometacarpiana (Trapeziometacarpal Arthrosis Symptoms and Disability, T ASD, em inglês), e os desfechos secundários, pela versão abreviada do Questionário de Incapacidades do Braço, Ombro e Mão (Disabilities of the Arm, Shoulder, and Hand, QuickDASH, em inglês) e a Escala Visual Analógica (EVA).

**Resultados** Não houve diferença estatisticamente significantativa entre os grupos nos resultados do T ASD, QuickDASH e EVA e ambas as técnicas demonstraram bons resultados funcionais e da dor. Não houve complicações que necessitassem de nova abordagem cirúrgica. Verificamos uma correlação positiva entre as pontuações no T ASD e no QuickDASH, o que sugere que ambos são eficazes na avaliação da funcionalidade e incapacidade da rizartrose

**Conclusão** A trapeziectomia com técnica de Kuhns e com a interposição tendínea se mostraram efetivas no tratamento cirúrgico da rizartrose. Não houve diferença significativa entre as técnicas em relação aos resultados funcionais.

## Palavras-chave

- ▶ osso trapézio
- ▶ osteoartrite
- ▶ procedimentos ortopédicos

## Introduction

Rhizarthrosis is a progressive degenerative condition affecting the carpometacarpal joint of the thumb. It is one of the most common forms of osteoarthritis in the hands, and it is more frequent in women older than 50 years of age. This condition can be debilitating, limiting the subjects' daily living and work capacity.<sup>1</sup> Several therapeutic options are available to manage symptoms and improve quality of life, including lifestyle changes, motor and analgesic physical therapy, orthoses, pharmacological treatments, infiltrations, and surgery.<sup>2</sup>

Surgical treatment is an option for patients with rhizarthrosis who do not obtain symptom relief through nonsurgical therapies.<sup>2</sup> Trapeziectomy is a resection arthroplasty of the trapezium that was first described by Gervis, with good outcomes.<sup>3</sup> Some authors described the metacarpal's proximal migration and the compromised functional outcomes as complications of this technique.<sup>4</sup> Thus, to prevent these complications, techniques associated with trapeziectomy were described, including tendon interposition,<sup>5</sup> ligament reconstruction,<sup>6</sup> arthroplasty with implants,<sup>7</sup> the distraction hematoma formation technique (Kuhns technique),<sup>8</sup> ligament reconstruction with tendon interposition,<sup>6</sup> and acellular material interposition.<sup>5</sup> Although these associated techniques can be effective, they may increase the risk of other complications, such as infection, pain, implant loosening, and muscle strength loss.<sup>4</sup> The current literature has no studies with conclusive evidence regarding the most effective technique for the surgical treatment of rhizarthrosis.<sup>4</sup>

Therefore, this study aimed to evaluate two surgical techniques: trapeziectomy with the Kuhns technique and

with tendon interposition, in 39 patients, with a minimum follow-up period of 6 months.

These techniques were chosen due to the high number of patients who underwent these procedures to provide a comprehensive and representative analysis, allowing a significant comparison between them and contributing to validating the obtained results.

## Materials and Methods

This retrospective cohort study evaluated 39 patients. Inclusion criteria were the following: patients with clinical and imaging diagnosis of Eaton and Littler grade-III and -IV rhizarthrosis,<sup>9</sup> from both sexes, who underwent surgical treatment with trapeziectomy using the Kuhns technique<sup>8</sup> (group 1, n = 18), or trapeziectomy with tendon interposition<sup>5</sup> (group 2, n = 21), from 2018 to 2022, operated by four experienced hand surgeons, with a minimum follow-up period of 6 months.

Data were collected from the electronic records of the study's hospital, searching for patients with the following diagnosis codes of the International Classification of Diseases, Tenth Revision (ICD-10): M18.0, M18.1, and M19.9. Retrieved information included age, sex, operated side, dominant side, degree of osteoarthritis, and type of surgical procedure (– **Table 1**).

Exclusion criteria were the following: patients with rheumatological, traumatic, or neurological diseases affecting hand or wrist joints, those who underwent previous surgery in the thumb region, those lost to postoperative follow-up, and those who did not sign the informed consent form for the study.

**Table 1** Demographics of the patients

Data	Group 1	Group 2	Total	p-value
Mean age (in years)	62.4	58.0	60.1	0.056
Sex	94.5% F	90.5% F	92.4% F	0.052
	5.5% M	9.5% M	7.6% M	
Dominant side	94.5% R	81% R	87.2% R	0.921
	5.5% L	19% L	12.8% L	
Operated side	44.5% R	42.9% R	43.6% R	0.209
	55.5% L	57.1% L	56.4% L	
Arthrosis grade (Eaton and Littler)	67% III	72% III	70% III	0.748
	33% IV	28% IV	30% IV	
Questionnaire application time (in months)	Mean: 39.8	Mean: 21.6		
	Minimum: 8	Minimum: 6	Mean: 30	0,08
	Maximum: 62	Maximum: 50		

Abbreviations: III, grade III; IV, grade IV; F, female; L, left; M, male; R, right.

After inclusion, we invited patients for an in-person assessment in a single outpatient visit with four residents in hand surgery. We asked the patients to answer questionnaires about their clinical and functional outcomes, including the Trapeziometacarpal Arthrosis Symptoms and Disability (TASD),<sup>10</sup> the shortened version of the Disabilities of the Arm, Shoulder, and Hand (QuickDASH),<sup>11</sup> and the Visual Analog scale (VAS)<sup>12</sup> for pain, as shown in the **Annexes 1, 2, and 3**. The average time between the surgical procedure and the application of the questionnaires was 30 months.

For statistical analysis, we imported data to the IBM SPSS Statistics for MacOS (IBM Corp., Armonk, NY, USA) software, version 25.0. The descriptive statistics of categorical data included absolute and relative frequency. Continuous data underwent the Shapiro-Wilk normality test, and their description used mean  $\pm$  standard deviation (SD), median, and 25th and 75th percentiles. Data with parametric distribution underwent the Student *t* test for two independent samples, while those with nonparametric distribution underwent the Mann-Whitney test for two independent samples. A difference was statistically significant when the type-I error, that is, the *p*-value, was lower than 0.05.

## Surgical Procedures Performed and Evaluated in the Study

For the first group, we performed the surgical technique described by Kuhns<sup>8</sup> using a longitudinal dorsal approach of approximately 4 cm in the carpometacarpal joint of the thumb between the tendon of the abductor pollicis longus muscle and the tendon of the extensor pollicis brevis muscle. Next, we opened the joint capsule in a T shape to expose and remove the trapezius; fixation was performed under radio-scopic with two 1.5-mm Kirschner wires between the first

and second metacarpals, maintaining the carpometacarpal joint space (**Fig. 1**). The procedure ended with suturing the joint capsule and skin. Then, a sterile dressing and an antebrachioepalmar plaster splint, including the thumb, were applied and kept for 4 weeks. We removed the Kirschner wires in the outpatient clinic 4 weeks after the surgery.

As for the second group, a trapeziectomy and tenoarthroplasty technique was performed with the palmaris longus muscle,<sup>5</sup> using the same approach described for group 1. Then, we resected the tendon and the palmaris longus muscle using three approaches (**Fig. 2**), creating a ball, and interposing it between the scaphoid and metacarpal bones (**Figs. 3–4**). We closed the joint capsule (**Fig. 5**), sutured the incision, and applied a bandage and a plaster splint. The sutures were removed after 2 weeks and immobilization was maintained for 4 weeks.

## Clinical Outcomes


### Primary Outcome

The primary outcome was the TASD questionnaire,<sup>10</sup> which consists of a specific self-reported assessment of rhizarthrosis-related functional limitations. The TASD was translated and culturally adapted to Brazilian Portuguese in 2021.<sup>13</sup> It contains a series of questions about pain intensity and functional thumb capacity. The answers are scored from 0 to 100, with higher scores indicating higher dysfunction.


### Secondary Outcomes

The QuickDASH<sup>11</sup> assesses upper limb functional disability and pain, listing 11 related activities. The answers are scored from 0 to 100, and higher scores indicate greater functional disability and pain reported by the patients.

The VAS<sup>12</sup> assesses the intensity of pain reported by patients. It consists of a horizontal line with a scale from 0



**Fig. 1** Completed surgical procedure using the Kuhns technique. Intraoperative images demonstrating the space (red arrow) after trapezium bone resection and the two Kirschner wires (yellow arrow) used to fixate the first to the second metacarpal bones.



**Fig. 2** Image demonstrating the removal of the palmaris longus muscle for use as a graft. Palmaris longus muscle's musculotendinous graft (yellow arrow) removal from the three approaches.

to 10, in which 0 represents no pain, and 10, maximum pain. Patients were asked to mark the degree of pain they were feeling at that moment on the scale.

A single evaluator not linked to the study applied the questionnaires during outpatient visits to ensure the standardization of data collection.

The ethics committee of our institution approved the study under the CAAE number 71550023.1.0000.5487.

## Results

The average time to apply the questionnaire to patients was of 30 months postoperatively, with a minimum time of 8 and a maximum of 62 months for group 1, and a minimum time of 6 and a maximum of 50 months for group 2 (►Table 1).

In the analysis of the primary outcome, the mean TASD score was of  $25.2 \pm 27.5\%$  in the Kuhns technique group, and of  $24.9 \pm 22\%$  in the tendon interposition group (►Fig. 6).

In the analysis of the secondary outcomes, the mean QuickDASH score was of  $25.5 \pm 30.7\%$  in the Kuhns technique group and of  $31.6 \pm 24.6\%$  in the tendon interposition group (►Fig. 7).

As for pain, the mean VAS score was of  $3.2 \pm 3.2\%$  in the Kuhns technique group, and of  $3.0 \pm 2.7\%$  in the tendon interposition group (►Fig. 8).

We found a positive correlation between the TASD and QuickDASH scores: an increase in the score on one questionnaire corresponded to an increase in the score on the other questionnaire. This suggests that both tools are effective in measuring aspects related to functionality and disability in rhizarthrosis in a consistent manner (►Fig. 9).



**Fig. 3** Palmaris longus muscle graft before being placed at the trapezius site.



**Fig. 5** Joint capsule closure.



**Fig. 4** Graft placement at the site of the resected trapezium.


## Discussion

The study sample was homogeneous and representative, as described in the literature.<sup>1</sup> This study used the TASD questionnaire as the primary outcome, and the QuickDASH questionnaire and the VAS as secondary outcomes, since these tools assess function and potential limitations in daily living activities and pain in operated patients. The literature reports that these are the most appropriate tools to evaluate the effectiveness of surgical treatment.<sup>10,11</sup>


Our positive functional outcomes with the Kuhns technique<sup>8</sup> were consistent with those reported in the literature.<sup>5,8,14</sup> We observed as advantages of this technique, a shorter surgical time, the lack of need to make new incisions, and the exemption from tendon graft removal. These benefits simplify the procedure and result in a faster postoperative recovery, potentially reducing complications associated with additional incisions and grafting procedures. Its disadvantages are the potential complications inherent to the insertion and maintenance of the Kirschner wire and the need for a second procedure for pin removal, which occurred in the outpatient clinic of the institution in which the present study was conducted.

The technique of trapeziectomy with tenoarthroplasty of the palmaris longus muscle also seeks to avoid proximal migration of the first metacarpal bone by using the tendon graft as a biological spacer. The main advantage is the use of an autologous graft with no synthetic implant requirements. As the main disadvantages of this technique, we noted that, despite its performance on the same operated limb, it is limited for patients who lack the palmaris longus muscle and scarring complications from graft removal. The positive functional outcomes noted with the use of this technique in the present study are consistent with those of the literature, in which several authors<sup>5,15</sup> have reported good and excellent outcomes.

Comparing both surgical techniques, we observed that trapeziectomy with long palmar muscle tendon interposition was not superior to the Kuhns technique. Although we did not find studies specifically comparing both procedures, the outcomes were consistent with the literature addressing similar surgical techniques.<sup>16-19</sup> The findings of this review support the effectiveness of both approaches in rhizarthrosis treatment.




**Fig. 6** Analysis of the T ASD scores in patients who underwent the Kuhns or tendon interposition techniques.



**Fig. 7** Analysis of the QuickDASH scores in patients who underwent the Kuhns or tendon interposition techniques.

However, based on the Cochrane Review,<sup>20</sup> we cannot currently make recommendations regarding the superiority of any surgical procedure over another for this condition.

We observed a positive correlation between the T ASD and QuickDASH, a result consistent with the findings in the literature.<sup>10,21</sup>



**Fig. 8** Analysis of the VAS scores in patients who underwent the Kuhns or trapeziectomy + tendon interposition techniques.

## Conclusion

The two techniques evaluated proved effective for treating patients with rhizarthrosis per the T ASD, with an average postoperative follow-up of 30 months. There was no superiority in functional outcomes between the groups when comparing trapeziectomy techniques with tendon interposition or distraction. The specific T ASD and the generic


QuickDASH functional questionnaires proved equivalent to measuring the patients' degree of functional limitation.

### Financial Support

The authors declare that they have not received financial support from agencies in the public, private or non-profit sectors to conduct the present study.

### Conflict of Interests

The authors have no conflicts of interests to declare.



**Fig. 9** Correlation of the QuickDASH and T ASD scores in postoperative patients who underwent the Kuhns or tendon interposition techniques.

## References

- 1 Armstrong AL, Hunter JB, Davis TR. The prevalence of degenerative arthritis of the base of the thumb in post-menopausal women. *J Hand Surg [Br]* 1994;19(03):340–341
- 2 Bertozzi L, Valdes K, Vanti C, Negrini S, Pillastrini P, Villafañe JH. Investigation of the effect of conservative interventions in thumb carpometacarpal osteoarthritis: systematic review and meta-analysis. *Disabil Rehabil* 2015;37(22):2025–2043
- 3 Gervis WH. Excision of the trapezium for osteoarthritis of the trapezio-metacarpal joint. *J Bone Joint Surg Br* 1949;31B(04):537–539, illust
- 4 Conolly WB, Lanzetta M. Surgical management of arthritis of the carpo-metacarpal joint of the thumb. *Aust N Z J Surg* 1993;63(08):596–603
- 5 Dell PC, Muniz RB. Interposition arthroplasty of the trapeziometacarpal joint for osteoarthritis. *Clin Orthop Relat Res* 1987;(220):27–34
- 6 Burton RI, Pellegrini VD Jr. Surgical management of basal joint arthritis of the thumb. Part II. Ligament reconstruction with tendon interposition arthroplasty. *J Hand Surg Am* 1986;11(03):324–332

- 7 Pellegrini VD Jr, Burton RI. Surgical management of basal joint arthritis of the thumb. Part I. Long-term results of silicone implant arthroplasty. *J Hand Surg Am* 1986;11(03):309–324
- 8 Kuhns CA, Meals RA. Hematoma and distraction arthroplasty for basal thumb osteoarthritis. *Tech Hand Up Extrem Surg* 2004;8(01):2–6
- 9 Eaton RG, Glickel SZ. Trapeziometacarpal osteoarthritis. Staging as a rationale for treatment. *Hand Clin* 1987;3(04):455–471
- 10 Becker SJ, Teunis T, Ring D, Vranceanu AM. The Trapeziometacarpal Arthrosis Symptoms and Disability Questionnaire: Development and Preliminary Validation. *Hand (N Y)* 2016;11(02):197–205
- 11 Beaton DE, Wright JG, Katz JN Upper Extremity Collaborative Group. Development of the QuickDASH: comparison of three item-reduction approaches. *J Bone Joint Surg Am* 2005;87(05):1038–1046
- 12 Huskisson EC. Measurement of pain. *Lancet* 1974;2(7889):1127–1131
- 13 de Souza Almeida VA, Fernandes CH, Meireles LM, Faloppa F, Ejnisman B, Cohen M. Translation and cross-cultural adaptation of “Trapeziometacarpal Arthrosis Symptoms and Disability-TASD” into Brazilian Portuguese. *Adv Rheumatol* 2021;61(01):61
- 14 Kuhns CA, Emerson ET, Meals RA. Hematoma and distraction arthroplasty for thumb basal joint osteoarthritis: a prospective, single-surgeon study including outcomes measures. *J Hand Surg Am* 2003;28(03):381–389
- 15 Berkhout MJ, Bachour Y, Wessing D, Ritt MJPF. Distal Pole Resection of the Scaphoid for the Treatment of Scaphotrapeziotrapezoid Osteoarthritis. *Hand (N Y)* 2019;14(02):230–235
- 16 Davis TR, Brady O, Dias JJ. Excision of the trapezium for osteoarthritis of the trapeziometacarpal joint: a study of the benefit of ligament reconstruction or tendon interposition. *J Hand Surg Am* 2004;29(06):1069–1077
- 17 Jain A, Herrera FA. Cost Analysis and National Trends in the Treatment of Thumb Basal Arthritis: Comparing Ligament Reconstruction/Tendon Interposition and Trapeziectomy/Hematoma Distraction Arthroplasty. *Ann Plast Surg* 2021;86(6S, Suppl 5):S622–S624
- 18 Tolo ET. Ligament reconstruction and tendon interposition versus trapeziectomy and hematoma distraction arthroplasty for treatment of trapeziometacarpal arthritis. *Curr Opin Orthop* 2006;17:283–287
- 19 Sandvall BK, Cameron TE, Netscher DT, Epstein MJ, Staines KG, Petersen NJ. Basal joint osteoarthritis of the thumb: ligament reconstruction and tendon interposition versus hematoma distraction arthroplasty. *J Hand Surg Am* 2010;35(12):1968–1975
- 20 Wajon A, Vinycomb T, Carr E, Edmunds I, Ada L. Surgery for thumb (trapeziometacarpal joint) osteoarthritis. *Cochrane Database Syst Rev* 2015;2015(02):CD004631
- 21 Ratneswaran A, Rockel JS, Antflek D, et al. Investigating Molecular Signatures Underlying Trapeziometacarpal Osteoarthritis Through the Evaluation of Systemic Cytokine Expression. *Front Immunol* 2022;12:794792