

Synovial Chondromatosis of the Shoulder: Report of Two Cases

Condromatose sinovial do ombro: Relato de dois casos

Daniel Ferreira Fernandes Vieira^{1,2®} Beatriz Begnini Borsato^{3®} Luís Felipe Baldinu Caramujo^{3®} Paula de Andrade Castello^{3®} João Paulo Fernandes Guerreiro^{1,2®}

¹ Orthopedics Course, Pontifícia Universidade Católica do Paraná (PUC/PR), Londrina, PR, Brazil

²Hospital de Ortopedia Uniort.e, Londrina, PR, Brazil

³ School of Medicine, Pontifícia Universidade Católica do Paraná (PUC/PR), Londrina, PR, Brazil Address for correspondence Daniel Ferreira Fernandes Vieira, MSc, Hospital de Ortopedia Uniort.e, Avenida Higienópolis, 2600, Guanabara, 86050-000, Londrina, PR, Brazil (e-mail: danielffvieira@hotmail.com).

Rev Bras Ortop 2024;59(S2):e180-e183.

Abstract

Keywords ► arthroscopy

► shoulder

synovial

chondromatosis

synovial membrane

Synovial chondromatosis is an uncommon, progressive, benign condition favoring synovial metaplasia resulting from the production of cartilaginous tissue as loose bodies within the joints. In rare cases, it can affect the interior of the shoulder joint and present with pain, edema, and impaired mobility. The diagnosis is challenging, requiring imaging techniques. Its confirmation often occurs only after surgical treatment and anatomopathological examination. The authors report two cases of patients with similar clinical presentations of pain, edema, and mobility loss in the shoulder joint. The investigation included imaging exams, such as radiographs and magnetic resonance imaging, and an anatomopathological examination confirming the diagnostic hypothesis. Arthroscopic surgical treatment with synovectomy and removal of loose bodies followed by physical therapy rehabilitation resulted in clinical improvement in both patients. This report emphasizes the importance of investigating suspected synovial chondromatosis due to its nonspecific clinical presentation. Comparing our outcomes with the literature, we concluded that surgical treatment with synovectomy, loose body removal, and physical therapy is effective, and that longterm outpatient follow-up is necessary to detect recurrence.

Work carried out at the Hospital de Ortopedia Uniort.e, Londrina,

PR, Brazil.

received June 18, 2021 accepted October 14, 2021 DOI https://doi.org/ 10.1055/s-0044-1790596. ISSN 0102-3616. © 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (https://creativecommons.org/licenses/by/4.0/). Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Resumo

A condromatose sinovial é uma afecção benigna incomum de caráter progressivo que favorece a metaplasia sinovial com a produção de tecido cartilaginoso em forma de corpos livres, encontrados nas articulações. Em casos raros, pode afetar o interior da articulação do ombro e apresentar-se com dor, edema e prejuízo da mobilidade. O diagnóstico é desafiador, requerendo técnicas de imagem e muitas vezes confirmado apenas após o tratamento cirúrgico e exame anatomopatológico. Os autores relatam dois casos de pacientes com quadros clínicos semelhantes de dor, edema e perda de mobilidade na articulação do ombro. A investigação contou com exames de imagem como radiografias e ressonância magnética, além do exame anatomopatológico, que confirmou a hipótese diagnóstica. O tratamento cirúrgico artroscópico com sinovectomia e remoção dos corpos livres, seguido de reabilitação fisioterapêutica, resultou na melhora clínica de ambas as pacientes. Com esse relato, os autores ressaltam a importância da investigação na suspeita de condromatose sinovial, devido ao seu quadro clínico inespecífico. Comparando os resultados obtidos com os propostos na literatura, concluímos que o tratamento cirúrgico com sinovectomia e remoção de corpos livres, associado a fisioterapia, é eficaz, e que o seguimento ambulatorial a longo prazo é necessário para detecção de recidiva.

Palavras-chave ► artroscopia

- condromatose sinovial
- membrana sinovial
- ► ombro

Introduction

Synovial chondromatosis (SC) is an exceptional arthropathy, usually monoarticular, resulting from the proliferation and metaplasia of synovial tissue and the formation of loose cartilaginous bodies in the tendon sheath or joint spaces.¹⁻³ The involvement pattern includes diarthrodial joints, especially the knee, hip, and elbow; descriptions of shoulder involvement are scarce in the literature.² SC etiology remains unknown, and the highest SC incidence occurs in the third to fifth decade of life. In addition, SC is three times more common in men.³ The clinical picture consists of pain, crepitus, edema, and joint movement limitation,² often with no apparent cause.¹ Malignant transformation is uncommon, and there is no direct relationship with trauma or inflammatory processes.⁴ Clinical diagnosis is difficult, as history and physical examination findings are not specific. Therefore, imaging methods, including radiography, computed tomography, and magnetic resonance imaging (MRI), become essential to identify the different types of SC lesions and stages.⁵ Diagnostic confirmation relies on histological examination of the synovial tissue, and surgery is the treatment of choice for symptomatic patients.^{1,3}

Case Reports

Case 1: A 62-year-old female patient with a history of pain in her right shoulder after exertion. On examination, the shoulder presented mild edema and limited movement. The Neer impingement test and the Jobe and Speed tests were positive. Radiographs showed signs of impingement and calcifications. MRI revealed a rupture of the supraspinatus tendon and multiple calcifications in the bursa measuring 0.4 to 1.2 cm (**Figs. 1** and **2**). The patient underwent video arthroscopic surgical treatment consisting of synovectomy, chondroma removal, and supraspinatus lesion repair. The anatomopathological examination confirmed synovial chondromatosis. The patient started physical therapy two weeks after surgery. During the outpatient follow-up, she presented improvement in pain, range of motion, and strength in her right shoulder.

Case 2: A 56-year-old female patient with a history of pain in her right shoulder. The physical examination revealed mild edema, limited movement, and a positive Jobe test. Radiographs showed signs of impingement and a hooked acromion with a spur. MRI revealed intense proliferative synovitis with small intra-articular nodules in the axillary



Fig. 1 Magnetic resonance imaging of the shoulder (coronal view, T2-weighted image).



Fig. 2 Magnetic resonance imaging of the shoulder (axial view, T2-weighted image)



Fig. 3 Intra-articular arthroscopic view.

recesses, bursitis, supraspinatus tendon rupture, extensive partial tear of the subscapularis tendon, and ruptured tendon of the long head of the biceps. The patient underwent video arthroscopic treatment for supraspinatus tendon injury repair, synovectomy, subacromial decompression, and removal of multiple loose bodies (**-Figs. 3**, **4**, and **5**). The anatomopathological examination confirmed synovial chondromatosis. The patient began physical therapy rehabilitation one month after surgery. Pain improved immediately after surgery, and shoulder function improved after six months of rehabilitation.

Discussion

Shoulder CS is a rare report in the literature (<5%) due to its atypical location.³ However, SC is a condition with



Fig. 4 Subacromial arthroscopic view.

well-defined characteristics. The classification into primary and secondary forms considers cartilaginous body number, shape, and size or the presence of a pre-existing disease. In shoulders, the secondary form is the most common, incidentally diagnosed in tests for a previous underlying disease.⁵ Although CS is more common in men, we reported two female patients. The monoarticular characteristic is consistent with the literature. In both cases, CS had an association with rotator cuff syndrome.

The cartilaginous bodies originating from synovial metaplasia³ may increase in size and undergo calcification^{3,4} or endochondral ossification, causing joint erosions, pain, stiffness, and movement restriction³ Therefore, imaging tests are essential, and their findings depend on the stage of the disease. Our patients underwent shoulder radiography and MRI, which detected loose bodies. In both cases, the MRI showed rupture of the supraspinatus tendon. Rotator cuff injury may result from the persistent presence of loose bodies in the subacromial region and impingement.⁶

Although there is controversy about the best therapy for SC^{2,7–9} and reports of spontaneous remission,⁷ most of the literature supports surgical treatment.^{1,2,9} Most cases under conservative therapies remain symptomatic or present with symptoms worsening before surgery.⁹ Everything indicates that arthroscopic treatment is the gold standard,⁹ but the need for synovectomy is not yet well established.^{2,8,9} Disease recurrence has been reported since SC can affect the tendon sheath of the long head of the biceps and escape detection or due to its incomplete treatment with an isolated arthroscopic technique by not using arthrotomy or mini-open techniques when necessary.9 Several authors described loose bodies removal with synovectomy.⁸ However, Jeffreys (1967) concluded that only removing loose bodies was successful.¹⁰ Milgram (1977) apud Maurice et al.⁸ (1988) recommended synovectomy with free body removal for the initial SC stage and the isolated removal of these bodies in the late stage.

Some authors, such as Ramos et al.⁷ (1997), prefer the simple removal of articular free bodies. However, when these



Fig. 5 Macroscopic appearance of the removed chondromas.

bodies are close to the synovium, as in our cases, we propose the addition of arthroscopic synovectomy to increase procedural precision. Arthroscopy involves small incisions, allows assessing the entire glenohumeral joint, and facilitates rapid rehabilitation.^{1,2} We believe that synovectomy and removing all loose bodies is the best therapy for shoulder CS, and resection of the bursa with the nodules in the subacromial region can minimize the occurrence of future rotator cuff injuries.⁶ There are reports of recurrence when the synovial membrane is not excised,³ favoring malignant transformation to synovial chondrosarcoma, although this event is rare.^{1,3} The association of radiotherapy with treatment is questioned because there is little benefit in its use since metastasis developed by patients with previous CS is rare.¹

Rehabilitation with physical therapy is critical for shoulder function recovery and provides the good outcomes described in the literature.⁹ In both cases, there was an improvement in pain and glenohumeral joint mobility. Long-term outpatient monitoring of these subjects is valuable since we must not neglect a potential recurrence. We should consider evaluation with imaging tests every 2 or 3 years for SC treatment.^{2,5,8}

Financial Support

The authors declare that there was no financial support from public, commercial, or not-for-profit sources.

Conflicts of Interest The authors have no conflicts of interest.

References

- 1 Brasil Filho R, Filardi Filho CS, Menitti EL, Baptista MV, Daher SS. Condromatose sinovial: relato de um caso. Rev Bras Ortop 1997; 32(11):921–923
- 2 Paim AE, Ferreira DC, Paim A, Almeida RM. Tratamento artroscópico da condromatose sinovial do ombro: relato de caso. Rev Bras Ortop 2008;43(04):146–149
- 3 Acharya BM, Devkota P, Shrestha SK, Pradhan NS, Ahmad S. Condromatose sinovial simétrica bilateral do ombro: relato de caso. Rev Bras Ortop 2018;53(05):647–650
- 4 Fuerst M, Zustin J, Lohmann C, Rüther W. Synoviale Chondromatose. Orthopade 2009;38(06):511–519
- 5 McKenzie G, Raby N, Ritchie D. A pictorial review of primary synovial osteochondromatosis. Eur Radiol 2008;18(11):2662–2669
- 6 Horii M, Tamai M, Kido K, Kusuzaki K, Kubo T, Hirasawa Y. Two cases of synovial chondromatosis of the subacromial bursa. J Shoulder Elbow Surg 2001;10(02):186–189
- 7 Ramos MRF, Ramos RRM, Santos LA. Condromatose sinovial. Relato de caso. Rev Bras Ortop 1997;32(09):749–750
- 8 Maurice H, Crone M, Watt I. Synovial chondromatosis. J Bone Joint Surg Br 1988;70(05):807–811
- 9 Sanchez-Munoz E, Prado MA, Martinho G, Pérez YG, Miró RL. Condromatose sinovial do ombro. Descrição de dois casos e revisão da literatura. Rev Port Ortop Traumatol 2014;22(03):406–414
- 10 Jeffreys TE. Synovial chondromatosis. J Bone Joint Surg Br 1967;49 (03):530–534