

Refractory Gastrointestinal Bleeding due to Gastric Neuroendocrine Tumor Treated with Application of Hemostatic powder

Sanish Ancil¹ Sachin Hosahally Jayanna¹ Sai Vamshi Krishna Chigurupati¹ Ritesh Acharya¹
Rajesh Gupta¹ Surinder S. Rana¹

¹ Departments of Gastroenterology, and *GI Surgery HPB and Liver Transplantation, Postgraduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

Address for correspondence Prof Surinder S. Rana, MD, DM, FASGE, AGAF, Department of Gastroenterology Postgraduate Institute of Medical Education and Research (PGIMER), Chandigarh 160012, India (e-mail: rana.surindersingh@pgimer.edu.in).

J Digest Endosc 2024;15:263–264.

Abstract

Keywords

- endoscopy
- neuroendocrine tumor
- hemoclip

Hemostatic powder is a new hemostatic modality that is easy to use and covers a large surface area. It causes hemostasis by mechanical tamponade over the bleeding vessel and increases the concentration of the clotting factor. In this image, we report a case of gastric neuroendocrine bleeding presenting with refractory bleeding and hemostasis achieved with the endoscopic application of hemostatic powder.

A 64-year-old gentleman, postcoronary artery bypass graft surgery and on antiplatelet therapy, presented with hematemesis and low hemoglobin (10.6 g/dL). Emergency gastroscopy revealed an umbilicated lesion measuring 1.5 cm in the body of the stomach with an active ooze of blood (► **Fig. 1**). Hemostasis was achieved with combined hemoclip application and adrenaline saline injection. However, he rebled 24 hours later and computed tomography angiography of the abdominal vessels did not reveal any abnormality. Repeat gastroscopy revealed persistent oozing of blood from the lesion despite the presence of hemoclips (► **Fig. 2**). A hemostatic powder (Hemospray; Cook Medical, Winston-Salem, North Carolina, United States) was sprayed over the lesion and hemostasis was achieved (► **Fig. 3**). Thereafter, an endoscopic biopsy was taken from the edge of the lesion and histopathological examination revealed a grade II neuroendocrine tumor. There were no further episodes of gastrointestinal bleeding and a week later, the patient underwent a successful surgical resection of the gastric lesion (► **Fig. 4**).

Hemostatic powder is a new hemostatic modality that is easy to use and covers a large surface area. It causes



Fig. 1 Endoscopy: Actively bleeding umbilicated lesion measuring 1.5 cm in size in the body of the stomach.

hemostasis by mechanical tamponade over the bleeding vessel and increases the concentration of the clotting factors.¹ It is effective in achieving hemostasis in bleeding ulcers, varices, and cancer-related bleeding.¹




Fig. 2 Endoscopy: Persistent oozing of the blood from the lesion despite the presence of hemoclips.




Fig. 3 Hemospray sprayed over the lesion and hemostasis achieved.

Authors' Contributions

S.A. drafted the manuscript and was involved in data collection. S.H.J. was involved in critical revision of the manuscript for intellectual content. S.V.K.C., R.A., and R.G. were involved in data collection. S.S.R. was involved in




Fig. 4 Surgically resected lesion (arrow).

collection and interpretation of data and critical evaluation of the manuscript for intellectual content.

Funding

None.

Conflict of Interest

None declared.

Reference

- 1 Aziz M, Weissman S, Mehta TI, et al. Efficacy of Hemospray in non-variceal upper gastrointestinal bleeding: a systematic review with meta-analysis. *Ann Gastroenterol* 2020;33(02):145–154