



Treatment Options for Intracranial Epidural Hematoma - An Integrative Review of the Past Three Decades

Opções de tratamento para hematoma epidural intracraniano - Uma revisão integrativa das últimas três décadas

Carlos Umberto Pereira¹ Lauro Roberto de Azevedo Setton²

¹Neurosurgery Service, Hospital de Urgências de Sergipe (HUSE), Aracaju, SE, Brazil

²Otorhinolaryngology Service, Hospital Santo Antônio Obras Irmã Dulce, Salvador, BA, Brazil

Address for correspondence Carlos Umberto Pereira, MD, PhD, Neurosurgery Service, Hospital de Urgências de Sergipe (HUSE), Av. Augusto Maynard, 245/404. Bairro São José, CEP: 49015-380 Aracaju, SE, Brazil (e-mail: umberto@infonet.com.br).

Arq Bras Neurocir 2024;43(4):e308–e314.

Abstract

Keywords

- ▶ acute intracranial epidural hematoma
- ▶ head trauma
- ▶ treatment
- ▶ craniotomy
- ▶ conservative treatment
- ▶ endovascular embolization

Resumo

Introduction Acute epidural intracranial hematoma (IEH) has been considered one of the most relevant neurosurgical emergencies in recent decades due to its high potential for morbidity and mortality. Early diagnosis followed by appropriate treatment results in a more favorable prognosis considering its rapid progression.

Objective To describe the various treatment modalities for IEH in the last three decades and their updates.

Methods Integrative literature review on therapeutic options in IEH treatment. The terms “Epidural hematoma,” “Traumatic brain injury,” and “treatment” were used in the Medline/PubMed, Google Scholar, and SciELO platforms, resulting in 90 articles.

Results Appropriate treatment for IEH depends directly on the Glasgow Coma Scale score obtained during admission, bleeding location, lesion size, presence of associated intracranial injuries, and the neurosurgeon’s experience.

Conclusion Initial treatment for IEH is predominantly surgical, with conservative treatment indicated in selected cases. Both neurosurgeons and clinicians must identify characteristic signs and symptoms promptly to avoid treatment delay. Moreover, minimally invasive approaches have gained prominence in recent decades, associated with image-guided procedures, and when well-indicated, result in rapid recovery and lower morbidity.

Introdução O hematoma epidural intracraniano agudo (HEIA) tem sido considerado uma das emergências neurocirúrgicas de maior relevância das últimas décadas devido ao seu alto potencial de morbimortalidade. O diagnóstico precoce seguido de

received
February 15, 2024
accepted
October 18, 2024

DOI <https://doi.org/10.1055/s-0044-1796652>.
ISSN 0103-5355.

© 2024. Sociedade Brasileira de Neurocirurgia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

tratamento adequado resulta em prognóstico mais favorável, tendo em vista sua progressão galopante.

Objetivo Descrever as diversas modalidades de tratamento do HEIA nas últimas três décadas e suas atualizações.

Métodos Revisão integrativa de literatura sobre as opções terapêuticas no tratamento do HEIA. Foram utilizados os termos: “Hematoma epidural”, “Traumatismo cranioencefálico” e “tratamento” nas plataformas Medline/PubMed, Google Scholar e SciELO, resultando em 90 artigos.

Resultados O tratamento adequado para o HEIA, depende diretamente do escore na escala de coma de Glasgow obtido durante a admissão, localização do sangramento, tamanho da lesão, presença de lesões intracranianas associadas e da experiência do neurocirurgião.

Conclusão O tratamento inicial para o HEIA é eminentemente cirúrgico. Sendo o tratamento conservador indicado em casos selecionados. É de suma importância que tanto neurocirurgiões quanto clínicos devam saber identificar os sinais e sintomas característicos para não retardar o tratamento. Além disso, a abordagem minimamente invasiva vem ganhando notoriedade nas últimas décadas, associada aos procedimentos guiados por imagem, e, quando bem indicadas, resultam numa rápida recuperação e menor morbidade.

Palavras-chave

- ▶ hematoma epidural agudo intracraniano
- ▶ traumatismo cranioencefálico
- ▶ tratamento
- ▶ craniotomia
- ▶ tratamento conservador
- ▶ embolização endovascular

Introduction

Intracranial epidural hematoma is defined as a collection of blood located between the dura mater and the cranial bones.^{1,2} Acute intracranial epidural hematoma (IEH) accounts for 0.2% to 6% of all traumatic brain injuries (TBIs) and 9% to 12% of severe TBIs.²⁻⁶ It predominantly affects the second and third decades of life.^{3,7-9} In 70% to 90% of cases, it is due to motor vehicle accidents and accidental falls.^{3,7,9,10} Its location is supratentorial in 90% and infratentorial in 10% of cases. Approximately 95% are unilateral and 5% are bilateral.^{9,11}

The classic clinical picture of loss of consciousness after TBI followed by a lucid interval, accompanied by homolateral mydriasis to the hematoma and contralateral hemiparesis, with a decreased level of consciousness, occurs in only 20% of cases.² CT is the method of choice for initial diagnosis and treatment.² Treatment varies according to the size, location, evolution, neurological status of the patient, and findings on neuroimaging exams. The authors discuss the various means of invasive and non-invasive treatment of IEH.

Methods

Justification: IEH has been frequent in cases of traumatic brain injury (TBI). Various procedures have been performed with excellent results. **Objective:** To describe the various modalities of IEH treatment. **Methods:** Integrative literature review on therapeutic options in the treatment of IEH. The terms “Epidural hematoma,” “Traumatic brain injury,” and “treatment” were used in the Medline/PubMed, Google Scholar, and SciELO platforms, resulting in 90 articles selected based on inclusion and exclusion criteria, as well as their respective citation impact and content. **Inclusion criteria:**

Articles written in Portuguese, English, and Spanish, published in the last three decades (1980-2023), absence of tangential discussion of the topic. **Exclusion criteria:** Articles published in languages other than Portuguese, English, and Spanish, publications outside the specified time frame, and tangential discussions of the topic. **Conclusion:** The treatment of IEH depends on a series of factors such as Glasgow Coma Scale score on admission, size, location, presence of associated intracranial lesions, findings of neuroimaging exams, and the experience of the neurosurgical team.

Discussion

IEH is a lesion that requires neurosurgical intervention in most cases. When it reaches a certain volume, it can cause elevated intracranial pressure (ICP) and lead the patient to a state of coma or cerebral herniation, and in some cases, death. Craniotomy has been the most indicated procedure for cerebral decompression.¹ The treatment of IEH ranges from conservative treatment to endovascular therapy, depending on the clinical picture and findings in neuroimaging exams. The main treatment options for IEH include conservative management, immediate surgery, exploratory craniotomy, conventional surgery, decompressive craniectomy, arterial embolization, ultrasound or computed tomography (CT) guided aspiration, and endoscopic drainage.^{2,12-22}

Conservative Treatment

Several factors are reported to influence the strategic approach to IEH treatment. Selection criteria for conservative versus surgical treatment remain controversial.^{12,23} The decision for conservative treatment and the timing for delayed intervention can be made on an individual basis

and depend on parameters such as patient age, size and location of the hematoma, neurological status of the patient upon admission, and case evolution.^{2,24} Shahid et al.²⁵ observed that young patients who underwent early surgery with no or minimal associated brain injury recovered better than those who underwent surgery late.

Conservative treatment is indicated in patients with a preserved level of consciousness, without focal neurological deficit, absence of associated intracranial lesion, and CT showing IEH volume below <30 ml, thickness below 15 mm, and midline shift below 5 mm six hours or more after trauma, but with constant clinical observation and CT monitoring; in case of neurological decompensation, immediate surgery is indicated. Bullock et al.¹³ demonstrated that a volume between 12–38 ml was suitable for conservative treatment, but Chen et al.¹² suggest that hematoma larger than 30 ml, thickness greater than 15 mm, and midline shift greater than 5 mm constitute a strong indication for surgical drainage. There is still disagreement regarding the management of IEH located in the posterior fossa; several authors indicate surgical treatment in all cases due to the potential for a considerable mass effect in a small space. Wong²⁶ reported that IEH located in the posterior fossa, with a volume smaller than 10 ml, can undergo conservative treatment. In cases with associated intracranial lesions, it indicates greater severity of trauma, and conservative treatment is contraindicated.^{8,27,28} Samadi-Motlagh et al.²⁹ used tranexamic acid in selected IEH patients for conservative treatment and observed a decrease in hematoma volume expansion compared to the control group that did not use the medication, concluding that tranexamic acid plays an important role in bleeding control in IEH patients and improves their prognosis.

An important consideration should be made regarding the timing of repeat CT scans in cases of conservative treatment. Hematoma volume increases in 23% of cases, beginning eight hours after trauma and completing within 36 hours.^{30–32} Spontaneous resolution rarely occurs in cases undergoing conservative treatment.^{33–35}

Conventional Surgery

Surgical treatment is indicated based on neurological status and head CT findings (► **Table 1**):

Surgical treatment is performed through osteoplastic craniotomy above the site of the hematoma, with coagulation of the lacerated vessel always necessary. Dura mater suturing at the edges of the craniotomy and in the center of the bone

Table 1 Indications for urgent surgery in cases of intracranial epidural hematoma

1. Coma with anisocoria and CT demonstrating IEH necessitate urgent surgery.
2. Coma and worsening neurological status in IEH with a volume > 25 ml.
3. Volume of IEH > 30 ml, even in the absence of symptoms.
4. Volume of IEH > 25 ml, located in the posterior fossa or temporal region.
5. Midline shift > 4 mm, with worsening neurological status.
6. Increase in volume of the IEH.

flap to prevent its recurrence. In cases of bleeding from venous sinuses, control is achieved through tamponade using gelfoam or surgicel, and the patient's head is elevated in bed to prevent air embolism.^{10,13,36,37} It is important to emphasize that drains should not be placed in postoperative cases of IEH, as they may lead to local infection and have not been effective in cases of IEH recurrence.

Immediate Surgery and Exploratory Trepanation

In cases where a patient presents in the emergency room with unilateral pupillary dilation, decerebrate posture, and signs of elevated intracranial pressure, emergency action may involve making a small incision in the suspected area above the hematoma. A quick trephine hole is made, allowing for partial drainage of the hematoma to relieve intracranial hypertension, followed by craniotomy, definitive hematoma drainage, and coagulation of the injured vessel.^{14,38–40} In the presence of associated intracranial lesions (subdural hematoma, brain contusion, intracerebral hematoma) seen on CT scan, if necessary, surgical drainage is performed. Intraoperative ultrasound may sometimes be useful to identify deep lesions. In cases of decompressive craniectomy (DC) where the bone flap cannot be immediately repositioned, it should be preserved in the freezer or the fatty layer of the abdominal wall for later repositioning. This occurs when there is cerebral edema or an existing lesion seen on CT observed during surgery, or in cases of refractory intracranial hypertension developed perioperatively. A new CT scan is indicated to determine the extent of hematoma drainage and to identify late-evolving hematomas.

Decompressive Craniectomy

Considerable controversy exists regarding the initial treatment of IEH, especially the optimal surgical treatment option, particularly in patients with cerebral herniation. The recommended treatment for IEH is osteoplastic craniotomy and hematoma drainage with replacement of the bone flap. Experiences with decompressive craniectomies (DC) in IEH have been sparsely reported in the literature.^{41–43} Korde et al.¹⁵ reported their experience in ten cases of IEH with admission Glasgow Coma Scale (GCS) scores below 5, who underwent DC initially, and concluded that these patients showed better outcomes.

Other authors^{2,44,45} recommend DC as the first choice in cases of massive hematoma and low GCS, and it has been beneficial. In some cases, hematoma drainage with additional DC may effectively prevent and/or alleviate postoperative cerebral infarction; however, further investigation through randomized clinical trials is needed for confirmation. Korde et al.¹⁵ suggest DC in patients with admission GCS scores below 5, with anisocoria for more than two hours from the time of the first neurological examination, or bilateral mydriasis at presentation, as they independently contribute to massive cerebral infarction and diffuse cerebral edema.

Predictive imaging factors for DC indication include hematoma volume above 100 ml or the presence of the swirl sign. Evidence of infarction in the middle cerebral artery or posterior cerebral artery territory also serves as an indicator for DC, as

does midline structure deviation, indicating elevated intracranial pressure. The incidence of post-traumatic cerebral infarction secondary to IEH is 18.2%, even higher in patients with risk factors such as transtemporal location, preoperative hypovolemic shock for more than 30 minutes, bilateral mydriasis, and preoperative cerebral herniation.^{13,46,47}

DC plays a crucial role in controlling elevated intracranial pressure and is recommended when feasible in severe cases of post-traumatic cerebral infarction secondary to IEH.⁴⁶⁻⁵⁰ According to Vilcinis et al,¹⁶ DC may effectively reduce intracranial pressure, but its effects on outcomes remain unknown. However, DC associated with IEH drainage in patients with profound coma appears to have inferior outcomes compared to osteoplastic craniotomy. Therefore, DC associated with initial IEH drainage may represent the best surgical option in well-selected cases, although more prospective studies are needed for conclusive evidence.

Inappropriately indicated DC as initial hematoma drainage may lead to inevitable complications, such as cerebral hemodynamic abnormalities, subsequent cerebral necrosis, and infarction, as well as the need for cranioplasty. The decision to drain the hematoma with or without DC, especially in patients with cerebral herniation, remains controversial.^{46,51}

Arterial Embolization

Recent technological advancements have enabled endovascular treatment in a variety of cerebrovascular lesions, including IEH.^{18,52-55} Middle meningeal artery embolization plays a crucial role as an alternative treatment, particularly in certain coexisting disease scenarios where conventional surgical treatment of IEH is precluded.^{55,56} For other authors, endovascular treatment has proven useful in cases of small IEH, serving as an effective treatment option.^{17,18,54,57}

Endovascular treatment of the middle meningeal artery was initially described in 2004 by Suzuki et al,¹⁸ and subsequent studies have reinforced its indication.^{17,18,53,58} Endovascular intervention is a method that can be effective in treating IEH without hematoma progression compared to traditional craniotomy.^{59,60} According to Madison et al,⁵⁹ treatment decisions depend on the severity of the patient's injury and their neurological conditions. Ye et al⁶¹ reported that endovascular intervention yields good results in treating IEH complicated by oronasal hemorrhage due to skull base fracture. Other authors recommend endovascular intervention in the treatment of IEH or its progression.^{18,62} Peres et al¹⁷ treated 80 patients with IEH, primarily in temporal locations, with excellent results. Suzuki et al¹⁸ used endovascular embolization in nine patients during the acute phase of IEH and achieved excellent outcomes. The potential role of endovascular embolization in the treatment of IEH in carefully selected patients has been beneficial.

Ultrasound and/or CT-Guided Aspiration

In cases of neonates, needle aspiration guided by transcranial ultrasound can be performed with excellent results.^{63,64} Spontaneous resolution without the need for neurosurgical intervention may also occur, but in cases of voluminous IEH

with mass effect, surgical treatment is indicated to prevent neurological deterioration and IEH ossification.⁶⁵

Zhao et al¹⁹ used traumatic IEH drainage in 33 selected patients, with puncture guided by CT and instillation of urokinase during drainage, concluding that this procedure is safe, simple, rapid, and accurate, as reported by other authors as well.^{1,14,66}

In cases associated with cephalohematoma, aspiration can be performed due to the common finding of communication between these hematomas.⁶⁷⁻⁷¹ It has been described that between 61% to 70% of IEH in young children coexist with cephalohematoma, and half of these have communication between them.⁶⁸ Smets and Vanhanwaert⁷² suggest that in the absence of neurological signs and symptoms, aspiration of the cephalohematoma to drain the communicating IEH in newborns can avoid a more aggressive surgical intervention.

Endoscopic Drainage and Minimally Invasive Surgery

Endoscopic treatment for IEH drainage has been rare. This procedure has been performed in cases of acute spinal epidural hematoma and subdural and intraparenchymal hematomas.^{22,46,73} Oshima et al⁵⁵ performed a combined approach (endoscopy and embolization of the middle meningeal artery), under local anesthesia for IEH drainage, in an elderly patient with poor clinical conditions, with excellent results.

Minimally Invasive Surgery (MIS) has been increasingly performed in patients with intracranial hematoma. It has the advantage of better-preserving brain tissue compared to more invasive procedures. According to Huang et al,⁷⁴ MIS is suitable for uncomplicated IEH, while for IEH involving the temporal base or major venous sinuses, craniotomy is the preferred initial choice. Wang et al⁴⁶ reported good results in 59 cases of IEH treated with aspiration and drainage through MIS, using urokinase in eight cases with excellent results.

Tseng et al²¹ concluded in their study that MIS assisted with an endoscope in traumatic hemorrhage are efficient and effective in carefully selected cases. According to Tseng et al,²¹ its advantages include early decompression, reduced surgical time, less blood loss, small incisions, early recovery, short hospital stays, and lower operational costs. Lu et al⁷⁵ performed aspiration and drainage through MIS, being effective in treating 58 patients with IEH involving the transverse sinus. Other authors performed trepanation followed by urokinase instillation and concluded it to be an efficient, easily performed technique effective in IEH treatment in selected cases.^{14,76,77} With the evolution of new devices and techniques, it will be a procedure to be successfully performed in experienced hands in the future.⁵⁵

Prognosis

IEH is an important cause of morbidity and mortality in patients with TBI.⁴ Factors that significantly influence prognosis include age, low GCS score on admission, presence of associated intradural injury, time elapsed between trauma and symptom onset, hematoma size, and location.^{3,49,78,79} Mortality from IEH varies from 0% to 21%.^{12,80-84} Stephanov⁸⁵

demonstrated that pre-CT era mortality ranged from 16% to 52%, while post-CT era mortality ranged from 8% to 14%. This author concluded that rapid patient transport to a neurosurgical referral center was the most important factor in reducing mortality. Jones et al⁸⁶ reported a decrease in mortality from 29% to 8% over the last 35 years. Other authors have shown that the absence of an early diagnosis, anisocoria, preoperative GCS score, time between injury and surgery, and presence of associated intradural injuries contribute to high morbidity and mortality.^{87–90}

Conclusion

The treatment of IEH is considered one of the most rewarding neurosurgical interventions for the neurosurgeon because when treated early and appropriately, the patient transitions from a severe neurological state to a normal one. Therefore, it is of utmost importance that both neurosurgeons and clinicians be able to identify the characteristic signs and symptoms to avoid treatment delay. Furthermore, it is concluded that treatment is invariably surgical in many cases. Additionally, it is worth noting that minimally invasive approaches have gained prominence in recent decades, associated with image-guided procedures, and when well-indicated, result in rapid recovery and lower morbidity.

Conflict of Interest

The authors report no conflicts of interest.

References

- Zhu Q, von Spreckelsen N, Huang P, et al. Minimally invasive puncture with twist intraosseous drill needle combined with hematoma drainage in the treatment of acute epidural hematoma in pediatric patients: A technical note. *Clin Neurol Neurosurg* 2023;226:107626
- Bisen YT, Korde P, Dighe O, Iratwar S, Bisen G. Decompressive craniectomy in the management of low Glasgow Coma Score patients with extradural hematoma: a review of literature and guidelines. *Cureus* 2023;15(01):e33790. Doi: 10.7759/cureus.33790
- Bricolo AP, Pasut LM. Extradural haematoma: toward zero mortality. *Neurosurgery* 1984;14(01):8–12
- Kalkan E, Cander B, Gul M, Girisgin S, Karabaghi H, Sahin B. Prediction of prognosis in patients with epidural hematoma by a new stereological method. *Tohoku J Exp Med* 2007;211(03):235–242
- Kvarnes TL, Trumpy JH. Extradural haematoma. Report of 132 cases. *Acta Neurochir (Wien)* 1978;41(1-3):223–231
- Paşaoğlu A, Orhon C, Koç K, Selçuklu A, Akdemir H, Uzunoğlu H. Traumatic extradural haematomas in pediatric age group. *Acta Neurochir (Wien)* 1990;106(3-4):136–139
- Baykaner K, Alp H, Çeviker N, Keskil S, Seçkin Z. Observation of 95 patients with extradural hematoma and review of the literature. *Surg Neurol* 1988;30(05):339–341
- Lobato RD, Rivas JJ, Cordobes F, et al. Acute epidural hematoma: an analysis of factors influencing the outcome of patients undergoing surgery in coma. *J Neurosurg* 1988;68(01):48–57
- Rivas JJ, Lobato RD, Sarabia R, Cordobés F, Cabrera A, Gómez P. Extradural hematoma: analysis of factors influencing the courses of 161 patients. *Neurosurgery* 1988;23(01):44–51
- Rocchi G, Caroli E, Raco A, Salvati M, Delfini R. Traumatic epidural hematoma in children. *J Child Neurol* 2005;20(07):569–572
- Gupta SK, Tandon SC, Mohanty S, Asthana S, Sharma S. Bilateral traumatic extradural haematomas: report of 12 cases with a review of the literature. *Clin Neurol Neurosurg* 1992;94(02):127–131
- Chen TY, Wong CW, Chang CN, et al. The expectant treatment of “asymptomatic” supratentorial epidural hematomas. *Neurosurgery* 1993;32(02):176–179, discussion 179
- Bullock MR, Chesnut R, Ghajar J, et al; Surgical Management of Traumatic Brain Injury Author Group. Surgical management of acute epidural hematomas. *Neurosurgery* 2006;58(03):S7–S15, discussion Si-iv
- Liu JT, Tyan YS, Lee YK, Wang JT. Emergency management of epidural haematoma through burr hole evacuation and drainage. A preliminary report. *Acta Neurochir (Wien)* 2006;148(03):313–317, discussion 317
- Korde PA, Iratwar SW, Patil A, Mundhe VM, Rathod CT. Decompressive craniectomy for traumatic acute extradural haematoma: Decision making and outcomes. *J Clin Diagn Res* 2020;14(01):PRO01–PRO03
- Vilcinis R, Bunevicius A, Piliponis L, Tamasauskas A. Influence of decompressive craniectomy post evacuation of epidural hematoma in comatose patients. *World Neurosurg* 2021;151:e753–e759. Doi: 10.1016/j.wneu.2021.04.109
- Peres CMA, Caldas JGMP, Puglia P, et al. Endovascular management of acute epidural hematomas: clinical experience with 80 cases. *J Neurosurg* 2018;128(04):1044–1050
- Suzuki S, Endo M, Kurata A, et al. Efficacy of endovascular surgery for the treatment of acute epidural hematomas. *AJNR Am J Neuroradiol* 2004;25(07):1177–1180
- Zhao X, Jiang H, Liu G, Wang T. Efficacy analysis of 33 cases with epidural hematoma treated by brain puncture under CT surveillance. *Turk Neurosurg* 2014;24(03):323–326
- Wang WH, Hu LS, Lin H, et al. Risk factors for post-traumatic massive cerebral infarction secondary to space-occupying epidural hematoma. *J Neurotrauma* 2014;31(16):1444–1450
- Tseng WL, Kuo LT, Chen CM, et al. Surgical application of endoscopic-assisted minimally-invasive neurosurgery to traumatic brain injury: Case series and review of literature. *J Formos Med Assoc* 2022;121(07):1223–1230
- Codd PJ, Venteicher AS, Agarwalla PK, Kahle KT, Jho DH. Endoscopic burr hole evacuation of an acute subdural hematoma. *J Clin Neurosci* 2013;20(12):1751–1753
- Pang D, Horton JA, Herron JM, Wilberger JE Jr, Vries JK. Nonsurgical management of extradural hematomas in children. *J Neurosurg* 1983;59(06):958–971
- Korinth M, Weinzierl M, Gilsbach JM. [Treatment options in traumatic epidural hematomas]. *Unfallchirurg* 2002;105(03):224–230
- Shahid A, Mumtaz A, Mohd I. Acute extradural haematoma: Factors affecting the outcome. *J Postgrad Med Inst* 2005;19:208–211
- Wong CW. The CT criteria for conservative treatment—but under close clinical observation—of posterior fossa epidural haematomas. *Acta Neurochir (Wien)* 1994;126(2-4):124–127
- Zwayed ARH, Lucke-Wold B. Conservative management of extradural hematoma: A report of sixty-two cases. *Neurol Clin Neurosci* 2018;2(02):5–9
- Cucciniello B, Martellotta N, Nigro D, Citro E. Conservative management of extradural haematomas. *Acta Neurochir (Wien)* 1993;120(1-2):47–52
- Samadi-Motlagh P, Shimia M, Shaken M, et al. Role of tranexamic acid in conservative treatment of patients with epidural hematoma. *Iran J Neurosurg* 2016;2(03):8–10
- Sullivan TP, Jarvik JG, Cohen WA. Follow-up of conservatively managed epidural hematomas: implications for timing of repeat CT. *AJNR Am J Neuroradiol* 1999;20(01):107–113
- Ndoumbe A, Patience EMV, Simen C, Takongmo S. Outcome of surgically treated acute traumatic epidural hematomas based on the Glasgow coma scale. *Open J Mod Neurosurg* 2018;8:109–118

- 32 Pozzati E, Tognetti F. Spontaneous healing of acute extradural hematomas: study of twenty-two cases. *Neurosurgery* 1986;18(06):696–700
- 33 Muñoz-Suárez D, Moreno-García S. Resolución espontánea de un hematoma epidural traumático en 17 horas de evolución en un niño de 9 años. Reporte de caso. *Rev Chil Neurocir* 2020;46:83–87
- 34 Manne S, Musali SR, Gollapudi PR, Karla R. Spontaneous resolution of epidural hematoma. A rare case. *Asian J Neurosurg* 2019;14(01):292–294
- 35 Yokoyama T, Sugimoto T, Yoneyama T, Futami M, Takeshima H. An unusual case of acute epidural hematoma showing rapid spontaneous resolution with delayed recurrence. *No Shinkei Geka. No Shinkei Geka* 2018;46(05):405–411
- 36 Costa Clara JM, Claramunt E, Ley L, Lafuente J. Traumatic extradural hematomas of the posterior fossa in children. *Childs Nerv Syst* 1996;12(03):145–148
- 37 Grevsten S, Pellettieri L. Surgical decision in the treatment of extradural haematoma. *Acta Chir Scand* 1982;148(02):97–102
- 38 Springer MF, Baker FJ. Cranial burr hole decompression in the emergency department. *Am J Emerg Med* 1988;6(06):640–646
- 39 Usman B, Mohamed B, Daibu U. Outcomes of evacuating subacute extradural hematoma through a micraniectomy. A 5-year study. *Indian J Neurotrauma* 2023;•••. Doi: 10.1055/s – 0043–1760742
- 40 Nelson JA. Local skull trephination before transfer is associated with favorable outcomes in cerebral herniation from epidural hematoma. *Acad Emerg Med* 2011;18(01):78–85
- 41 Soon WC, Marcus H, Wilson M. Traumatic acute extradural haematoma - Indications for surgery revisited. *Br J Neurosurg* 2016;30(02):233–234
- 42 Zhang Y, Li Q, Zhao R, et al. Novel minimally invasive treatment strategy for acute traumatic epidural hematoma: Endovascular embolization combined with drainage surgery and use of urokinase. *World Neurosurg* 2018;110:206–209
- 43 Gurer B, Kertmen H, Yilmaz ER, Dolgun H, Hasturk AE, Sekerci Z. The surgical outcome of traumatic extradural hematomas causing brain herniation. *Turk Neurosurg* 2017;27(01):37–52
- 44 Yang C, Huang X, Feng J, et al. Prospective randomized evaluation of decompressive ipsilateral craniectomy for traumatic acute epidural hematoma (PREDICT_AEDH) study protocol for a randomized controlled trial. *Trials* 2021;22(01):421. Doi: 10.1186/s13063-021-05359-6
- 45 Okamoto T, Umezawa K, Ogita S, Kurosaki K, Takegami T, Kimura S. Two cases of difficult-to-treat acute epidural hematoma and a review of decompressive craniectomy with hematoma evacuation. *No Shinkei Geka* 2018;46(03):227–234
- 46 Wang W. Minimally invasive surgical treated of acute epidural hematoma. *Case series. BioMed Res Int* 2016;2016:6507350. Doi: 10.1155/2016/6507350
- 47 Otani N, Takasato Y, Masaoka H, et al. Surgical outcome following a decompressive craniectomy for acute epidural hematoma patients presenting with associated massive brain swelling. *Acta Neurochir Suppl (Wien)* 2010;106:261–264. Doi: 10.1007/978-3-211-98811-4-49
- 48 Hutchinson PJ, Koliatsos AG, Tajsic T, et al. Consensus statement from the international consensus meeting on the role of decompressive craniectomy in the management of traumatic brain injury. *Acta Neurochir (Wien)* 2019;161(07):1261–1274
- 49 Servadei F. Prognostic factors in severely head injured adult patients with epidural haematoma's. *Acta Neurochir (Wien)* 1997;139(04):273–278
- 50 Cooper DJ, Rosenfeld JV, Murray L, et al; DECRA Trial Investigators Australian and New Zealand Intensive Care Society Clinical Trials Group. Decompressive craniectomy in diffuse traumatic brain injury. *N Engl J Med* 2011;364(16):1493–1502
- 51 Lin H, Wang WH, Hu LS, et al. Novel clinical scale for evaluating pre-operative risk of cerebral herniation from traumatic epidural hematoma. *J Neurotrauma* 2016;33(11):1023–1033
- 52 Lammy S, McConnell R, Kamel M, Rennie I, Al-Haddad S. Extradural haemorrhage: is there a role for endovascular treatment? *Br J Neurosurg* 2013;27(03):383–385
- 53 Misaki K, Muramatsu N, Nitta H. Endovascular treatment for traumatic ear bleeding associated with acute epidural hematoma. *Neurol Med Chir (Tokyo)* 2008;48(05):208–210
- 54 Ross IB. Embolization of the middle meningeal artery for the treatment of epidural hematoma. *J Neurosurg* 2009;110(06):1247–1249
- 55 Ohshima T, Tajima H, Fujii K, et al. Combined endovascular and endoscopic surgery for acute epidural hematoma in a patient with poor health. *Neurol Med Chir (Tokyo)* 2012;52(11):829–831
- 56 Park TJ, Lee SP, Baek J, Ryou K, Kim SH. Middle meningeal artery embolization to treat progressive epidural hematoma: a case report. *J Cerebrovasc Endovasc Neurosurg* 2020;22(01):20–25. Doi: 10.7461/jcen.2020.22.2.20
- 57 Offner PJ, Pham B, Hawkes A. Nonoperative management of acute epidural hematomas: a “no-brainer”. *Am J Surg* 2006;192(06):801–805
- 58 Bortoluzzi M, Pavia M. Endovascular treatment of incoercible epistaxis and epidural cerebral hematoma. A case report. *Interv Neuroradiol* 2006;12(03):233–236
- 59 Madson MT, Graupman PC, Carroll JM, Torok CM, Touchette JC, Nussbaum ES. Traumatic epidural hematoma treated with endovascular coil embolization. *Surg Neurol Int* 2021;12(322):322
- 60 Fan G, Wang H, Ding J, et al. Application of absolute alcohol in the treatment of traumatic intracranial hemorrhage via interventional embolization of middle meningeal artery. *Front Neurol* 2020;11:824
- 61 Ye Z, Jin H, Chen Y, Ji H, Xu H, Jin Y. Endovascular treatment of traumatic oronasal hemorrhage complicated with progressive acute epidural hematoma. *Medicine (Baltimore)* 2022;101(03):e28654
- 62 Zussman BM, Goldschmidt E, Faraji AH, Salvetti DJ, Jankowitz BT. Middle meningeal artery embolization for the treatment of an expanding epidural hematoma. *World Neurosurg* 2019;128:284–286
- 63 Noguchi M, Inamasu J, Kawai F, et al. Ultrasound-guided needle aspiration of epidural hematoma in a neonate after vacuum-assisted delivery. *Childs Nerv Syst* 2010;26(05):713–716
- 64 Vachharajani A, Mathur A. Ultrasound-guided needle aspiration of cranial epidural hematoma in a neonate: treating a rare complication of vacuum extraction. *Am J Perinatol* 2002;19(08):401–404
- 65 Yu DK, Heo DH, Cho SM, Cho YJ. Rapidly calcified epidural hematoma in a neonate. *J Korean Neurosurg Soc* 2008;44(02):98–100
- 66 Li JG, Wang XD, Luo YM. Clinical analysis of microinvasive puncture and drainage for acute traumatic epidural hematoma with YL-1 puncture needle. *Chin J Mod Operative Surg* 2010;14:469–470
- 67 Oh KW, Kim HM. Epidural hematoma treated by aspiration of accompanying cephalohematoma in a newborn infant. *Korean J Pediatr* 2007;50(11):1125–1128
- 68 Yamamoto T, Enomoto T, Nose T. Epidural hematoma associated with cephalohematoma in a neonate—case report. *Neurol Med Chir (Tokyo)* 1995;35(10):749–752
- 69 Heyman R, Heckly A, Magagi J, Pladys P, Hamlat A. Intracranial epidural hematoma in newborn infants: clinical study of 15 cases. *Neurosurgery* 2005;57(05):924–929, discussion 924–929
- 70 Negishi H, Lee Y, Itoh K, et al. Nonsurgical management of epidural hematoma in neonates. *Pediatr Neurol* 1989;5(04):253–256
- 71 Aoki N. Epidural hematoma communicating with cephalohematoma in a neonate. *Neurosurgery* 1983;13(01):55–57
- 72 Smets KJ, Vanhauwaert D. Treatment of cranial epidural hematoma in a neonate by needle aspiration of a communicating cephalohematoma. *Eur J Pediatr* 2010;169(05):617–619
- 73 Miki K, Nonaka M, Kobayashi H, et al. Optimal surgical indications of endoscopic surgery for traumatic acute subdural hematoma in

- elderly patients based on a single-institution experience. *Neurosurg Rev* 2021;44(03):1635–1643
- 74 Huang APH, Huang SJ, Hong WC, et al. Minimally invasive surgery for acute noncomplicated epidural hematoma: an innovative endoscopic-assisted method. *J Trauma Acute Care Surg* 2012;73(03):774–777
- 75 Lu Z, Zhu G, Qiu Y, Cheng X. Minimally-invasive aspiration and drainage for management of traumatic epidural hematoma straddling transverse sinus. *Neurol India* 2013;61(02):111–116
- 76 Shrestha D, Feng L. A novel surgical approach to traumatic intracranial epidural hematoma. *Open Access Library Journal* 2017;4:e3820. Doi: 10.4236/oalib.1103820
- 77 Park J, Kim GJ, Hwang SK. Thrombolytic evacuation of post-craniotomy epidural haematomas using closed suction drains: a pilot study. *Acta Neurochir (Wien)* 2008;150(04):359–366, discussion 366
- 78 Dubey A, Pillai SV, Kolluri SV. Does volume of extradural hematoma influence management strategy and outcome? *Neurol India* 2004;52(04):443–445
- 79 Andrioli GC, Zuccarello M, Trinica G, Fiore D. Extradural hematomas in elderly. A statistical analysis of 58 cases. *Adv Neurosurg* 1984;12:218–223
- 80 Bezircioğlu H, Erşahin Y, Demirçivi F, Yurt I, Dönertaş K, Tektaş S. Nonoperative treatment of acute extradural hematomas: analysis of 80 cases. *J Trauma* 1996;41(04):696–698
- 81 Alliez JR, Hilal N, Kaya JM, Leone M, Reynier Y, Alliez B. Hématomes intracrâniens extra-duraux: À propôs de 100 cases récents. *Neurochirurgie* 2005;51:464–470
- 82 Beni-Adani L, Flores I, Spektor S, Umansky F, Constantini S. Epidural hematoma in infants: a different entity? *J Trauma* 1999;46(02):306–311
- 83 Dhellemmes P, Lejeune JP, Christiaens JL, Combelles G. Traumatic extradural hematomas in infancy and childhood. Experience with 144 cases. *J Neurosurg* 1985;62(06):861–864
- 84 Jamjoom A. The influence of concomitant intradural pathology on the presentation and outcome of patients with acute traumatic extradural haematoma. *Acta Neurochir (Wien)* 1992;115(3-4):86–89
- 85 Stephanov S. Post-operative mortality in acute extradural haematoma. *Br J Neurosurg* 1993;7(05):461–463
- 86 Jones NR, Molloy CJ, Kloeden CN, North JB, Simpson DA. Extradural haematoma: trends in outcome over 35 years. *Br J Neurosurg* 1993;7(05):465–471
- 87 Bejjani GK, Donahue DJ, Rusin J, Broemeling LD. Radiological and clinical criteria for the management of epidural hematomas in children. *Pediatr Neurosurg* 1996;25(06):302–308
- 88 Cheung PSY, Lam JMY, Yeung JHH, Graham CA, Rainer TH. Outcome of traumatic extradural haematoma in Hong Kong. *Injury* 2007;38(01):76–80
- 89 Chowdhury NKSM, Raihan MZ, Chowdhury FH, Ashadullah Atm, Sarkar MH, Hossain SS. Surgical management of traumatic extradural hematoma: experience of 610 patients and prospective analysis. *Indian J Neurotrauma* 2008;5:75–79
- 90 Ayub S, Ali M, Ilyas M. Acute extradural hematoma: factors affecting the outcome. *J Postgrad Med Inst* 2005;19:208–211