MUSCULOSKELETAL RADIOLOGY

Case report: MRI of decubital ischemic fasciitis

Ozden Sila Ulus, Ercan Karaarslan, Yesim Saglican¹, Abdullah Yakupoglu²

Department of Radiology, Acibadem University, Departments of ¹Pathology and ²Radiology, Acibadem Maslak Hospital, Istanbul, Turkey

Correspondence: Dr. Ozden Sila Ulus, Gülsuyu Mah, Fevzi Çakmak Cad, Divan Sok. No: 1 Maltepe, Istanbul, Turkey. E-mail: silaulus@hotmail.com

Abstract

The MRI findings in a case of decubital ischemic fasciitis located posterolateral to the right greater trochanter, in a 72-year-old woman, are presented. Decubital ischemic fasciitis is an uncommon entity encountered mostly in debilitated, elderly patients, in the deep subcutaneous tissue, at pressure points or bony prominences. It can simulate soft-tissue sarcomas. Recognition of this lesion radiologically is important to prevent unnecessary interventions.

Key words: Decubital ischemic fasciitis; hip; MRI; soft-tissue tumor

Introduction

Decubital ischemic fasciitis is a rare entity occurring mostly in the elderly and in debilitated patients, in the deep subcutaneous tissue, at pressure points or bony prominences.^[1-4] It can simulate a soft-tissue sarcoma clinically and histopathologically.^[1-4] Imaging helps to rule out malignancies and therefore to prevent unnecessary interventions. We would like to highlight the MRI findings of this entity.

Case Report

A 72-year-old woman presented with a palpable mass over the right hip region. She was being treated for Parkinson's disease. Physical examination revealed a 4- to 5-cm, hard, and fixated mass in the right proximal thigh, located posterolateral to the greater trochanter. Radiography of the pelvis was normal. MRI revealed diffuse hyperintense signals on the T2W images [Figures 1A and 1B] consistent

Access this article online	
Quick Response Code:	
	Website: www.ijri.org
	DOI: 10.4103/0971-3026.82289

with edema and inflammation, in the muscles and subcutaneous tissues, with prominent, peripheral, and illdefined enhancement [Figure 2], over an area measuring approximately 4–7 cm², at the level of the greater trochanter. A CT-guided needle biopsy showed coagulative necrosis, hemorrhage, and fibrosis involving the adipose tissue. The capillary and muscular vessels within the lesion also showed fibrinoid necrosis in their walls, without evidence of primary vasculitis [Figure 3]. Scattered reactive fibroblasts with enlarged nuclei were present in or around the necrosis. No neoplastic process was detected. These findings, along with the clinical features, were considered compatible with the diagnosis of ischemic fasciitis (atypical decubital fibroplasia).

Discussion

Decubital ischemic fasciitis, also called atypical decubital fibroplasia, is a distinctive fibroplasia occurring predominantly in the elderly and debilitated patients, confined to bed or wheelchair-bound.^[1-3] The lesions occur in the deep subcutaneous tissue at pressure points or bony prominences. It is a rare condition, and 40 cases have been reported in the literature^[1] until 2009. Ilaslan *et al.* have reported the MRI features of decubital ischemic fasciitis in three patients.^[4]

On MRI, the mass-like area in our patient was isointense compared to muscle on the T1W images, and hyperintense on the T2W images. After administration of gadolinium,



Figure 1 (A,B): Axial (A) and coronal (B) fat-suppressed T2W images show diffuse hyperintense signals (arrows) in the muscles and subcutaneous tissues, adjacent to the right greater trochanter



Figure 3: Histopathologic slide (H and E stain) shows coagulative necrosis and hemorrhage with capillary and muscular vessels showing fibrinoid necrosis in their walls (arrows)

subtle and peripheral enhancement was seen in the subcutaneous and muscular tissues. The bone adjacent to the affected soft tissue (greater trochanter) was not involved. Lesions occurring in the shoulder, sacral area, posterior chest wall, and vulvovagina have been reported.^[5]

The diagnosis of decubital ischemic fasciitis with typical findings on MRI prevents the already debilitated patient from undergoing further and unnecessary interventions. Consideration of this entity by the clinician and the radiologist helps the pathologist rule out malignancy. We want to point out that since MRI is a common and



Figure 2: Axial contrast-enhanced T1W, fat-suppressed MRI shows subtle, peripheral enhancement (arrow) in an area measuring approximately 4–7 cm², at the level of the greater trochanter

appropriate imaging method used to examine soft-tissue masses, all radiologists, especially those evaluating musculoskeletal MRI examinations, should be aware of this entity and be familiar with the clinical features, common locations, and MRI findings.

References

- 1. Znati K, Daoudi A, Elfatemi H, Chbani L, Bennis S, Boutayeb F, *et al.* Ischemic fasciitis. A case report. Ann Chir Plast Esthet 2009;54:392-5.
- Montgomery EA, Meis JM, Mitchell MS, Enzinger FM. A typical decubital fibroplasias. A distinctive fibroblastic pseudotumor occurring in debilitated patients. Am J Surg Pathol 1992;16:708-15.
- 3. Perosio PM, Weiss SW. Ischemic fasciitis: A juxta-skeletal fibroblastic proliferation with a predilection for elderly patients. Mod Pathol 1993;6:69-72.
- Ilaslan H, Joyce M, Bauer T, Sundaram M. Decubital ischemic fasciitis: Clinical, pathologic, and MRI features of pseudosarcoma. Am J Roentgenol 2006;187:1338-41.
- Scanlon R, Kelehan P, Flannelly G, McDonald D, McCluggage WG. Ischemic fasciitis: An unusual vulvovaginal spindle cell lesion. Int J Gynecol Pathol 2004;23:65-7.

Cite this article as: Ulus OS, Karaarslan E, Saglican Y, Yakupoglu A. Case report: MRI of decubital ischemic fasciitis. Indian J Radiol Imaging 2011;21:111-2.

Source of Support: Nil, Conflict of Interest: None declared.