

Etiologic and epidemiologic study of hepatocellular carcinoma in West Azarbaijan of Iran (2006–2011)

Nasim Valizadeh,
Rahim Mehdioghli¹,
Ramin Behroozian²

Departments of Hematology
and Medical Oncology, ¹Internal
Medicine, ²Gastroenterology,
Urmia University of Medical
Science, Urmia, Iran

Address for correspondence:

Dr. Nasim Valizadeh,
Department of Hematology and
Medical Oncology, Urmia University
of Medical Sciences, Urmia, Iran.
E-mail: nседaha0@gmail.com

ABSTRACT

Background: Hepatocellular carcinoma (HCC) is one of the most common cancers worldwide. Epidemiologic and etiologic studies help in the prevention of disease and improve surveillance of affected patients. The aim of this study was to study the etiology and epidemiology of HCC in West Azarbaijan of Iran. **Materials and Methods:** In this cross-sectional, retrospective study, medical records of 60 patients with HCC who were diagnosed between 2006 and 2011 in West Azarbaijan of Iran were analyzed for age, sex, location, presence of liver cirrhosis and/or well-known risk factors of HCC. Data were analyzed by SPSS 17 software. **Results:** From 60 patients with HCC, 75% were male and 25% were female. Mean age of the patients was 63.47 years. 51.66% were from rural areas and 48.33% were from urban areas. 43.33% of HCC patients were cirrhotic while 56.66% patients were non-cirrhotic. Of the 26 cirrhotic patients, 57.69% were seropositive for hepatitis B surface antigen (HBsAg), 15.38% not only had history of excess alcohol intake but were also HBsAg positive, 3.84% had history of alcoholism, 3.84% had history of Wilson disease and the rest (19.24%) had cryptogenic cirrhosis. Of the 34 non-cirrhotic patients, 35.29% were positive for HBsAg. None of the 60 patients with HCC had positive serology for hepatitis C virus (HCV) or history of hepatitis C infection. **Conclusion:** In our study, most of the patients (>50%) with HCC did not have underlying liver cirrhosis. In the cirrhotic patients, hepatitis B infection was the most common predisposing factor, followed by hepatitis B infection concomitant with alcoholism, alcoholism alone and Wilson disease. In the remaining cirrhotic patients, we could not find any predisposing factor for cirrhosis. Infection with HCV was not found in our studied group. We concluded that etiology of HCC in our province is different from that in the western countries.

Key words: Etiology, epidemiology, hepatocellular carcinoma, risk factor

INTRODUCTION

Hepatocellular carcinoma (HCC) is the fifth most common cancer and the second cause of cancer-related death of males.^[1,2]

In females, HCC is the seventh most common cancer and the sixth cause of cancer-related death.^[2] Eighty percent of patients with HCC have underlying chronic hepatitis B or C infection.^[3] HCC distribution is different among regions

of a country due to difference in exposure to hepatitis viruses and environmental pathogens.^[4] The HCC rate is higher in men than in women, with a 2:1 predominancy. Explanation for this male dominancy is that males have a higher possibility of being infected with hepatitis B virus (HBV) and hepatitis C virus (HCV), alcohol intake, smoking and hemochromatosis and have higher levels of androgenic hormones.^[5-7] Most cases of HCC are due to viral hepatitis (B and C) or cirrhosis.^[8] Etiology of HCC in patients with and without cirrhosis is different.^[9-12]

Of the important predisposing factors of HCC in western countries, which included HBV, HCV and alcohol, only HBV seems to directly lead to liver cell transformation.^[13]

Other risk factors of HCC included excessive alcohol consumption, aflatoxin intake, nonalcoholic steatohepatitis (NASH) and hereditary hemochromatosis.^[14]

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.107086

MATERIALS AND METHODS

In this cross-sectional, retrospective study, medical records of 60 patients with HCC who were diagnosed between 2006 and 2011 in the Imam Khomeini Hospital of Urmia City were analyzed for age, sex, living in a city or village, smoking, presence of liver cirrhosis and/or known risk factors of HCC. Data were analyzed by SPSS software. Diagnosis of HCC had been made in 50 patients with live biopsy and in the remaining 10 patients with solitary or multiple liver masses in abdominal imaging (size >2 cm) and markedly elevated alfa-fetoprotein (more than 500 ng/mL). Cirrhosis was defined by typical view of liver in sonography, prolonged prothrombin time (PT) and presence of ascites.

RESULTS

Of 60 patients with HCC, 75% ($n=45$) were male and 25% were female ($n=15$). Mean age of the patients was 63.47 years (range: 17–80 years). Mean age for females was 60.2 years (range: 20–80 years) and mean age for males was 66.75 years (range: 17–80 years). 51.66% ($n=31$) were from rural areas and 48.33% ($n=29$) were from urban areas. Sixty percent ($n=36$) of the patients had a history of smoking. 43.33% ($n=26$) of the patients were cirrhotic while 34 patients (56.66%) were non-cirrhotic [Table 1].

Of the 26 cirrhotic patients, 57.69% ($n=15$) had positive hepatitis B surface antigen (HBsAg), 15.38% ($n=4$) had HBs antigenemia combined with history of excess alcohol intake, 3.84% ($n=1$) had history of excess alcohol consumption, 3.84% ($n=1$) had history of Wilson disease and 19.24% ($n=5$) had cryptogenic cirrhosis. Of the 34 non-cirrhotic patients, 35.29% ($n=12$) were positive for HBsAg. None of the 60 patients with HCC had positive serology for HCV or history of hepatitis C infection [Table 2]. In our study, most of the patients (>50%) with HCC had no underlying liver cirrhosis. Of the cirrhotic patients ($n=26$), hepatitis B infection, alcoholism and Wilson disease were the causative agents and in 19.20% ($n=4$), we could not find any etiology of cirrhosis. Infection with HCV was not found in our studied group. We concluded that etiology of HCC in our province (West Azarbaijan of Iran) is different from that of other western countries.

DISCUSSION

Goetze *et al.* from eastern Germany studied 640 patients with HCC. In their study, 223 of the patients (83.5%) had liver cirrhosis. Alcohol-induced chronic liver disease followed by hepatitis C infection, NASH, hepatitis B infection, cryptogenic liver cirrhosis, alcohol in combination

Table 1: Clinical data of 60 patients with HCC

Male	45 (75%)
Female	15 (25%)
Mean age	63.47 (range:17–80)
Smoking	36 (60%)
Excess alcohol intake	5 (8.33%)
Cirrhotic	26 (43.33%)
Non-cirrhotic	34 (56.66%)
Rural area	31 (51.66%)
Urban area	29 (48.33%)

HCC - Hepatocellular carcinoma

Table 2: Risk factors for HCC in our studied group

Cirrhotic 43.33% ($n=26$)		Non-cirrhotic 56.66% ($n=34$)	
HBsAg positive	$n=15$	HBsAg positive	$n=12$
Excess alcohol intake	$n=1$	Other risk factors	$n=0$
HBsAg positive + excess alcohol intake	$n=4$		
Wilson disease	$n=1$		
HCV Ab positive	$n=0$		
Cryptogenic cirrhosis	$n=5$		

HCC - Hepatocellular carcinoma; HCV - Hepatitis C virus; $n=60$ patients

with viral hepatitis, hemochromatosis and hepatitis B and C coinfection were the causative agents of HCC.^[14]

Chiesa *et al.* from Italy studied 174 patients with newly diagnosed HCC. They found liver cirrhosis in 142 cases, excluded in 21 cases and remained undefined in 11 cases. They showed that most HCC cases are due to HBV, HCV or alcohol intake in both the presence and the absence of cirrhosis.^[15]

Iqbal *et al.* from north of Pakistan studied 100 patients with HCC. Seventy-four percent of them were HCV positive, 13% were HBsAg positive, 4% patients were HCV and HBsAg positive and in the remaining 9%, no predisposing cause was found. Eighty percent of them were cirrhotic.^[16]

Uzunlimoğlu *et al.* from Turkey studied 207 patients with HCC; they concluded that HBV infection is the highest leading cause of HCC in Turkey, followed by hepatitis C infection and alcoholic liver disease.^[17]

A metaanalysis was performed on 27,881 HCC patients for seroprevalence of HBsAg and HCV Ab from 90 studies. A predominance for HBsAg was found in HCC patients from Asian, African and Latin American countries, but anti-HCV dominance was found in Europe, North America, Japan, Pakistan, Mongolia and Egypt. Anti-HCV was found to be higher than HBsAg in Europe and in the United States.^[18]

In this study, we found that most of the patients with

HCC did not have hepatitis B and none of them had HCV infection; thus, it seems likely that high dietary exposure with aflatoxin and other environmental factors and toxins are possible predisposing factors for HCC. Aflatoxin was found in mouldy peanuts, wheat, soya beans, corn and rice; thus, we recommend searching for aflatoxin in these foods in West Azarbaijan of Iran. Giving information to people for proper care of crops and foods' storage can prevent from HCC in this province of Iran.

ACKNOWLEDGMENT

This work is a result from a thesis which was supported by Urmia University of Medical Sciences. Our special thank to staff of Imam Khomeini hospital that helped us in providing medical records of patients.

REFERENCES

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011;61:69.
- Epidemiology and etiologic association of associations of hepatocellular carcinoma. Up to date.011; 19.2 version.
- Perz JF, Armstrong GL, Farrington LA, Hutin YJ, Bell BP. Contribution of Hepatitis B virus and Hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J Hepatol* 2006;45:529.
- Munoz N, Bosch, X. Epidemiology of hepatocellular carcinoma. In: Neoplasms of the Liver, In: Okuda, K, Ishak, KG, Editors, Tokyo: Springer; 1989. p.3.
- McGlynn KA, Tsao L, Hsing AW, Devesa SS, Fraumeni JF Jr. International trends and patterns of primary liver cancer. *Int J Cancer* 2001;94:290-6.
- Yu MW, Chen CJ. Elevated serum testosterone levels and risk of hepatocellular carcinoma. *Cancer Res* 1993;53:790-4.
- Yu MW, Yang YC, Yang SY, Cheng SW, Liaw YF, Lin SM, *et al.* Hormonal markers and hepatitis B virus-related hepatocellular carcinoma risk: A nested case-control study among men. *J Natl Cancer Inst* 2001;93:1644-51
- Kumar V, Fausto N, Abbas A. Robbins and Cotran Pathologic Basis of Disease (7th ed). Philadelphia, Saunders 2003. p.914- 7
- Trevisani F, D'Intino PE, Caraceni P, Pizzo M, Stefanini GF, Mazziotti A, *et al.* Etiologic factors and clinical presentation of hepatocellular carcinoma. *Cancer (Phila.)* 1995;75:2220-32.
- Nzeako UC, Goodman ZD, Ishak KG. Hepatocellular carcinoma in cirrhotic and noncirrhotic livers. A clinical-histopathological study of 804 North American patients. *Am J Clin Pathol* 1996;105:65-75.
- Brechot C, Jaffredo F, Lagorce D, Gerken G, Meyer zum BÜschenfelde K, Papakonstantinou A, *et al.* Impact of HBV, HCV and GBV-C/HGV on hepatocellular carcinomas in Europe: Results of a European concerted action. *J Hepatol* 1998;29:173-83.
- Stroffolini T, Andreone P, Andriulli A, Ascione A, Craxi A, Chiaramonte M, *et al.* Characteristics of hepatocellular carcinoma in Italy. *J Hepatol* 1998;29:944-52.
- Idilman R, De Maria N, Colantoni A, Van Thiel DH. Pathogenesis of hepatitis B and C-induced hepatocellular carcinoma. *J Viral Hep* 1998;5:285-99.
- Goetze T, Schuette K, Bornschein J, Kipper M, Mohnike K, Malfertheiner P, *et al.* Epidemiology and clinical characterization of HCC in a tertiary center in eastern Germany. *J Clin Oncol* 29:2011 (suppl; abstr e12010)
- Chiesa R, Donato F, Tagger A, Favret M, Ribero ML, Nardi G, *et al.* Etiology of Hepatocellular Carcinoma in Italian Patients with and without Cirrhosis. *Cancer Epidemiol Biomarkers Prev* 2000;9:213-6.
- Iqbal S, Iman N, Khan H, Rahman S, Hameed A, Wadud Khan A, *et al.* Predisposing Factors for Hepatocellular Carcinoma in North Of Pakistan. *J Med Sci* 2010;18:63-6.
- Uzunlifoğlu O, Yurdaydin C, Çetinkaya H, Bozkaya H, Sahin T, Colakoğlu S, *et al.* Risk Factors for Hepatocellular Carcinoma in Turkey. *Dig Dis Sci* 2001;46:1022-8.
- Franceschi S, Raza SA. Epidemiology and prevention of hepatocellular carcinoma. *Cancer Lett* 2009;286:5-8

How to cite this article: Valizadeh N, Mehdioghli R, Behroozian R. Etiologic and epidemiologic study of hepatocellular carcinoma in West Azarbaijan of Iran (2006-2011). *Indian J Med Paediatr Oncol* 2012;33:221-3.

Source of Support: Nil, **Conflict of Interest:** None declared.

"Quick Response Code" link for full text articles

The journal issue has a unique new feature for reaching to the journal's website without typing a single letter. Each article on its first page has a "Quick Response Code". Using any mobile or other hand-held device with camera and GPRS/other internet source, one can reach to the full text of that particular article on the journal's website. Start a QR-code reading software (see list of free applications from <http://tinyurl.com/yzlh2tc>) and point the camera to the QR-code printed in the journal. It will automatically take you to the HTML full text of that article. One can also use a desktop or laptop with web camera for similar functionality. See <http://tinyurl.com/2bw7fn3> or <http://tinyurl.com/3ysr3me> for the free applications.