

Bacteriological profile and antibiotic susceptibility patterns of clinical isolates in a tertiary care cancer center

Vivek Bhat, Sudeep Gupta,
Rohini Kelkar¹,
Sanjay Biswas¹, Navin Khattry,
Aliasgar Moiyadi,
Prashant Bhat,
Reshma Ambulkar,
Preeti Chavan,
Shubadha Chiplunkar,
Amol Kotekar, Tejpal Gupta

ACTREC, Tata Memorial Centre,
¹TMH, Tata Memorial Centre,
Kharghar, Mumbai, Maharashtra,
India

Address for correspondence:

Dr. Sudeep Gupta,
ACTREC, Tata Memorial Centre,
Kharghar, Mumbai - 410 210,
Maharashtra, India.
E-mail: sudeepgupta04@yahoo.com

INTRODUCTION

In spite of the vast advances made by medical science in cancer treatment, infections remain a major cause of morbidity and mortality in patients diagnosed with cancer. The cancer patient is immunocompromised because of the nature of the disease itself and also due to interventions in the form of chemotherapy etc., in addition, there are usually other associated risk factors for acquiring infection such as long term catheterization, mucositis due to cytotoxic agents, neutropenia, and stem cell transplantation.^[1]

This increased risk of bacterial infections is further compounded by the rising trends of antibiotic resistance in commonly implicated organisms all over the world. This is particularly true in the case of members of *Enterobacteriaceae* group like *Escherichia coli* and *Klebsiella pneumoniae* and the nonfermenter group of organisms such as *Acinetobacter* spp. in the Indian setting. There is already widespread resistance to the cephalosporins as shown by ESBL (extended spectrum β -lactamase) and Amp C producers

ABSTRACT

Introduction: This increased risk of bacterial infections in the cancer patient is further compounded by the rising trends of antibiotic resistance in commonly implicated organisms. In the Indian setting this is particularly true in case of Gram negative bacilli such as *Escherichia coli*, *Klebsiella pneumoniae* and *Acinetobacter* spp. Increasing resistance among Gram positive organisms is also a matter of concern. The aim of this study was to document the common organisms isolated from bacterial infections in cancer patients and describe their antibiotic susceptibilities. **Methods:** We conducted a 6 month study of all isolates from blood, urine, skin/soft tissue and respiratory samples of patients received from medical and surgical oncology units in our hospital. All samples were processed as per standard microbiology laboratory operating procedures. Isolates were identified to species level and susceptibility tests were performed as per Clinical Laboratory Standards Institute (CLSI) guidelines -2012. **Results:** A total of 285 specimens from medical oncology (114) and surgical oncology services (171) were cultured. *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Acinetobacter* spp. were most commonly encountered. More than half of the *Acinetobacter* strains were resistant to carbapenems. Resistance in *Klebsiella pneumoniae* to cephalosporins, fluoroquinolones and carbapenems was >50%. Of the *Staphylococcus aureus* isolates 41.67% were methicillin resistant. **Conclusion:** There is, in general, a high level of antibiotic resistance among gram negative bacilli, particularly *E. coli*, *Klebsiella pneumoniae* and *Acinetobacter* spp. Resistance among Gram positives is not as acute, although the MRSA incidence is increasing.

Key words: *Antibiotics, resistance, susceptibility testing*

among the *Enterobacteriaceae*.^[2] Rampant use of antibiotics has unfortunately led to increasing resistance to the carbapenems as well, and this is generally due to carbapenemase production by the organisms.^[3] Prevalence of Metallo-beta-lactamase (MBL) producing organisms including New Delhi MBL-1 (NDM-1) is also on the rise in India.^[4]

Increasing resistance among Gram-positive organisms is also a matter of concern. High rates of Methicillin-Resistant *Staphylococcus aureus* (MRSA) in clinical samples have been noted in one study from North East India.^[5] Similarly, resistance to the glycopeptide antibiotics such as vancomycin and ticoplanin among clinical isolates of enterococci is also increasing.^[6] Empirical treatment of infection in the cancer patient is often arbitrarily attempted by administration of broad spectrum or combination antibiotics until culture, and

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Bhat V, Gupta S, Kelkar R, Biswas S, Khattry N, Moiyadi A, et al. Bacteriological profile and antibiotic susceptibility patterns of clinical isolates in a tertiary care cancer center. Indian J Med Paediatr Oncol 2016;37:20-4.

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.177010

susceptibility results are available. This could be made more evidence-based if the clinician has adequate information on the spectrum of microorganisms and the antimicrobial susceptibility patterns prevalent in that particular setting. It is possible that these patterns may differ from one geographical region to another and even from one hospital to another. The aim of this study is to document the common organisms isolated in cancer patients and describe their antibiotic susceptibilities.

MATERIALS AND METHODS

We conducted a 6 month study of all isolates from samples of patients received from medical and surgical oncology units from July 2013 to December 2013. Medical oncology included all hematolymphoid malignancies and hematopoietic stem cell transplant recipients. Surgical oncology included all other patients with solid tumors. All relevant samples were collected as per hospital sample collection protocol from various clinical areas; these included blood, pus/wound swabs, sputum, bronchoalveolar lavage, urine, etc., all samples were processed as per standard microbiology laboratory operating procedures.^[7] Isolates were generally identified up to species level by means of various biochemical tests. Susceptibility tests were performed as per Clinical Laboratory Standards Institute (CLSI, USA) guidelines 2012.^[8] Disc diffusion technique was the default method used for antibiotic susceptibility testing. In brief, lawn cultures of appropriate inoculum of respective organisms were performed in Mueller Hinton Agar (or Mueller-Hinton Blood agar for fastidious organisms) and antibiotic discs of required strengths were placed on the surface of the inoculated media and these were then incubated overnight. Zones of inhibition were measured the next day and were correlated with CLSI interpretive breakpoints to characterize them as Sensitive, Intermediate, and Resistant. For drugs such as colistin for which CLSI breakpoints are not available, the European Committee on Antimicrobial Susceptibility Testing interpretive breakpoints were used. In the case of other drugs like cefoperazone-sulbactam, cefepime-tazobactam, etc., interpretive breakpoints were provided by the manufacturer. *S. aureus* ATCC 25923, *E. coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used for Quality control.

For Gram-positive organisms, the antibiotics to be tested and reported were chosen from the following (depending on the organism isolated): Penicillin (10 units), erythromycin (15 µg), clindamycin (2 µg), amoxicillin-clavulanate (20/10 µg), ampicillin (10 µg), cefoxitin (30 µg), ciprofloxacin (5 µg), gentamicin (10 µg), vancomycin (30 µg/MIC), teicoplanin (30 µg), linezolid (30 µg), and co-trimoxazole (1.25/23.75 µg).

For Gram-negative, the antibiotics for respective organisms were chosen from the following: Amoxicillin-clavulanate (20/10 µg), ciprofloxacin (5 µg), levofloxacin (5 µg),

gentamicin (10 µg), amikacin (30 µg), netilmycin (30 µg), cefuroxime (30 µg), cefotaxime (30 µg), ceftazidime (30 µg), cefepime (30 µg), cefoperazone-sulbactam (75/25 µg), cefepime-tazobactam (30/10 µg), imipenem (10 µg), and meropenem (10 µg). Colistin susceptibility was performed by MIC (minimum inhibitory concentration) method, with *E*-test strips.

RESULTS

A total of 285 specimens from medical oncology (114) and surgical oncology services (171) were cultured. Table 1 shows the numbers of various specimens received from medical and surgical oncology services. Table 2 profiles the organisms isolated from all patients. ESBL producers among *E. coli* and *K. pneumoniae* were 38.46 and 35.17%, respectively.

Carbapenem resistance in *K. pneumoniae* was >50%. Of the *S. aureus* isolates 41.67% were MRSA (Methicillin Resistant

Table 1: Breakup of specimens received for bacterial culture and sensitivity

Specimen	Medical oncology	Surgical oncology
Blood	54	19
Respiratory	20	13
Skin and soft tissue	33	117
Urine	07	22
Total	114	171

Table 2: Organism profiles for medical and surgical oncology

Organism	Medical oncology	Surgical oncology	Total
Gram-positive			
<i>Coagulase negative staphylococcus</i>	20	17	37
<i>Enterococcus</i> spp.	7	5	12
<i>Other streptococci</i>	8	3	11
<i>Staphylococcus aureus</i>	5	19	24
Gram-negative			
<i>Escherichia coli</i>	18	43	61
<i>Klebsiella pneumoniae</i>	31	33	64
<i>Pseudomonas aeruginosa</i>	6	52	58
<i>Acinetobacter</i> spp.	9	3	12
<i>Stenotrophomonas maltophilia</i>	2	0	2
<i>Shewanella putrefaciens</i>	1	0	1
<i>Klebsiella ozonae</i>	2	0	2
<i>Klebsiella oxytoca</i>	4	1	5
<i>Proteus vulgaris</i>	4	4	8
<i>Proteus mirabilis</i>	3	7	10
<i>Citrobacter</i> spp.	1	1	2
<i>Providencia stuartii</i>	1	0	1
<i>Morganella morganii</i>	3	3	6
<i>Pseudomonas</i> spp.	2	1	3
<i>Chryseobacterium</i> spp.	0	11	11
<i>Serratia marcescens</i>	0	1	1

S. aureus). Figures 1-5 provide the detailed antibiotic susceptibility patterns for *S. aureus*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, and *Acinetobacter* spp. respectively.

DISCUSSION

Pneumonia and bacteremia are common infections seen in cancer patients followed by a urinary tract, skin and soft tissue and gastrointestinal infections.^[9] Our study population showed higher numbers of skin and soft tissue infections and lesser numbers of urinary tract infections [Table 1]. The greater numbers of stool cultures in our study reflect also the surveillance cultures that are performed for all our bone marrow transplant patients.

A wide variety of Gram-positive and Gram-negative organisms were isolated from clinical samples in our study [Table 2]. It is striking to note the high rates of resistance of enterobacteriaceae particularly *E. coli* and *K. pneumoniae* to the third generation cephalosporins (cefotaxime/cefazidime) and also to the β -lactam- β -lactamase inhibitor combinations such as cefoperazone-sulbactam and piperacillin-tazobactam. Similar high rates of

resistance of these organisms to the third generation cephalosporins have been noted in a multicentric study across Karnataka^[10] and another study from Bhopal.^[11] Approximately, half of the *E. coli* and *K. pneumoniae* isolates in the above studies were ESBL producers; lesser rates were seen in this study suggesting other methods of cephalosporin resistance also playing an important role in our setting. Fortunately, more than half of *E. coli* and *P. aeruginosa* isolates in our study retained clinically useful susceptibility to aminoglycosides (gentamicin, amikacin) and these still have an important role to play in the antibiotic treatment of these organisms in our set up. However >50% of *K. pneumoniae* and *Acinetobacter* isolates were resistant to gentamicin.

The rate of carbapenems resistant *Enterobacteriaceae* in the *Enterobacteriaceae* group of organisms particularly *E. coli* and *K. pneumoniae* is a worrying factor. It was much higher than a chinese study that showed only 6.6% carbapenem resistance with less than half of them producing the carbapenemases KPC-2, IMP-4, and NDM-1.^[12] One study has identified NDM-1 from different sites in India, mostly among *E. coli* and *K. pneumoniae* and these were highly resistant to all




Figure 1: *Staphylococcus aureus* — susceptibility patterns




Figure 2: *Escherichia coli* susceptibility patterns




Figure 3: *Klebsiella pneumoniae* susceptibility patterns




Figure 4: *Pseudomonas aeruginosa* — susceptibility patterns




Figure 5: *Acinetobacter* spp. susceptibility patterns

antibiotics except tigecycline and colistin.^[13] We did not perform molecular studies to identify the carbapenemases in our setting and were, therefore, unable to characterize them. Although the high rate of resistance of *P. aeruginosa* to piperacillin-tazobactam, amikacin and carbapenems have been described,^[14,15] most isolates in our setting were susceptible to primary antipseudomonals, aminoglycosides, and the carbapenems [Figure 4].

The resistance of *Acinetobacter* spp. to aminoglycosides and carbapenems was seen to be high in our study [Figure 5]. One study from Odisha, India showed very high rates of resistance of *Acinetobacter* to ceftazidime (93%), gentamicin (76%) and meropenem (22%).^[16] Colistin resistance was seen in rare isolates of *Acinetobacter* spp. and *K. pneumoniae*, and is a cause for concern because there are hardly any antibiotic alternatives left for these patients. We have tested select organisms for older drugs such as chloramphenicol, tetracycline, and chloramphenicol in such cases with limited success.

The problem of antibiotic resistance is fortunately not as high among the Gram-positive organisms. High rates of methicillin resistance (Manipal [54%], Puducherry [72.34%]),^[17,18] and the emergence of vancomycin intermediate strains of *S. aureus* have been reported from India. We did not encounter any vancomycin resistance among staphylococci, and MRSA rates have been approximate 41.67% [Figure 1].

There is a paucity of studies describing the microbiological profiles and antibiotic susceptibility patterns of organisms isolated from infections in the Indian oncology setting and more reports from similar centers would provide greater insights to this very important emerging issue in this patient population. Important factors leading to the development of resistance include misuse of antibiotics in clinics and hospitals; abundant use of antibiotics in animal farms, aquaculture and poultry; and overuse of anti-infectives and disinfectants.^[19] Attempts are being made

to formulate antibiotic policies at the national level. The “Chennai declaration” initiative provides directives and recommendations to tackle the menace of antimicrobial resistance at this level.^[20] However, it is also important for every healthcare setting to formulate antibiotic policies based on local antibiotic susceptibility patterns to so that arbitrary use of antibiotics is avoided and resistance is kept to a minimum.

CONCLUSION

There is, in general, a high level of antibiotic resistance among Gram-negative bacilli, particularly *E. coli*, *K. pneumoniae* and *Acinetobacter* spp. to the cephalosporins, β -lactam- β -lactam inhibitor combinations and carbapenems group of drugs. Occasional isolates resistant to colistin have also been noted. Resistance among Gram-positive organisms is not as acute, although the MRSA incidence is increasing.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Available from: <http://www.medscape.org/viewarticle/580631>. [Last accessed on 2015 Feb 15].
- Rudresh SM, Nagarathamma T. Extended spectrum β -lactamase producing *Enterobacteriaceae* & antibiotic co-resistance. Indian J Med Res 2011;133:116-8.
- Tzouveleki LS, Markogiannakis A, Psychogiou M, Tassios PT, Daikos GL. Carbapenemases in *Klebsiella pneumoniae* and other *Enterobacteriaceae*: An evolving crisis of global dimensions. Clin Microbiol Rev 2012;25:682-707.
- Kumar SG, Adithan C, Harish BN, Sujatha S, Roy G, Malini A. Antimicrobial resistance in India: A review. J Nat Sci Biol Med 2013;4:286-91.
- Tsering DC, Pal R, Kar S. Methicillin-resistant *Staphylococcus aureus*: Prevalence and current susceptibility pattern in Sikkim. J Glob Infect Dis 2011;3:9-13.
- Muto CA, Jernigan JA, Ostrowsky BE, Richet HM, Jarvis WR, Boyce JM, *et al.* SHEA guideline for preventing nosocomial transmission of multidrug-resistant strains of *Staphylococcus aureus* and *Enterococcus*. Infect Control Hosp Epidemiol 2003;24:362-86.
- Forbes BA, Sahm DF, Weissfeld AS. Bailey and Scott's Diagnostic Microbiology. 12th ed. Elsevier-Mosby, St. Louis, Missouri 63043, USA; 2007.
- Clinical Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twenty Second Informational Supplement M100-S22. Wayne, PA: Clinical Laboratory Standards Institute; 2012.
- Yadegarynia D, Tarrand J, Raad I, Rolston K. Current spectrum of bacterial infections in patients with cancer. Clin Infect Dis 2003;37:1144-5.
- Rao SP, Rama PS, Gurushanthappa V, Manipura R, Srinivasan K. Extended-spectrum beta-lactamases producing *Escherichia coli* and *Klebsiella pneumoniae*: A multi-centric study across Karnataka. J Lab Physicians 2014;6:7-13.
- Shashwati N, Kiran T, Dhanvijay AG. Study of extended spectrum β -lactamase producing *Enterobacteriaceae* and antibiotic co-resistance in a tertiary care teaching hospital. J Nat Sci Biol Med 2014;5:30-5.

12. Zhao ZC, Xu XH, Liu MB, Wu J, Lin J, Li B. Fecal carriage of carbapenem-resistant *Enterobacteriaceae* in a Chinese university hospital. *Am J Infect Control* 2014;42:e61-4.
13. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R, *et al.* Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: A molecular, biological, and epidemiological study. *Lancet Infect Dis* 2010;10:597-602.
14. Manoharan A, Chatterjee S, Mathai D; SARI Study Group. Detection and characterization of metallo beta lactamases producing *Pseudomonas aeruginosa*. *Indian J Med Microbiol* 2010;28:241-4.
15. Pawar M, Mehta Y, Purohit A, Trehan N, Rosenthal VD. Resistance in gram-negative bacilli in a cardiac intensive care unit in India: Risk factors and outcome. *Ann Card Anaesth* 2008;11:20-6.
16. Dash M, Padhi S, Pattnaik S, Mohanty I, Misra P. Frequency, risk factors, and antibiogram of *Acinetobacter* species isolated from various clinical samples in a tertiary care hospital in Odisha, India. *Avicenna J Med* 2013;3:97-102.
17. Menezes GA, Harish BN, Sujatha S, Vinothini K, Parija SC. Emergence of vancomycin-intermediate *Staphylococcus* species in Southern India. *J Med Microbiol* 2008;57(Pt 7):911-2.
18. Eshwara VK, Munim F, Tellapragada C, Kamath A, Varma M, Lewis LE, *et al.* *Staphylococcus aureus* bacteremia in an Indian tertiary care hospital: Observational study on clinical epidemiology, resistance characteristics, and carriage of the Pantone-Valentine leukocidin gene. *Int J Infect Dis* 2013;17:e1051-5.
19. Choudhury R, Panda S, Singh DV. Emergence and dissemination of antibiotic resistance: A global problem. *Indian J Med Microbiol* 2012;30:384-90.
20. Ghafur A, Mathai D, Muruganathan A, Jayalal JA, Kant R, Chaudhary D, *et al.* The Chennai declaration: A roadmap to tackle the challenge of antimicrobial resistance. *Indian J Cancer* 2013;50:71-3.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1024 kb. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than **4096 kb (4 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.