

# Exercise in cancer

P. Rajarajeswaran,  
R. Vishnupriya

College of Physiotherapy, Mother  
Theresa Post Graduate and  
Research Institute of Health  
Sciences, Puducherry – 6, India

## Address for correspondence:

Dr. P. Rajarajeswaran,  
College of Physiotherapy, Mother  
Theresa Post Graduate and  
Research Institute of Health  
Sciences, Puducherry – 6, India.  
E-mail: physio.raja@yahoo.com

## ABSTRACT

Physical exercise has attracted increased interest in rehabilitation of oncological patients. The purpose of this paper is to review the literature and summarize the evidence of physical exercise in preventing cancer, its ability in attenuating the effect of cancer and its treatments and to provide guidelines for exercise prescription. Review of recent literature by electronic search of MEDline (Pub Med), Cancer lit, Cochrane libraries, CINAHL were done using Keywords and the variables were identified and systematically evaluated. There is strong evidence for reduced risk of colorectal and breast cancer with possible association for prostate, endometrial and lung cancer with increasing physical activity. Exercise helps cancer survivors cope with and recover from treatment; exercise may improve the health of long term cancer survivors and extend survival. Physical exercise will benefit throughout the spectrum of cancer. However, an understanding of the amount, type and intensity of exercise needed has not been fully elucidated. There is sufficient evidence to promote exercise in cancer survivors following careful assessment and tailoring on exercise prescription.

**Key words:** *Exercise, cancer, prevention, rehabilitation*

**DOI:** 10.4103/0971-5851.60050

## INTRODUCTION

More than 10 million people are diagnosed with cancer worldwide; with improvement in early detection and treatment, increasing numbers of patients can be expected to be alive five years after they are diagnosed with cancer.<sup>[1]</sup> These individuals will join the expanding number of cancer survivors, estimated at about 25 million.<sup>[1]</sup> Current cancer treatment, although increasingly efficacious for improving survival are toxic in numerous ways and produce negative short and long term physiologic and or psychological effects, including pain, decreased cardio respiratory fitness, cancer related fatigue, reduced Quality Of Life (QOL) and suppressed immune function.<sup>[2]</sup> Interest in physical activity as a means for primary prevention of cancer is increasing as the evidence for its protective effect is rapidly accumulating. The International agency for research on cancer (IARC) estimates that 25% of cancer cases worldwide are caused by overweight or obesity and a sedentary lifestyle.<sup>[3]</sup> Physical activity is an attractive cancer preventive strategy because it potentially benefits many health's end points in addition to reducing the risk of certain cancers.<sup>[4]</sup> Physical activity may have benefits throughout the spectrum of living with cancer, but cancer survivors are often at increased risk for becoming too sedentary for several reasons.

Physical exercise has attracted increased interest in rehabilitation of oncological patients in general and also in palliative care.<sup>[5]</sup> In a growing body of research that has investigated exercise in cancer patients; dramatic improvements in physiologic and psychological functioning have been documented in patients participating in exercise programs. At least 15 meta-analysis have been published reviewing 100 studies showing the nearly universal to multifactorial benefits of exercise in this patient population. Evidence of the benefits of exercise for cancer survivors in areas of psychological and quality of life (QOL) outcomes,<sup>[6]</sup> cancers related fatigue,<sup>[7]</sup> physical functioning,<sup>[8]</sup> body weight and composition,<sup>[9]</sup> muscle strength and endurance,<sup>[10]</sup> immune function<sup>[11]</sup> and cardiovascular<sup>[12-13]</sup> fitness have been reported. It may reduce the risk of cancer recurrence, second primary cancers and other chronic diseases<sup>[4,8]</sup> as well as prolong survival.<sup>[5]</sup> Exercise may also alleviate symptoms that interfere with daily life of cancer patients and survivors such as lack of appetite, diarrhea, paresthesia, constipation, physical fatigue, mental fatigue, treatment related fatigue, muscle pain, arthralgia and other pain, depression, anxiety and insomnia.<sup>[12,13,14]</sup> The purpose of this paper is to review the literature and summarize the evidence of physical exercise in preventing cancer, its ability in attenuating the effect of cancer and its

treatments and to provide guidelines for exercise prescription.

## METHOD

Electronic search of MEDline (Pub Med), Cancer lit, Cochrane library, CINAHL were conducted using the following Keywords and its combinations- physical activity, exercise, prevention, intervention, cancer, neoplasm, quality of life, rehabilitation, chemotherapy, symptoms, side effects and biological mechanisms. Sources included references list of all relevant articles and reviews, Clinical Practice guidelines and Books. Review of recent systematic reviews, Meta analysis and studies on the topic that have been published in recent literature were reviewed and relevant articles are included. To be included in this review, a study had to be published in English Language between 2000 to present (Aug 09). Physical activities in cancer with multiple intervention and Pilot studies were excluded.

The following variables were identified and systematically evaluated in each paper: cancer type, age, gender, oncological treatment, QOL, biological mechanisms, type of exercise program and frequency, intensity, type and time (FIT) outcomes. Papers that met inclusion criteria and quality were studied. Definition of cancer survivor: As suggested by the national coalition of cancer survivorship to refer to any individual diagnosed with cancer from the time of discovery and for the balance of life.<sup>[15]</sup> Physical activity (PA) is defined as a bodily movement produced by skeletal muscle, which results in a substantial increase in energy expenditure over resting level. Physical exercise is defined as planned, structured, repetitive and purposeful physical activity.

## RESULTS

### Exercise in cancer prevention

Exercise may reduce the risk of developing a primary cancer, Nearly 150 studies have examined the relation between physical activity and cancer prevention at specific cancer sites, studies that meet the inclusion criteria and quality were studied and the results are summarized in Table 1 along with possible biological mechanism. The mechanism stated for association and cancer has not been established. These include changes in endogenous sexual and metabolic hormone levels,<sup>[35,36]</sup> growth factors,<sup>[60]</sup> decreased obesity and central adiposity<sup>[37]</sup> and possible changes in immune function.<sup>[65]</sup>

### Exercise benefits in cancer survivors

A majority of studies tested interventions and most of these studies used supervised exercise programs. Almost

all of the studies tested aerobic exercise programs although several combined aerobic and resistance exercise programs. Most of the early studies had significant methodological limitations. Despite these limitations however, the studies have consistently demonstrated that exercise has beneficial effects on a wide variety of physical fitness and QOL endpoints in cancer survivors including functional capacity, muscular strength, body weight and composition, flexibility, fatigue, nausea, diarrhea, pain, physical well-being, functional well being, depression, anxiety, rigor, anger, mood, self esteem, satisfaction with life and overall quality of life. These studies have resulted in exercise being recommended to cancer survivors by American Cancer Society and also as a therapy for fatigue in cancer survivors.

Table 2 summarizes the recent systematic reviews of physical activity in cancer survivors

The general consensus of these reviews is that physical activity has modest positive effect on supportive care outcomes including aerobic fitness, physical functioning, muscular strength, fatigue and some aspects of quality of life but the results are not as strong as the post-adjunct setting. In the review study by L.M. Oldervoll reported that some promising effect of physical exercises on overall Quality of life, fatigue, physical functioning, physical capacity and/or muscular fitness during and after cancer treatment.<sup>[75]</sup> C. Anderson *et al.* concluded that a six-week exercise intervention for cancer patient with or without disease and who are undergoing chemotherapy could lead to a reduction in symptoms and side effects of chemotherapy.<sup>[14]</sup> Physical activity may also help cancer survivors manage symptoms, improve mobility, slow functional decline and maintain quality of life at the end of life.<sup>[66]</sup> In a prospective phase II study, Olderswall *et al.* showed that structured physical exercise program by physiotherapist is a promising intervention for palliative cancer patient with short life expectancy and after six weeks there was a significant decrease in physical fatigue and improvements in physical and emotional functioning and concluded that physical exercise (Resistance exercise) is a feasible intervention in a palliative care setting and may be beneficial.<sup>[76]</sup> Retraining physical function and independence in activities of daily living are important factors in palliative patient.<sup>[5,77]</sup> Figure 1 shows how and when physical exercise may affect cancer experience.

These studies suggest that physical activity may help cancer survivors live longer by: reducing the risk of cancer recurrence or slowing cancer progression and reducing the risk of other life threatening diseases including second

**Table 1: Epidemiologic evidence on association between physical activity and cancer and possible biological mechanisms**

Cancer site	Average risk reduction %	Overall level of scientific evidence	Possible mechanisms involved	Rationale
Colon <sup>[16-19]</sup>	40-50	Convincing	Decreased gastrointestinal transit time Decreased ratio of prostaglandins Lowered bile acid secretion or enhanced acid metabolism	Physical activity increases gut motility and reduces mucosal exposure time to carcinogens. Strenuous exercise may increase prostaglandin (PG) F <sub>2</sub> , which inhibits colonic cell proliferation and increases gut motility while not increasing PGE <sub>2</sub> , which affects colonic cell proliferation, opposite to the effect of PGF <sub>2</sub> . Bile acid concentrations may be decreased in physically active (Confounding by diet) persons.
Breast <sup>[20-37]</sup>	30-40	Convincing	Decreased lifetime exposure to estrogen	Physical activity delays menarche, reduces the number of ovulatory cycles, and reduces ovarian estrogen production. It also reduces body fat and could reduce fat-produced estrogens. It increases the production of sex hormone-binding globulin, resulting in less biologically available estrogen.
Prostate <sup>[38-46]</sup>	10-30	Probable	Reduced exposure to testosterone	Physical activity increases production of sex hormone-binding globulin, resulting in lower levels of free testosterone
Endometrium <sup>[47,48]</sup>	30-40	Possible	Decreased percent body fat	Fat storage of carcinogens can occur in visceral fat, which can be released in overweight individuals.
Ovary <sup>[49-51]</sup>	20-30	Insufficient		
Lung <sup>[52,53]</sup>	30-40	Possible	NE	NE
Testis <sup>[54]</sup>	10-30	Insufficient	NE	NE
All cancers <sup>[55-65]</sup>	NE	NE	Genetic predisposition of habitually active people Exercise-induced increase in antitumor immune defenses Improved antioxidant defense systems Decreased circulating insulin and glucose Decreased insulin and insulin-like growth factors	Constitutional factors influence athletic selection or interest in physical activity and susceptibility to cancer. Exercise may increase number and activity of macrophages, lymphokine-activated killer cells and their regulating cytokines; it may increase mitogen-induced lymphocyte proliferation. Strenuous exercise increases the production of free radicals, whereas chronic exercise improves free radical defenses by up-regulating both the activities of free scavenger enzymes and antioxidant levels. Increased exercise may decrease levels of insulin and bioavailable IGF-I, both of which enhance division of normal cells and inhibit cell death.

Definitions adapted from the World Cancer Research Fund and American Institute for Cancer Research<sup>(1)</sup>. Convincing evidence is defined as evidence that is conclusive; probable evidence indicates evidence is strong enough to conclude that a causal relation is likely; possible evidence indicates a causal relation may exist; insufficient evidence indicates evidence is suggestive but too sparse to make a more definitive judgment. NE not examined, IGF: Insulin-like growth factors

primary cancers. The results generally show that the higher physical activity is associated with lower rate of breast and colon cancer recurrences, cancer specific mortality and all causes of mortality.

### Exercise program and prescriptions

The American College of Sports Medicine (ACSM) recommends that an exercise prescription consist of five components: frequency, intensity, time, type (FITT Principle) and progression.

Table 3 summarizes the FITT principle of exercise prescription in clinical programs.

These components should be used to prescribe health related exercise training program for both healthy and chronically ill population to include patients with cancer on treatment. Suggestions to prescribe aerobic exercises for patients in early stage of cancer have been published and to date no one has published guidelines for resistance or flexibility exercise protocol in patient with cancer or recovering from cancer. It is recommended that patients should undergo a symptom-limited graded exercise test, which serves as a basis for exercise prescription. Patients present physical status and the individual's current phase of treatment or recovery must also be considered.

**Table 2: Systematic reviews of physical activity in cancer survivors, published 2004–2008**

Study	Studies reviewed	Authors' conclusions
Kirshbaum, 2007 <sup>[66]</sup>	Systematic review of 29 intervention and observational studies in breast cancer survivors	Affirmation of the central proposition that exercise seems to be beneficial and safe for a variety of breast cancer patients continues.
Markes <i>et al.</i> 2006 <sup>[67]</sup>	Systematic review and meta-analysis of 9 controlled trials in breast cancer survivors during adjuvant therapy	Improvement can be expected in physical fitness and the resulting capacity for performing activities of daily life. An improvement for other outcomes is still tenable.
McNeely <i>et al.</i> 2006 <sup>[68]</sup>	Systematic review and meta-analysis of 14 randomized trials in breast cancer survivors	Exercise is an effective intervention to improve quality of life, cardio respiratory fitness, physical functioning, and fatigue.
Conn <i>et al.</i> 2006 <sup>[69]</sup>	Systematic review and meta-analysis of 30 intervention studies in cancer survivors	Exercise interventions resulted in small positive effects on health and well-being outcomes.
Schmitz <i>et al.</i> 2005 <sup>[70]</sup>	Systematic review and meta-analysis of 32 controlled trials in cancer survivors	Physical activity improves cardio respiratory fitness during and after cancer treatment, symptoms and physiologic effects during treatment, and vigor post-treatment.
Knols <i>et al.</i> 2005 <sup>[71]</sup>	Systematic review of 34 controlled trials in cancer survivors	Cancer patients may benefit from physical exercise both during and after treatment.
Douglas, 2005 <sup>[72]</sup>	Systematic review of 21 intervention studies in cancer survivors	There is a growing body of evidence to justify the inclusion of exercise programs in the rehabilitation of cancer patients returning to health after treatment.
Galvao and Newton, 2005 <sup>[73]</sup>	Systematic review of 26 intervention studies in cancer survivors	Preliminary positive physiologic and psychological benefits from exercise when undertaken during or after traditional cancer treatment.
Stevinson <i>et al.</i> 2004 <sup>[74]</sup>	Systematic review and meta-analysis of 33 controlled trials in cancer survivors	Exercise interventions for cancer patients can lead to moderate increases in physical function and are not associated with increased symptoms of fatigue.
Oldervoll <i>et al.</i> 2004 <sup>[75]</sup>	Systematic review of 12 randomized controlled trials in cancer survivors	Cancer patients benefit from maintaining physical activity balanced with efficient rest periods

**Table 3: FITT exercise prescription for apparently healthy individuals**

FITT principal components	Frequency	Intensity	Time	Type
Cardio respiratory (Aerobic training)	3 to 5 days per week	40% or 50-85% HRR 40% or 50-85% VO <sub>2</sub> R 55% or 65-90% HRmax	20 to 60 minutes	Dynamic use of large muscle groups
Muscular strength and endurance (Resistance training)	2 to 3 days per week	12-16 RPE	1 set of 3 to 20 repetitions (e.g. 3 to 5, 8 to 10, 12 to 15)	8 to 10 exercises (All major muscles)
Flexibility	2 to 7 days per week	Stretch to tightness at the end of the range of motion but not to pain	15 to 30 seconds 2 to 4 times/ stretch	Static stretches (All major muscles)

FITT = frequency, intensity, time, type; HRR = heart rate reserve; HR max = maximum heart rate; RPE = rate of perceived exertion; VO<sub>2</sub>R = maximum oxygen consumption reserve. Data adapted from ACSM.

### Type or mode of exercise

The main health related types of exercise are aerobic, resistance and flexibility. The best mode of exercise depends on the person's goals, health status, and exercise history and cancer experience. Aerobic training is defined as a method of improving cardio respiratory system (e.g. Cycling, walking)


Resistance training is defined as a method to maintain or improve muscular strength, endurance or power, which is performed against relatively high resistance and few repetitions. Resistance exercise is a potent physiological intervention to increase muscle mass and attenuate muscle wasting. Flexibility training is defined as a method of maintaining or improving length of the muscle.

Radiation and chemotherapy may cause scar formation in joint, which may result in limitation in range of motion and this limitation can be prevented and normal range of motion can be gained by flexibility training. The best mode of exercise for the patient with cancer on treatment has not been determined because of the lack of research. However, no mode of exercise has been determined to be harmful to patient with cancer on treatment either.

### Frequency

American College of Sports Medicine recommends apparently healthy individual to engage in aerobic training 3-5 days/week. In a deconditioned population, however, several shorter exercise sessions per day are generally better

**When: cancer-related time period**



**Figure 1:** Framework PEACE: an organizational model for examining when and how physical exercise may affect the cancer experience. [Adapted from Courneya K.S, Friedenreich C.M.2]

tolerated. It further recommends individuals to undertake two to three non-consecutive days per week of resistance training. Flexibility training ranges from two to three days/week up to five to seven days/week.<sup>[78]</sup>

**Intensity**

The intensity recommended for aerobic exercise is 40-85% of maximum heart rate calculated by Karvonen method, which calculates % of heart rate reserve.

**Karvonen formula**

Age predicted maximum heart rate (APMHR) = 220 – age.

Target heart rate range = [(APHMR – resting heart rate) × per cent intensity] + resting heart rate.

However, any medication that affect’s heart rate, for e.g. Beta-blockers, invalidates the formula. For these patient subjective means of gauging intensity such as rate of perceived exertion (RPE) assessed by BORG Scale can be used.<sup>[79]</sup> These individuals should be motivated or advised to exercise on intensity between 12 and 16 (somewhat hard to hard) of Borg scale. For resistance exercise, 50 -70% of 1- repetition maximum in two or three sets with 8-12 repetitions per set has been shown effective. The 1- repetition maximum refers to maximum load that a person able to lift once. Lower range of intensity is recommended for older and debilitated survivors and higher range is recommended for apparently healthy survivors. Survivors who are confined to bed or who

fatigue with mild exertion may not be candidates for recommended intensity aerobic training but they may benefit from low level of physical activity. These survivors require supervision in early stages of their recovery by a physiotherapist. These severely compromised survivors may benefit from range of motion exercises and gentle resistance training within their tolerance levels in early stage of rehabilitation. The low level training will allow them to gradually build up their tolerance for activity. These survivors in early stages of recovery may later progress to short bouts walking or bicycling several times per day in order to gradually build endurance and allow them to advance to moderate intensity aerobic exercise.

For persons undergoing chemotherapy or radiation treatment the goals of exercise is to maintain function and prevent loss of endurance and strength these survivors may be benefited from routine physiotherapy and occupational therapy. Brisk walking and static cycling are some safe mode of aerobic exercise. Machine resistance and or free weights are used for resistance exercise of large muscle groups of lower and upper extremities.

**Duration of exercise**

American College of Sports Medicine recommends that apparently healthy cancer survivors should exercise aerobically between 20-60 minutes, lower range for less fit and old and duration increases according to fitness and age.<sup>[78]</sup> Resistance training should be less than 60 minutes for whole bodywork. Flexibility training is given for two

to four repetitions with each stretch holding for about 15-30 seconds.<sup>[78]</sup>

### Exercise progression

It depends on many variables such as the person's goals, exercise tolerance and age. Things to remember about progression are that every person will adopt differently to exercise stimulus, hence the rate of adaptation dictate the rate of progression. Treatment for cancer progresses far less predictably, and often non-linearly, because of multiple factors including the treatment schedule, fluctuating blood counts and varying symptom experiences. Definitive guidelines for progression exist except for patient with early stage disease. The guidelines state 1) Frequency and duration should be increased before intensity. 2) Progression should be slower and more gradual for the deconditioned patient and those who are experiencing severe side effects of treatment.

### Contraindications to exercise

Medical screening should be conducted for all survivors prior to their participation in an exercise program. General contraindications to exercises are cardiovascular insufficiency (e.g. uncontrolled symptomatic heart failure, acute myocarditis, and recent myocardial infarction), acute infectious diseases, metabolic diseases (e.g. thyrotoxicosis, myxedema), mental or physical impairment leading to inability to exercise.<sup>[78]</sup> In addition to general contraindication certain contraindication and precautions are specific to cancer survivors they are; Exercise within two hours of chemotherapy or radiation therapy as increased circulation may increase the effects of treatment, Intravenous chemotherapy within previous 24 hours is also a contraindication for cancer survivors.<sup>[80]</sup> Survivors with anemia (Hemoglobin <8g/dl) should not exercise until anemia is improved (Hemoglobin >10g/dl),<sup>[81]</sup> Bedside exercise programs may be prescribed for these individuals with frequent and short sessions. Hematological values where Absolute Neutrophil count less than  $0.5 \times 10^9/\mu\text{l}$  and Platelet count less than  $50 \times 10^9/\mu\text{l}$  is contraindication for exercise as well.<sup>[81]</sup> Acute onset of nausea during exercise and vomiting within previous 24-36 hours, unusual fatigability or muscular weakness, disorientation, blurred vision, faintness, pallor, night pain or pain not associated with injury are also signs of contraindication to exercise.<sup>[80]</sup> Survivors with immunosuppressants should avoid public gyms until their white blood cell count return to safe level ( $>500/\text{mm}^3$ ),<sup>[80]</sup> bone marrow transplanted survivors should avoid exposure to public places with risk of microbial contamination for one year after transplantation. Survivors with indwelling catheter should avoid resistance exercise of muscle in the area to avoid dislodgement of catheter. Survivors with significant peripheral neuropathies should avoid exercise of the part because of weakness

or loss of balance, stationary bicycle may be used in this situation.<sup>[81]</sup>

### Adherence to exercise

The studies that reported adherence to the exercise program obtained a high level of adherence and a low dropout rate.<sup>[81-86]</sup> Adherence rate after completion of cancer treatment was 95% or higher and during active therapies was between 72-86%, which is still in acceptable level. Lower adherence rate are expected during active treatment when the patients are experiencing more symptoms. Structured exercise protocol might benefit these patients with many or severe symptoms and who struggle to regain their normal function such as those unable to resume work, or who suffer from chronic fatigue and low physical function after the end of treatment. Some cancer survivors can adopt an exercise program independently, many will benefit from referral to physiotherapist who can give recommendation of exercise program on type, frequency, duration, and intensity based on survivors age, previous fitness level, type of cancer, stage of treatment, type of therapy, comorbid conditions. Physiotherapy is essential for survivors with injuries, pain or specific post surgical conditions such as lymphedema or amputation.

The study by Pichett M *et al.* suggests that individual who lead sedentary lifestyle may benefit from structured exercise programs that include information and support related to exercise adherence strategy.<sup>[87]</sup> Counseling patients is one such strategy that effectively increased adherence to exercise and increased physical activity in general practice.<sup>[88]</sup> Most survivors preferred that their Oncologist initiate the discussion of exercise and such discussion also appears to increase exercise level during treatment.<sup>[89]</sup> Cancer survivors have unique and varied exercise counseling and programming preference. In the study by Jones W.L (307 survivors) 98% preferred recreational exercises, 81% preferred walking, 57% preferred unsupervised exercise.<sup>[90]</sup> These preferences for individual survivors should be considered before exercise program prescription.

Providing reassurance that exercise is safe and beneficial modality may improve exercise adherence for inactive cancer survivors and exercise program prescribed should build confidence by slowly increasing the intensity. It should be noted that adherence to exercise program is necessary to obtain improved aerobic fitness. A point to be considered is transfer of local exercise training into activities of daily living for example, resistance exercise may improve muscle strength, endurance and physical functioning but it is known that without integration of functional training improved muscle strength does not result in improved

functional task performance for efficient performance of activities of daily living.<sup>[91]</sup> An individual must be able to perform basic movement and also combination of these in order to accomplish more complex tasks,<sup>[92]</sup> sports may provide training in such complex tasks. Sports are often included in exercise program to facilitate integration into daily life, as it is difficult to become physically active when sedentary.<sup>[93]</sup> Enjoyment of sports has also been reported to facilitate for adaptation of an active life style.<sup>[94]</sup> Sports might also have beneficial effect on physical activity level and physical health, develop sports specific skills, provide a sense of achievement and empowerment, develop self esteem and teach self discipline.<sup>[95]</sup> Preliminary trails on lifestyle intervention (incorporating short periods of moderate activity into their daily routine) are going on for cancer survivors and these studies have shown promising effect on improving physical functioning and quality of life and increasing physical activity.<sup>[96]</sup>

The health benefits of physical activity are independent of whether the physical activity is sport, household, occupational or recreational in nature. Some of the ways to increase physical activity that can be advised are – using stairs rather than an elevator, always walking to the destination when possible, exercising with friends and family, taking a ten minutes exercise break to stretch and quick walk, walking to visit nearby friends or co workers instead of calling them over phone, planning for active vacations rather than only driving trips, using a stationary bicycle while watching TV, planning the exercise routine to gradually increase the days per week and minutes per session.

#### Adverse event issues

Of the reviewed studies 14 commented on the presence or absence of adverse events during the period of intervention. In the 14 studies 12 indicated that no harm was observed as a result of exercise during or after cancer treatment. McNeely *et al.* reported that one participant complained of nausea during one session, with no further difficulties.<sup>[83]</sup> Courneya *et al.* noted that three participants developed lymphedema out of which two had undergone axillary irradiation, a strong risk factor for lymphedema.<sup>[84]</sup> The author commented that it was not clear whether the onset of lymphedema was due to exercise.

#### Exercise and diet

Overweight and obesity have been associated with many types of cancer, the ideal method to limit weight gain or loss weight is to unbalance the energy equation by combination of both diet and increasing physical activity. The expert group of WHO concluded that limiting weight gain during adult life, thereby avoiding overweight and obesity

reduces the risk of postmenopausal breast cancer, colon cancer, endometrial cancer, kidney (renal cell), esophagus (adenocarcinoma) and thyroid cancer.<sup>[83]</sup> The expert committee of American Cancer Society has concluded that increasing vegetable and fruits, increasing fiber, omega 3 fatty acid, soy and limiting total fat and saturated fat have possible benefit on preventing some cancer recurrence and overall survival,<sup>[97]</sup> but the information available is insufficient to conclude the benefits for some sites. Diet with this recommendation is recommended which dietitian must individualize as food intake may be compromised by the effects of disease or therapy and to achieve specific goals of individual exercise program. It should also be noted that benefits of exercise are independent of weight loss and diet.

---

## DISCUSSION

The study suggests that there is strong evidence for reduced risk of some cancers with increasing physical activity. The strongest evidence exists for colorectal and postmenopausal breast cancer with possible association for prostate, endometrial and lung cancer. The findings are supported by identified biological mechanisms. The field of oncology will benefit from understanding the importance of physical activity both for primary prevention as well as in helping cancer survivors cope with and recover from treatments, improve the health of long term cancer survivors and possibly even reduce the risk of recurrence and extend survival after a cancer diagnosis. However, an understanding of the amount, type, and intensity of activity needed has not been fully elucidated for primary prevention and for patients at different stages of disease progression is still lacking. There is sufficient evidence already to recommend that at least moderate intensity activity of 30 minutes/day for five days/week or more than 45-60 minutes vigorous activities for some cancer site is given. There is sufficient evidence to promote exercise in cancer survivors following careful assessment and tailoring on exercise prescription based on health status of individual. Additional studies will be needed to more firmly establish physical activity benefits to cancer survivors.

---

## ACKNOWLEDGMENTS

We thank our institute (M.T.P.G and R.I.H.S) for setting the stage for our performance and we thank Librarian Kumaravel C for his co-operation.

---

## REFERENCES

1. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74-108.

2. Courneya KS, Friedenreich CM. Framework PEACE: An organizational model for examining physical exercise across the cancer experience. *Ann Behav Med* 2001;23:263-72.
3. Vainio H, Bianchine F. *Wight control and physical activity*. Lyon: IARC Pres; 2008.
4. Chao A, Connell CJ, Jacobs EJ, McCullough ML, Patel AV, Calle EE, *et al*. Amount, type and timing of recreational Physical activity in relation to colon and rectal cancer in older adults: the cancer prevention study Nutrition cohort. *Cancer Epidemiol Biomarkers Prev* 2004;13:2187-95.
5. Cheville. A rehabilitation of patients with advanced cancer. *Cancer* 2001;92:1039-1048.
6. Knoff MT, Musanti R, Dorward J. Exercise and QOL outcomes in patients with cancer. *Semin oncol Nurs* 2007;23:285-29.
7. Mock V, Pickett M, Ropka ME, Muscari Lin E, Stewart KJ, Rhodes VA, *et al*. Fatigue and quality of life outcomes of exercise during cancer treatment. *Cancer Pract* 2001;9:119-27.
8. Schmitz KH, Holtzman J, Courneya KS, Mâsse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2005;14:1588-95.
9. Ingram C, Courneya KS, Kingston D. The effects of exercise on body weight and composition in breast cancer survivors. *Oncol Nurs Forum* 2006;33:937-50.
10. Visovsky C. Muscle strength, body composition and physical activity in women receiving chemotherapy for breast cancer. *Integr Cancer Ther* 2006;5:183-9.
11. McTiernan A. Physical activity after cancer: physiologic outcomes. *Cancer Invest* 2004;22:68-81.
12. Winningham ML, MacVicar MG, Bondoc M, Anderson JI, Minton JP. Effect of aerobic exercise on body weight and composition in patients with breast cancer on adjuvant chemotherapy. *Oncol Nurs Forum* 1989;16:683-9.
13. Courneya KS, Friedenreich CM, Sela RA, Quinney HA, Rhodes RE, Handman M. The group psychotherapy and home-based physical exercise (Group-Hope) trial in cancer survivors: Physical fitness and quality of life outcomes. *Psychooncology* 2003;12:357-74.
14. Andersen C, Adamsen L, Moeller T, Midtgaard J, Quist M, Tveteraas A, *et al*. The effect of a multidimensional exercise programme on symptoms and side effects in cancer patients undergoing chemotherapy, The use of semi-structural diaries. *Eur J Oncol Nurs* 2006;10:247-62.
15. National coalition for cancer survivorship (Homepage on the internet 3. Glossary C.1994-2004. Avariable from. [http://www.Canceradvocacyorg. Resources/glossary asp top](http://www.Canceradvocacyorg.Resources/glossary.asp). [cited on 2008 Apr].
16. Wannamethee SG, Shaper AG, Walker M. Physical activity and risk of cancer in middle-aged men. *Br J Cancer* 2001;85:1311-6.
17. Colbert LH, Hartman TJ, Malila N, Limburg PJ, Pietinen P, Virtamo J, *et al*. Physical activity in relation to cancer of the colon and rectum in a cohort of male smokers. *Cancer Epidemiol Biomarkers Prev* 2001;10:265-8.
18. Nilsen TI, Vatten LJ. Prospective study of colorectal cancer risk and physical activity, diabetes, blood glucose and BMI: exploring the hyperinsulinaemia hypothesis. *Br J Cancer* 2001;84:417-22.
19. Boutron-Ruault MC, Senesse P, Méance S, Belghiti C, Faivre J. Energy intake, body mass index, physical activity, and the colorectal adenoma-carcinoma sequence. *Nutr Cancer* 2001;39:50-7.
20. Moradi T, Adami HO, Ekblom A, Wedrén S, Terry P, Floderus B, *et al*. Physical activity and risk for breast cancer: a prospective cohort study among Swedish twins. *Int J Cancer* 2001;100:76-81.
21. Moradi T, Nyrén O, Zack M, Magnusson C, Persson I, Adami HO. Breast cancer risk and lifetime leisure-time and occupational physical activity. *Cancer Causes Contro* 2000;11:523-31.
22. Adams-Campbell LL, Rosenberg L, Rao RS, Palmer JR. Strenuous physical activity and breast cancer risk in African-American women. *J Natl Med Assoc* 2001;93:267-75.
23. Gilliland FD, Li YF, Baumgartner K, Crumley D, Samet JM. Physical activity and breast cancer risk in Hispanic and non-Hispanic white women. *Am J Epidemiol* 2001;154:442-50.
24. Matthews CE, Shu XO, Jin F, Dai Q, Hebert JR, Ruan ZX, *et al*. Lifetime physical activity and breast cancer risk in the Shanghai Breast Cancer Study. *Br J Cancer* 2001;84:994-1001.
25. Friedenreich CM, Bryant HE, Courneya KS. Case control study of lifetime physical activity and breast cancer risk. *Am J Epidemiol* 2001;154:336-47.
26. Lee IM, Rexrode KM, Cook NR. Physical activity and breast cancer risk: the Women's Health Study. *Cancer Causes Control* 2001;12:137-45.
27. Dirx MJ, Voorrips LE, Goldbohm RA, van den Brandt PA. Baseline recreational physical activity, history of sports participation, and postmenopausal breast carcinoma risk in the Netherlands Cohort Study. *Cancer* 2001;92:1638-49.
28. Verloop J, Rookus MA, van der Kooy K, van Leeuwen FE. Physical activity and breast cancer risk in women aged 20-54 years. *J Natl Cancer Inst* 2000;92:128-35.
29. Wyshak G, Frisch RE. Breast cancer among former college athletes compared to non-athletes: a 15-year follow-up. *Br J Cancer* 2000;82:726-30.
30. Hofvind SS, Thoresen SO. Physical activity and breast cancer. *Tidsskr Nor Laegeforen* 2001;121:1892-5.
31. Drake DA. A longitudinal study of physical activity and breast cancer prediction. *Cancer Nurs* 2001;24:371-7.
32. Lee IM, Cook NR, Rexrode KM, Buring JE. Lifetime physical activity and risk of breast cancer. *Br J Cancer* 2001;85:962-5.
33. Moore DB, Folsom AR, Mink PJ, Hong CP, Anderson KE, Kushi LH. Physical activity and incidence of postmenopausal breast cancer. *Epidemiology* 2000;11:292-6.
34. Luoto R, Latikka P, Pukkala E, Hakulinen T, Vihko V. The effect of physical activity on breast cancer risk: a cohort study of 30,548 women. *Eur J Epidemiol* 2000;16:973-80.
35. Hoffman-Goetz L, Apter D, Demark-Wahnefried W, Goran MI, McTiernan A, Reichman ME. Possible mechanisms mediating an association between physical activity and breast cancer. *Cancer* 1998;83:621-8.
36. Toniolo PG. Endogenous estrogens and breast cancer risk: the case for prospective cohort studies. *Environ Health Perspect* 1997;105:587-92.
37. Friedenreich CM. Review of anthropometric factors and breast cancer risk. *Eur J Cancer Prev* 2000;10:15-32.
38. Lund Nilsen TI, Johnsen R, Vatten LJ. Socio-economic and lifestyle factors associated with the risk of prostate cancer. *Br J Cancer* 2000;82:1358-63.
39. Norman A, Moradi T, Gridley G, Dosemeci M, Rydh B, Nyrén O, *et al*. Occupational physical activity and risk for prostate cancer in a nationwide cohort study in Sweden. *Br J Cancer* 2002;86:70-5.
40. Clarke G, Whittemore AS. Prostate cancer risk in relation to anthropometry and physical activity: the National Health and Nutrition Examination Survey. I: epidemiological follow-up study. *Cancer Epidemiol Biomarkers Prev* 2000;9:875-81.
41. Bairati I, Larouche R, Meyer F, Moore L, Fradet Y. Lifetime occupational physical activity and incidental prostate cancer. *Cancer Causes Control* 2000;11:759-64.
42. Liu S, Lee IM, Linson P, Ajani U, Buring JE, Hennekens CH. A prospective study of physical activity and risk of prostate cancer in US physicians. *Int J Epidemiol* 2000;29:29-35.
43. Lee IM, Sesso HD, Paffenbarger RS. A prospective cohort study of physical activity and body size in relation to prostate cancer risk. *Cancer Causes Control* 2001;12:187-93.
44. Putnam SD, Cerhan JR, Parker AS, Bianchi GD, Wallace RB, Cantor KP, *et al*. Lifestyle and anthropometric risk factors for prostate cancer in a cohort of Iowa men. *Ann Epidemiol*



- 2000;10:361-9.
45. Lacey JV Jr, Deng J, Dosemeci M, Gao YT, Mostofi FK, Sesterhenn IA, *et al.* Prostate cancer, benign prostatic hyperplasia and physical activity in Shanghai, China. *Int J Epidemiol* 2001;30:341-9.
  46. Friedenreich CM, Thune I. A review of physical activity and prostate cancer risk. *Cancer Causes Control* 2001;12:461-75.
  47. Moradi T, Weiderpass E, Signorello LB, Persson I, Nyrén O, Adami HO. Physical activity and postmenopausal endometrial cancer risk. *Cancer Causes Control* 2000;11:829-37.
  48. Littman AJ, Voigt LF, Beresford SA, Weiss NS. Recreational physical activity and endometrial cancer risk. *Am J Epidemiol* 2001;154:924-33.
  49. Cottreau CM, Ness RB, Kriska AM. Physical activity and reduced risk of ovarian cancer. *Obstet Gynecol* 2000;96:609-14.
  50. Bertone ER, Willett WC, Rosner BA, Hunter DJ, Fuchs CS, Speizer FE, *et al.* Prospective study of recreational physical activity and ovarian cancer. *J Natl Cancer Inst* 2001;93:942-8.
  51. Tavani A, Gallus S, La Vecchia C, Dal Maso L, Negri E, Pelucchi C, *et al.* Physical activity and risk of ovarian cancer: an Italian case-control study. *Int J Cancer* 2001;91:407-11.
  52. Kubík AK, Zatloukal P, Tomásek L, Petruzelka L. Lung cancer risk among Czech women: a case-control study. *Prev. Med* 2002;34:436-44.
  53. Colbert LH, Hartman TJ, Tangrea JA, Pietinen P, Virtamo J, Taylor PR, *et al.* Physical activity and lung cancer risk in male smokers. *Int J Cancer* 2002;98:770-3.
  54. Srivastava A, Kreiger N. Relation to physical activity to risk of testicular cancer. *Am J Epidemiol* 2000;151:78-87.
  55. Bergström A, Terry P, Lindblad P, Lichtenstein P, Ahlbom A, Feychting M, *et al.* Physical activity and risk of renal cell cancer. *Int J Cancer* 2001;92:155-7.
  56. Michaud DS, Giovannucci E, Willett WC, Colditz GA, Stampfer MJ, Fuchs CS. Physical activity, obesity, height, and the risk of pancreatic cancer. *J Am Med Assoc* 2001;286:921-9.
  57. Hanley AJ, Johnson KC, Villeneuve PJ, Mao Y; Canadian Cancer Registries Epidemiology Research Group. Physical activity, anthropometric factors and risk of pancreatic cancer: results from the Canadian enhanced cancer surveillance system. *Int J Cancer* 2001;94:140-7.
  58. Rossing MA, Remler R, Voigt LF, Wicklund KG, Daling JR. Recreational physical activity and risk of papillary thyroid cancer. *Cancer Causes Control* 2001;12:881-5.
  59. Shors AR, Solomon C, McTiernan A, White E. Melanoma risk in relation to height, weight, and exercise. *Cancer Causes Control* 2001;12:599-606.
  60. Yu H, Rohan T. Role of the insulin-like growth factor family in cancer development and progression. *J Natl Cancer Inst* 2000;92:1472-89.
  61. Giovannucci E. Insulin, insulin-like growth factors and colon cancer: a review of the evidence. *J Nutr* 2001;131:3109-20.
  62. Duclos M. Effects of physical training on endocrine functions. *Ann Endocrinol* 2001;62:19-32.
  63. IARC Working Group IARC Handbook of Cancer Prevention, Weight Control and Physical Activity. Vol. 6. Lyon: IARC; 2002.
  64. Marrett L, Theis B, Ashbury F. An Expert Panel Workshop report: physical activity and cancer prevention. *Chronic Dis Can* 2000;21:143-9.
  65. Shephard RJ, Rhind S, Shek PN. The impact of exercise on the immune system: NK cells, interleukins 1 and 2, and related responses. *Exerc Sport Sci Rev* 1995;23:215-41.
  66. Kirshbaum MN. A review of the benefits of whole body exercise during and after treatment for breast cancer. *J Clin Nurs* 2007;16:104-21.
  67. Markes M, Brockow T, Resch KL. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database of Syst Rev* 2006;18:CD005001.
  68. McNeely ML, Campbell KL, Rowe BH, Klassen TP, Mackey JR, Courneya KS. Effects of exercise on breast cancer patients and survivors: a systematic review and meta-analysis. *CMAJ* 2006;175:34-41.
  69. Conn VS, Hafdahl AR, Porock DC, McDaniel R, Nielsen PJ. A meta-analysis of exercise interventions among people treated for cancer. *Support Care Cancer* 2006;14:699-712.
  70. Schmitz KH, Holtzman J, Courneya KS, Mâsse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2005;14:1588-95.
  71. Knols R, Aaronson NK, Uebelhart D, Fransen J, Aufdemkampe G. Physical exercise in cancer patients during and after medical treatment: a systematic review of randomized and controlled clinical trials. *J Clin Oncol* 2005;23:3830-42.
  72. Douglas E. Exercise in cancer patients. *Phys Ther Rev* 2005;10:71-88.
  73. Galvao DA, Newton RU. Review of exercise intervention studies in cancer patients. *J Clin Oncol* 2005;23:899-909.
  74. Stevinson C, Lawlor DA, Fox KR. Exercise interventions for cancer patients: systematic review of controlled trials. *Cancer Causes Control* 2004;15:1035-56.
  75. Oldervoll LM, Kaasa S, Hjermstad MJ, Lund JA, Loge JH. Physical exercise results in the improved subjective well-being of a few or is effective rehabilitation for all cancer patients? *Eur J Cancer* 2004;40:951-62.
  76. Oldervoll LM, Loge JH, Paltiel H, Asp MB, Vidvei U, Wiken AN, *et al.* The effect of physical exercise program is management palliative care. *J Pain Symptom Manage* 2006;1:421-46.
  77. Coben SR, Leis A. What determine qualities of life QOL of terminally ill cancer patient from their own perspective? *J Palliative Care* 2002;18:48-58.
  78. American College of Sport Medicine. ACSM's guidelines for exercise testing and Prescription Ed T. Philadelphia PA: Lippincott Williams and Wilkin; 2006.
  79. Borg GA. Psychological basis of physical exertion. *Med Sci Sports Exerc* 1982;14:377.
  80. Winningham ML, MacVicar MG, Burke MB. Exercise for cancer patients: guidelines and precautions. *Physician Sports Med* 1986;14:125-32.
  81. Courneya KS, Mackey JR, Jones LW. Coping with cancer: can exercise help? *Phys Sportsmed* 2000;28:49-73.
  82. Segal R, Evans W, Johnson D, Smith J, Colletta S, Gayton J, *et al.* Structured exercise improves physical functioning in women with stages I and II breast cancer: results of a randomized controlled trial. *J Clin Oncol* 2001;19:657-65.
  83. McNeely ML, Parliament M, Courneya KS, Seikaly H, Jha N, Scrimger R, *et al.* A pilot study of a randomized controlled trial to evaluate the effects of progressive resistance training on shoulder dysfunction caused by spinal accessory neuropraxia, neurectomy in head and neck cancer survivors. *Head Neck* 2004;20:518-30.
  84. Courneya KS, Mackey JR, Bell GJ, Jones LW, Field CJ, Fairey AS. Randomized controlled trial of exercise training in postmenopausal breast cancer survivors: cardiopulmonary and quality of life outcomes. *J Clin Oncol* 2003;21:1660-8.
  85. Burnham TR, Wilcox A. Effects of exercise on physiological and psychological variables in cancer survivors. *Med Sci Sports Exerc* 2002;34:1863-7.
  86. Dimeo FC, Stieglitz RD, Novelli-Fischer U, Fetscher S, Keul J. Effects of physical activity on the fatigue and psychologic status of cancer patients during chemotherapy. *Cancer* 1999;85:2273-7.
  87. Pickett M, Mock V, Ropka ME, Cameron L, Coleman M, Podewils L. Adherence to moderate-intensity exercise during breast cancer therapy. *Cancer Pract* 2002;10:284-92.
  88. Elley CR, Kerse N, Arroll B, Robinson E. Effectiveness of counseling patients on physical activity in general practice: cluster randomised controlled trial. *BMJ* 2003;326:793.
  89. Jones LW, Courneya KS. Exercise discussions during cancer treatment consultations. *Cancer Pract* 2002;10:66-74.

90. Jones LW, Courneya KS. Exercise counseling and programming preferences of cancer survivors. *Cancer Pract* 2002;10:208-15.
91. de Vreede PL, Samson MM, van Meeteren NL, Duursma SA, Verhaar HJ. Functional-task exercise versus resistance strength exercise to improve daily function in older women: a randomized controlled trial. *J Am Geriatr Soc* 2005;53:2-10.
92. Schwartz AL. Physical activity after a cancer diagnosis: psychosocial outcomes. *Cancer Invest* 2004;22:82-92.
93. van der Ploeg HP, Streppel KR, van der Beek AJ, van der Woude LH, Vollenbroek-Hutten MM, van Harten WH, *et al.* Counselling increases physical activity behaviour nine weeks after rehabilitation. *Br J Sports Med* 2006;40:223-9.
94. Lewis BA, Marcus BH, Pate RR, Dunn AL. Psychosocial mediators of physical activity behavior among adults and children. *Am J Prev Med* 2002;23:26-35.
95. Jackson NW, Howes FS, Gupta S, Doyle JL, Waters E. Interventions implemented through sporting organisations for increasing participation in sport. *Cochrane Database Syst Rev* 2005;2:CD004812.
96. Basen-Engquist K, Taylor CL, Rosenblum C, Smith MA, Shinn EH, Greisinger A, *et al.* Randomized pilot test of a life style physical activity intervention for breast cancer survivors. *Patient Educ Couns* 2006;64:225-34.
97. Brown JK, Byers T, Doyle C, Coumeya KS, Demark-Wahnefried W, Kushi LH, *et al.* Nutrition and physical activity during and after cancer treatment: An American Cancer Society Guide for Informed Choices. *CA Cancer J Clin* 2003;53:268-91.

**Source of Support:** Nil, **Conflict of Interest:** None declared.

### Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than **2048 kb (2 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.