# Dentists' knowledge and practice regarding prevention of infective endocarditis

Soukaina Ryalat<sup>1</sup>, Yazan Hassona<sup>1</sup>, Mohammad Al-Shayyab<sup>1</sup>, Mais Abo-Ghosh<sup>2</sup>, Faleh Sawair<sup>1</sup>

> <sup>1</sup>Department of Oral and Maxillofacial Surgery, Oral Medicine, and Periodontics, Faculty of Dentistry, The University of Jordan, Amman, Jordan, <sup>2</sup>Internist at the Department of Oral and Maxillofascial Surgery, Faculty of Dentistry, The University of Jordan, Amman, Jordan

**Correspondence:** Dr. Soukaina Ryalat Email: salryalat@yahoo.com

#### **ABSTRACT**

**Objectives:** The objective of this study is to assess the knowledge and practice of dentists from Jordan, regarding prevention of infective endocarditis (IE) in dental practice. **Materials and Methods:** A sample of Jordanian dentists was interviewed regarding their IE knowledge and practice using a validated and pretested survey instrument. **Results:** Most of the dentists have encountered a patient with IE who needed prophylactic antibiotic (PA) and have prescribed antibiotics to prevent IE. Jordanian dentists' approach to patients in need for PA varied between the National Institute for Health and Clinical Excellence recommendations and the American Heart Association to a lesser degree, but still a relatively high percent (39%) did not know any guidelines to follow although 74% have encountered patients who needed endocarditis prophylaxis. Patients with prosthetic heart valve were ranked on top of medical conditions that required PA (87.4%), and most dentists (94.5%) thought that dental extractions need PA followed by periodontal surgery (88.2%). **Conclusion:** There is a lack of consistency in the knowledge and practice of Jordanian dentists with regard to IE. There is a need to take actions to improve dentist's knowledge regarding this topic.

Key words: Antibiotic choice, antibiotic prophylaxis, dental procedures, dentist's knowledge, infective endocarditis

### **INTRODUCTION**

Infective endocarditis (IE), which is a life-threatening infection of the endocardium or heart valves, is usually associated with cardiac defects, congenital, or acquired.<sup>[1]</sup> Oral cavity, with more than 600 microbiological species have been recorded in the human mouth, is also home to microbial communities with multiple effects in human health and disease.<sup>[2]</sup> Various dental procedures have been linked to the occurrence of IE. Nevertheless, there seems to be a dilemma about which dental procedures and under what circumstances significant risk occurs.<sup>[3]</sup> Several guidelines regarding IE prevention

Access this article online				
Quick Response Code:				
	Website: www.eurjdent.com			

are available globally including, British Society for Antimicrobial Chemotherapy guidelines (BSAC) on 2006, National Institute for Health and Clinical Excellence guidelines (NICE) on 2008, European Society of Cardiology guidelines on 2009, the Australian guidelines, and the American Heart Association guidelines (AHA) on 2007; most of these guidelines recommended a more restrictive use of prophylaxis for IE.<sup>[4-7]</sup> The BSAC guidelines reduced cardiac conditions in which prophylaxis indicated to the presence of prosthetic valves, a history of IE, and some congenital

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Ryalat S, Hassona Y, Al-Shayyab M, Abo-Ghosh M, Sawair F. Dentists' knowledge and practice regarding prevention of infective endocarditis. Eur J Dent 2016;10:480-5.

**DOI:** 10.4103/1305-7456.195158

heart diseases.<sup>[8]</sup> AHA guidelines added another indication, valvulopathy in cardiac transplant recipients. <sup>[9]</sup> NICE guidelines did not recommend any prophylaxis to patients for IE prevention regardless their risk.<sup>[10]</sup>

The data about epidemiology of IE in Jordan are scarce, mostly due to the absence of nationwide studies in this regard. Similarly, little is known about the practice of general dental practitioners in Jordan, regarding prevention of IE and their compliance with international guidelines of IE prophylaxis.

The aim of this study was to assess the knowledge and practice of dentists from Jordan, regarding prevention of IE in dental practice.

### **MATERIALS AND METHODS**

This cross-sectional study was approved by the Faculty of Dentistry Research and Ethics Committee in accordance with the World Medical Declaration of Helsinki. Ethical protocol number (12/2015).

A total of 210 dentists were approached to participate in the study, and 127 agreed to participate. A survey instrument was used to interview participants. The survey instrument included questions about demographic data, knowledge about risk factors of IE, and predisposing dental procedures and their current practice of prescribing antibiotics to at-risk patients [Appendix 1].

Statistical analysis performed using SPSS 16.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics used to describe our study sample. To examine differences between groups, Chi-square test, one-way ANOVA, Student's *t*-test, and Spearman's rho test were used. Results with P < 0.05 considered statistically significant.

### **RESULTS**

The sample was composed of 52.8% males and 47.2% females. Mean age was 33.5 years (range 22–65 years) with 1–40 years of experience (mean 9 years) [Table 1].

As shown in Table 1, most of the respondents were working in private clinics and graduated from Jordanian universities.

Most of the dentists have encountered a patient with IE who needed prophylactic antibiotic (PA) and have prescribed PA to prevent IE, 37% were following the NICE guidelines, and 23.6% were following AHA guidelines. Majority of the respondents (51%) have

prescribed PA when encountering suspected patient to prevent IE, 44% have referred, and only 4.7% have refused to treat these patients [Table 2].

Table 1: Demographic characteristics of participating		
dentists		
Variable	Total, <i>n</i> (%)	
Gender		
Male	67 (52.8)	
Female	60 (47.2)	
Age (years)		
Mean	33.5	
Range	22-65	
SD	10.3	
Experience (years)		
Mean	9.2	
Range	1-40	
SD	9.1	
Place of work		
Private clinic	83 (65.4)	
Hospital-based clinic	31 (24.4)	
Public sector clinic	13 (10.2)	
Country of graduation		
Jordan	59 (46.5)	
Arab countries	45 (35.4)	
Eastern Europe	17 (13.4)	
Western Europe	6 (4.7)	
SD: Standard deviation		

Table 2: Experience with infective endocarditis			
Experience	n (%)		
Encountered a patient who needed PA for IE			
No	33 (26)		
Yes	94 (74)		
Ever prescribed PA to prevent IE			
No	45 (35.4)		
Yes	82 (64.6)		
Encountered a patient who developed IE following dental treatment			
No	104 (81.9)		
Yes	23 (18.1)		
Followed guidelines to prevent IE			
AHA	30 (23.6)		
NICE	47 (37)		
Others	4 (3.1)		
Do not know	46 (36.2)		
What they do when encountering a patient who needs PA to prevent IE**			
prescribe PA	65 (51.2)		
Refer the patient to his physician for consultation	56 (44.1)		
Refuse treating the patient and refer to another colleague	6 (4.7)		

\*\*The decision was not significantly affected by gender, age, experience, country of graduation, or place of work. IE: Infective endocarditis, PA: Prophylactic antibiotic, AHA: American Heart Association, NICE: National Institute for Health and Clinical Excellence

Patients with prosthetic heart valve were ranked on top of medical conditions that required PA (87.4%), and then patients with the previous history of IE, followed by patients with rheumatic heart disease (57.5), the least record was for patients having physiologic murmurs [Table 3].

Regarding dental procedures which needed PA [Table 4], most dentists (87.4%) thought that dental extractions need PA, followed by periodontal surgery (88.2%); however, only 17.3% thought that PA is needed for simple tooth restorations.

Amoxicillin was the antibiotic of choice for 57.5% of dentists to prevent IE, 17.3% prescribed clindamycin; a minority have prescribed lincomycin, metronidazole, gentamicin, and clarithromycin. Around 19% of dentists did not know which antibiotic to prescribe [Table 5].

## DISCUSSION

Dental procedures, especially the invasive ones, are being blamed for causing IE.<sup>[1,3,4]</sup> We aimed in this study to assess the knowledge and practice of dentists in Jordan, regarding prevention of IE in dental practice.

Although the NICE recommended the restriction of antibiotic prophylaxis (AP) use to very few predefined circumstances,<sup>[8]</sup> more than 50% of Jordanian dentists still prescribe PA widely to prevent IE.

Jordanian dentists' approach patients in need for PA varied between the NICE recommendations and the AHA to a lesser degree, but still a relatively high percent (39%) did not know any guidelines to follow although 74% have encountered patients who needed PA to prevent IE.

Only 37% of the respondents followed the NICE guidelines compared with a similar study in London, in which 95% of participants were aware of the NICE guidelines and 77% of them were compliant with those guidelines.<sup>[11]</sup>

This controversy, over the PA prescription for the different dental procedure, is a critical issue among dentists worldwide, which is affected by both the new guidelines and the IE incidence.<sup>[12]</sup>

Prosthetic heart valves and preexisting structural abnormality of the heart are major risk factors for IE,<sup>[13]</sup> a fact which is well appreciated by 87% of Jordanian dentists; comparable results were reported by an Iranian study.<sup>[11]</sup> While 14.2% of Jordanian dentists prescribe PA for cases of physiologic murmurs, 87.2% of their Iranian counterparts prescribe PA for this condition. This might be due to the variations of the universal recommendations or due to lack of knowledge regarding the clinical condition.

Dental procedures that needed PA, due to invasive nature and the risk of bacteremia were extractions and periodontal surgery; majority of Jordanian dentists agree with this fact.<sup>[12,14,15]</sup>

While Amoxicillin was the drug of choice as PA for 57.5% of respondents in this study, it was the drug of choice for 89% of Nigerian dentists.<sup>[16]</sup> In contrast, most

Table 3: Patients who need prophylactic antibiotics to prevent infective endocarditis				
Rank	Patient type	Dentists' thought the patient needs PA to prevent IE n (%)		
1	Patients with prosthetic valve	111 (87.4)		
2	Patients with previous history of infective endocarditis	81 (63.8)		
3	Patients with rheumatic heart disease	73 (57.5)		
4	Patients with history of open heart surgery	64 (50.4)		
5	Patients with stints	62 (48.8)		
6	Patients with cyanotic congenital heart disease	55 (43.3)		
7	Mitral valve prolapsed or regurgitation	53 (41.7)		
8	Cardiac transplantation	50 (39.4)		
9	Patients with pacemaker	47 (37)		
10	Cardiac bypass	41 (32.3)		
11	Patients with ventricular septal defect	38 (29.9)		
11	Patients with a cyanotic congenital heart disease	38 (29.9)		
12	Patients with atrial septal defects	35 (27.6)		
13	Patients with ischemic heart disease	31 (24.4)		
14	Any patient with history of cardiac disease	29 (22.8)		
15	Physiologic murmur	18 (14.2)		
Do not know		6 (4.7)		
15 1.6.10	La contra DA Decela la characteria della de			

IE: Infective endocarditis, PA: Prophylactic antibiotic

Rank	Dental procedure	Dentists' thought the patient needs PA to prevent IE, n (%)
1	Surgical tooth extraction	120 (94.5)
2	Periodontal surgery	112 (88.2)
3	Implants	107 (84.3)
4	Biopsy	83 (65.4)
4	Apicoectomy	83 (65.4)
5	Simple tooth extraction	72 (56.7)
6	Abscess drainage	71 (55.9)
7	Scaling and polishing	70 (55.1)
8	Dentoalveolar trauma	67 (52.8)
9	Reimplantation of avulsed tooth	64 (50.4)
10	Endodontic treatment	58 (45.7)
11	Intraligamentary anesthesia	55 (43.3)
12	Placement of matrix band and wedges	47 (37)
13	Inferior dental nerve block	43 (33.9)
14	Placement of rubber dam	40 (31.5)
15	Tooth preparation and impression taking	28 (22)
16	Placement of fixed orthodontic band	24 (18.9)
17	Simple and complex restorations	22 (17.3)
Do not know		4 (3.1)
IE: Infective endoc	arditis. PA: Prophylactic antibiotic	4 (3.1)

Table 4: Procedures that require the administration of prophylactic antibiotics to prevent infective endocarditis

Table 5: Antibiotics prescribed for prevention of

meetive endocarditis before dental procedures		
Antibiotic	n (%)	
Amoxicillin	73 (57.5)	
Clindamycin	22 (17.3)	
Erythromycin	2 (1.6)	
Metronidazole	1 (0.8)	
Clarithromycin	1 (0.8)	
Gentamycin	1 (0.8)	
Lincomycin	1 (0.8)	
No antibiotic	2 (1.6)	
Do not know	24 (18.9)	

general dentists in Japan prescribe cephams followed by Penicillin.<sup>[17]</sup>

Giving PA is not without harm although it is crucial in certain specific circumstances. Antimicrobial resistance is an emerging issue that must be dealt with to decelerate its progression. PA aim only to prevent IE due to viridans streptococci, and in its best case scenario can prevent up to 10% of all cases of IE.<sup>[18]</sup> Based on these statistics, it is not feasible to give PA to every cardiac patient, and it is worth to adhere with the guidelines to prevent IE in the specific patients, the PA target, without adding on antimicrobial resistance.

It has been reported that all observational studies such as cross-sectional surveys are prone to limitation and bias.<sup>[19]</sup> In this study, the number of participants is limited as a big number of dentists refused to participate; it may be due to their inadequate knowledge about this issue. Moreover, most dentists still prescribe PA in most cardiac patients although it is not recommended anymore in the majority of them; this discrepancy might be due to lack of update on their knowledge about the guidelines.

### CONCLUSION

Although the issue of AP to prevent IE is controversial and no general consensus exists, a large proportion of dentists still do not follow any guidelines. The majority of dentists encountered a patient who needs PA so that the level of knowledge and practice need to be upgraded in Jordan.

# **Financial support and sponsorship** Nil.

#### **Conflicts of interest**

There are no conflicts of interest.

#### REFERENCES

- Jain P, Stevenson T, Sheppard A, Rankin K, Compton SM, Preshing W, et al. Antibiotic prophylaxis for infective endocarditis: Knowledge and implementation of American Heart Association Guidelines among dentists and dental hygienists in Alberta, Canada. J Am Dent Assoc 2015;146:743-50.
- Germano F, Bramanti E, Arcuri C, Cecchetti F, Cicciù M. Atomic force microscopy of bacteria from periodontal subgingival biofilm: Preliminary study results. Eur J Dent 2013;7:152-8.
- 3. Bahammam MA, Abdelaziz NM. Awareness of antimicrobial

prophylaxis for infective endocarditis among dental students and interns at a teaching hospital in Jeddah, Saudi Arabia. Open Dent J 2015;9:176-80.

- Thornhill MH, Lockhart PB, Prendergast B, Chambers JB, Shanson D. NICE and antibiotic prophylaxis to prevent endocarditis. Br Dent J 2015;218:619-21.
- 5. Kazi DS, Bolger AF. Infective endocarditis and antibiotic prophylaxis. Lancet 2015;386:530.
- 6. Di Filippo S. Prophylaxis of infective endocarditis in patients with congenital heart disease in the context of recent modified guidelines. Arch Cardiovasc Dis 2012;105:454-60.
- 7. Diz Dios P. Infective endocarditis prophylaxis. Oral Dis 2014;20:325-8.
- Al-Fouzan AF, Al-Shinaiber RM, Al-Baijan RS, Al-Balawi MM. Antibiotic prophylaxis against infective endocarditis in adult and child patients. Knowledge among dentists in Saudi Arabia. Saudi Med J 2015;36:554-61.
- Ghaderi F, Oshagh M, Dehghani R, Hasanshahi R. Awareness of Iranian's general dentists regarding the latest prophylaxis guideline for prevention of infective endocarditis. J Dent (Shiraz) 2013;14:6-12.
- Lockhart PB, Hanson NB, Ristic H, Menezes AR, Baddour L. Acceptance among and impact on dental practitioners and patients of American Heart Association recommendations for antibiotic prophylaxis. J Am Dent Assoc 2013;144:1030-5.
- 11. Farook SA, Davis AK, Khawaja N, Sheikh AM. NICE guideline and current practice of antibiotic prophylaxis for high risk cardiac

patients (HRCP) among dental trainers and trainees in the United Kingdom (UK). Br Dent J 2012;213:E6.

- 12. Thornhill MH. Infective endocarditis: The impact of the NICE guidelines for antibiotic prophylaxis. Dent Update 2012;39:6-10, 12.
- Dayer MJ, Chambers JB, Prendergast B, Sandoe JA, Thornhill MH. NICE guidance on antibiotic prophylaxis to prevent infective endocarditis: A survey of clinicians' attitudes. QJM 2013;106:237-43.
- Ahmadi-Motamayel F, Vaziri S, Roshanaei G. Knowledge of general dentists and senior dental students in Iran about prevention of infective endocarditis. Chonnam Med J 2012;48:15-20.
- Hashemipour M, Korki M. Antibiotic prophylaxis for bacterial endocarditis: A study of knowledge of guidelines among dentists participated in the 47<sup>th</sup> international congress of dentistry. Dent J Tehran Univ Med Sci 2008;21:210-8.
- Adeyemo WL, Oderinu OH, Olojede AC, Ayodele AO, Fashina AA. Nigerian dentists' knowledge of the current guidelines for preventing infective endocarditis. Community Dent Health 2011;28:178-81.
- Nakano K, Ooshima T. Common knowledge regarding prevention of infective endocarditis among general dentists in Japan. J Cardiol 2011;57:123-30.
- 18. Pallasch T. Antibiotic prophylaxis. Endodontic Topics 2003;4:46-59.
- Al-Shamiri HM, Alaizari NA, Al-Maweri SA, Tarakji B. Knowledge and attitude of dental trauma among dental students in Saudi Arabia. Eur J Dent 2015;9:518-22.

# **APPENDIX 1: QUESTIONNAIRE**

- Age:
- Gender:
- Country of graduation:
- Years of practice:

Place of practice: Private clinics, hospital-based clinic, public sector clinic

Have you encountered a patient who needed prophylactic antibiotics to prevent infective endocarditis? Yes no

Have you ever prescribed prophylactic antibiotics to prevent infective endocarditis to your patients? Yes no

Have you ever encountered a patient who developed infective endocarditis following dental treatment? Yes no

Which guidelines regarding prevention of infective endocarditis do you follow in your practice? AHA NICE others I do not know

If you encounter a patient who needs prophylactic antibiotics to prevent infective endocarditis

- 1. I prescribe the prophylactic antibiotics
- 2. Refer the patient to his physician for consultation
- 3. Refuse treating the patient and refer him to another colleague.

Which of the following patients would need prophylactic antibiotics to prevent infective endocarditis?

- 1. Patients with prosthetic valve
- 2. Patients with stents
- 3. Patients with history of open heart surgery
- 4. Patients with ischemic heart disease
- 5. Patients with ventricular septal defect
- 6. Patients with atrial septal defects

- 7. Patients with rheumatic heart disease
- 8. Patients with cyanotic congenital heart disease
- 9. Patients with a cyanotic congenital heart disease
- 10. Any patient with history of cardiac disease
- 11. Patients with previous history of infective endocarditis
- 12. Cardiac bypass
- 13. Patients with pacemaker
- 14. Mitral valve prolapsed or regurgitation
- 15. Physiologic murmur
- 16. Cardiac transplantation
- 17. Do not know.

Which of the following procedures would require the administration of prophylactic antibiotics to prevent infective endocarditis?

- 1. Surgical tooth extraction
- 2. Periodontal surgery
- 3. Implants
- 4. Biopsy
- 5. Apicoectomy
- 6. Reimplantation of avulsed tooth
- 7. Scaling and polishing
- 8. Simple tooth extraction
- 9. Endodontic treatment
- 10. Abscess drainage
- 11. Dentoalveolar trauma
- 12. Placement of fixed orthodontic band
- 13. Simple and complex restorations
- 14. Tooth preparation and impression taking
- 15. Placement of rubber dam
- 16. Placement of matrix band and wedges
- 17. Intraligamentary anesthesia
- 18. Inferior dental nerve block
- 19. Do not know.

Which antibiotics do you prescribe for prevention of infective endocarditis before dental procedures?

- Type
- Dose
- Duration.