# Role of fetal surgery in spinal dysraphism

# A. Martina Messing-Jünger

Department of Neurosurgery, Asklepios Children's Hospital, Sankt Augustin, Germany

# A B S T R A C T

Open spinal dysraphism is a common and clinically challenging organo-genetic malformation. Due to the well-known multi-organ affection with significant implication on the lives of patients and their families, abortion after prenatal diagnosis became reality in most parts of the world. After publication of the Management of Myelomeningocele Study (MOMS) results fetal surgery seems to be a new option and a broad discussion arose regarding advantages and risks of in utero treatment of spina bifida. This paper tries to evaluate objectively the actual state of knowledge and experience. This review article gives a historical overview as well as the experimental and pathophysiological background of fetal surgery in open spinal dysraphism. Additionally clinical follow-up experience of foetoscopically treated patients are presented and discussed. After carefully outweighing all available information on fetal surgery for spina bifida, one has to conclude, in accordance with the MOMS investigators, that in utero surgery cannot be considered a standard option at present time. But there is clear evidence of the hypothesis that early closure of the spinal canal has a positive influence on spinal cord function and severity of Chiari malformation type II, has been proven. A persisting problem is the fetal risk of prematurity and the maternal risk of uterus damage. There is also evidence that due to technical restrictions, fetal closure of the spinal canal bears unsolved problems leading to a higher postnatal incidence of complication surgery. Finally, missing long-term results make a definite evaluation impossible so far. At the moment, fetal surgery in open spinal dysraphism is not a standard of care despite promising results regarding central nervous system protection due to early spinal canal closure. Many technical problems need to be solved in the future in order to make this option a safe and standard one.

Key words: Fetal surgery, myelomeningocele, open spinal dysraphism

# INTRODUCTION

Open spinal dysraphism is one of the most frequent congenital malformations with possible multi-organ manifestations. Therefore affected children develop significant disability and sometimes life-threatening complications. Lifelong medical support is needed. The fetal hit leading to spina bifida aperta happens during early organogenesis in neurulation phase 12 around gestational day (GD) 28, when the neural tube is formed from cranial down to the lumbar segments.

*In utero* ultrasound and magnetic resonance imaging studies have shown that in many cases of open neural tube defects lower limb development and movements are normal at early stages and tend to deteriorate toward late pregnancy.<sup>[1]</sup>

Access this article online		
Quick Response Code:	Website	
	www.ijns.in	
	DOI: 10.4103/2277-9167.110205	

This observation led to the hypothesis, that the amniotic environment secondarily damages the neural tissue and early closure could possibly prevent this. A second hypothesis was discussed related to the development of Chiari malformation type II, e.g., of hindbrain herniation, which most likely is a result of excessive cerebrospinal fluid (CSF) outflow from the spinal defect.

These observations lead to animal trials investigating the influence of *in utero* closure of experimental neural tube defects. Due to positive results in some of them, a multicenter trial to investigate the effect in human fetal surgery for open spinal dysraphism was conducted with some encouraging results.

After the publication of the Management of Myelomeningocele Study (MOMS) results, fetal surgery seems to be a novel option not only for treatment, but also in respect of long-term outcome of affected children. There is an ongoing discussion regarding advantages and risks of *in utero* treatment of spina bifida versus postnatal surgery, which becomes a crucial issue also in prenatal counseling. This paper tries to evaluate objectively the actual state of knowledge and experience.

Address for correspondence: Dr. A. Martina Messing-Jünger, Department of Neurosurgery, Asklepios Children's Hospital, Sankt Augustin, Germany. E-mail: m.messing@asklepios.com

# HISTORICAL BACKGROUND

During pregnancy follow up, the former diagnostic standard for spina bifida detection consisted of maternal serum screening for  $\alpha$ -fetoprotein around gestational week (GW) 16-18, followed by amniocentesis and determination of amnion fluid acetylcholinesterase. Nowadays, fetal anatomical ultrasound with high resolution and fetal magnetic resonance imaging with a detection rate of 98% are screening standards.

In the 1980s the first animal models for *in utero* repair of congenital defects were conducted by Adzick and Harrison.<sup>[2]</sup> In 1994 a fetal lamb model was introduced by Meuli and Adzick,<sup>[3,4]</sup> in which a myelomeningocele (MMC)-like defect was created a GD 75 (term 145-150 days). The defect was covered with a latissimus dorsi flap at GD 100. After cesarean section prior to term, near-normal neurological functions and a well-preserved anatomy and histology was found in study animals compared with controls showing human-like defects.

Additionally, it could be demonstrated that lumbar myelotomy leads to Chiari type II like hindbrain herniation in a sheep model. After fetal closure of the spinal defect, normal hindbrain anatomy was found.<sup>[5]</sup>

The first human fetal MMC closure was performed foetoscopically in 4 fetuses by Bruner.<sup>[6]</sup> He covered the dysraphic defect using a maternal flap. Two of the fetuses died and the series was stopped.

In 1998 the first open fetal surgery was performed at Children's Hospital of Philadelphia (CHOP) in GW 22 by Adzick.<sup>[7]</sup> Tulipan started his series of open fetal surgery later during pregnancy (GW 28-30) and found that all 4 fetuses were born prematurely with evidence of reversed Chiari type II. Two of the babies needed an early CSF shunt. In later series he found that the most striking result was the positive effect on hindbrain herniation.<sup>[8]</sup> It could be shown that surgery later than GW 25 has no benefit for neurological function and Chiari type II. Other groups had similar results.<sup>[9]</sup>

Based on these results and experience a randomized controlled multicenter study was initiated by the National Institute of Health (USA) in 2003. Participants of MOMS were the three centers with the largest surgical experience in fetal MMC repair: CHOP, University of California, San Francisco (UCSF) and Vanderbilt University (VU). Study duration was 8 years and costs were US \$22.5 million.

A total of 200 pregnancies/mothers were scheduled for

inclusion. The study was terminated after 183 included cases after reaching significance level. Initially 1,087 fetuses and mothers were screened. CHOP treated 77 mothers, UCSF 54 and VU 52 (http://www.nejm. org/doi/suppl/10.1056/NEJMoa1014379/suppl\_file/ nejmoa1014379\_protocol.pdf). In March 2011, study results were published and latest since then, a worldwide discussion is going on regarding the optimal treatment and prognosis for open spina bifida.

During MOMS, all other centers in North America discontinued comparable surgical activities. Parallelly open and foetoscopic surgery was performed in single centers around the globe, the largest series are reported from Poland<sup>[10]</sup> (open), Germany<sup>[11,12]</sup> (foetoscopic) and recently from Brazil,<sup>[13]</sup> but not under controlled study conditions.

Beside open spinal dysraphism a variety of other indications for *in utero* surgery exist and some of them are considered as standard treatment. Due to the essential sophisticated skills regarding anesthesia and surgery inside the amnion cavity and related problems in closing both amnion sac and uterus these operations remain reserved for highly specialized centers. The North American Fetal Therapy Network is a voluntary association of such medical centers with expertise in all kinds of fetal surgery in USA and Canada. In Europe major centers are located in Leuven (Belgium), Linz (Austria), Zürich (Switzerland) as well as Bonn and Hamburg (Germany).

# **RESULTS OF MOMS**

The rationale of MOMS was to reduce amnion fluid toxicity in order to minimize a second-hit damage to the neural tissue and to stop CSF-outflow out of the spinal defect in order to minimize hindbrain herniation as major manifestation of Chiari malformation type II.

Accordingly a positive effect on postnatal paresis, bladder and bowel dysfunction, and development of hydrocephalus, secondary tethered cord and possible Chiari-related complications was expected and thus investigated. The trial compared open prenatal MMC closure versus open postnatal reconstruction in 158 pregnant women including postnatal neurological and radiological tests of the infants 12 months after birth as well as mental and motor development tests after 30 months. Fetal closure was performed without surgical neurulation of the spinal cord, postnatal surgery included standard plastic reconstruction of placode and all tissue layers. The primary outcome criteria were shunt rate and death. In the fetal group both were 68% compared with 98% in the postnatal group (P<0.001). Tonsillar herniation has been found in 64% (fetal) compared with 96% (postnatal) [Table 1]. After 30 months a significantly improved mental and motor development was also found (P=0.007).

Fetal surgery resulted in all cases in preterm delivery, in average at GW 34. In 13% pregnancy was terminated before GW 30. The incidence of uterine dehiscence was 30% and both cases with fatal outcome occurred in the prenatal group. The causes were extreme prematurity at GW 23 and severe Chiari malformation type II. The authors found a higher incidence of surgical interventions due to early secondary tethered cord syndrome.<sup>[14]</sup>

The results need to be discussed and the authors themselves do not consider fetal surgery as standard of care yet, despite significant improvement of outcome parameters. So far, the long-term results in both groups are not available and bladder and bowel dysfunction were not study parameters. Looking at the reported shunt rate, a certain bias can be observed: Independent reviewers found that in the prenatal group an indication for hydrocephalus shunting was given in 65%, but only 40% received a shunt. In the postnatal group there was also a tendency to avoid a shunt, but not as pronounced as in prenatal patients (92% indicated vs. 82% shunted). The most important argument against considering fetal closure as a standard is the fact, that the presented data derive from three highly specialized centers and that expertise and technical background cannot be guaranteed in other centers.

#### Experience with foetoscopic MMC closure

Parallel to open fetal surgery endoscopic techniques were tested as well, mainly to reduce the uterine damage. Technical problems in respect of amniotic membrane and uterus closure and more important regarding the plastic covering of the dysraphic lesion

Table 1: Results of primary outcome criteriadeath/shunt and extent of tonsillar herniationafter fetal and postnatal surgery (management ofMyelomeningocele Study, Adzick et al. <sup>[14]</sup> )				
	Fetal	Postnatal		
Death/shunt %	68	98	P<0.001	
Death	2	0		
Shunt %	40	82		
Chiari II				
Tonsillar herniation %	64	96		
Mild	40	29		
Moderate	19	45		
Severe	6	22		

itself remained unsolved until recently. Therefore foetoscopy was considered less promising and not tested in a major study.

After the early experience with fetal surgery in North America and Europe, also with established foetoscopic procedures for other indications, a German pediatrician first developed a sheep model for foetoscopic MMC closure<sup>[15]</sup> and then started a series of individual treatment trials in human fetuses.<sup>[16,17]</sup> The fact that this ongoing series is not controlled, the true incidence of complications and the overall outcome remains obscure, e.g., there is no matched population of postnatally treated patients in a comparable medical and social environment. The endoscopic interventionalist himself indicates that he is offering endoscopic treatment to a very select group of patients.<sup>[11,12]</sup> In order to get more evidence regarding the clinical results of this technique, several independent colleagues, pediatric neurologists as well as neurosurgeons collected data from patients followed by them with those who were primarily treated foetoscopically. In these randomly collected patient groups the shunt incidence was as high as in regularly treated patients, although shunts were put in many cases after the sixth month. Severe courses with early need for Gardner decompression due to excessive tonsillar herniation were seen, and numerous patch complications and early detethering procedures in secondary tethered cord syndromes were also noticed. On the other side, good motor development with only moderate bladder dysfunction could be observed in good-risk patients, which means low and smaller lesions, with intact patch closure. In this foetoscopic series, at least in the beginning, fatal outcome and maternal problems, e.g., uterus and amniotic cavity complications were reported. All children were premature, some of them extremely.

There is one independent paper reporting upon patients from this series. Verbeek et al.<sup>[18]</sup> could demonstrate that segmental neurological function after foetoscopic repair was better preserved compared with matched pairs with standard postnatal treatment, some of them after natural birth. A total of 13 pairs were investigated regarding segmental neurological function and leg muscle ultrasound density (MUD), a technique developed by Verbeek for functional muscle evaluation, e.g., in small children with dysraphic disorders.<sup>[19,20]</sup> Additionally they determined the inter-individual difference in MUD (dMUD), which describes the difference between affected myotomes cranial and caudal to the MMC level. This parameter was interpreted as extent of neuronal damage. The results show a better neurological functioning in respect of preserved knee-jerk (P=0.006) and anal reflex (P<0.05) in the fetal group, the dMUD Table 2: Results of fetoscopic myelomeningocele closure performed at Department of Obstetrics, University of Bonn, Germany, between 2003 and 2009 (Verbeek *et al.*<sup>[18]</sup>)

n foetoscopic MMC interventions n fetal death		19
		3
n interruption by iatrogenic hemorrhages		3
Matched pairs analysis	Fetal	Postnatal
n	13	13
Mean age	14 months	
F/M ratio	1.6	
GW at birth (median)	32	39 ( <i>P</i> =0.001
Preserved knee-jerk reflex	$\uparrow$	↓ ( <i>P</i> =0.006
Preserved anal reflex	$\uparrow$	↓ ( <i>P</i> =0.032
dMUD (mean 24, 95% CI 15-33)	$\checkmark$	↑ ( <i>P</i> <0.05)
Complications*	$\uparrow$	$\checkmark$

MMC – Myelomeningocele; GW – Gestational week; dMUD – Difference in muscle ultrasound density. \*(chorioamnionitis, premature rupture of amniotic membranes, oligohydramnios, respiratory distress syndrome, in premature baby)

was smaller as well (P<0.05). An overview of the results is given in Table 2. Out of 19 pregnancies 3 fetal deaths and 3 interruptions due to iatrogenic hemorrhage resulted. The complication rate was high [Table 2].

## DISCUSSION

In general the standard treatment for open spinal dysraphism is postnatal plastic reconstruction after cesarean section, if possible. From neurosurgical point of view surgical neurulation and careful closure with sufficient CSF space around the placode is the best option for prevention of secondary tethering. Hydrocephalus shunting in due time is also recommended for optimal long-term mental and motor development. Beside Chiari-related circulation disorders, malresorptive hydrocephalus due to chemical CSF reactions caused by the open tube defects is also present in 80-90% of the patients.

Theoritically, fetal defect closure is able to reduce the negative impact of the amniotic fluid on neural tissue and CSF and also minimizes CSF-outflow with positive effect on hindbrain herniation formation. Due to the fact that the time window for fetal intervention lies between GW 22 and 25, some early damage of the neural structures cannot be prevented and potential fetal and maternal risks exist related to the early gestational state. Anesthesia as well as obstetric techniques must be very sophisticated and performed by a team with high expertise<sup>[21]</sup> in specialized centers. Even in good-risk patients treated under best clinical conditions fetal surgery for MMC inevitably results in significant prematurity and bears a 30% rate of uterus lesions. The *in utero* defect closure itself seems to be less appropriate both in open and in endoscopic surgery compared with postnatal treatment, as the increased rate of early interventions due to secondary tethering shows. Endoscopic coverage is technically more demanding and *in utero* and persisting postpartal CSF leakages frequent with significant complications and less successful prevention of hindbrain herniation have been reported.

No long-term outcome data exist so far as well as on bladder dysfunction. The fact that secondary tethering is an unsolved problem and hydrocephalus of later onset might be expected, secondary evaluation of the preliminary MOMS outcome data has to be awaited. Probably the best treatment for spina bifida will be prevention, e.g., folic acid supplementation and stem cell repair somewhere in the future.<sup>[22,23]</sup>

#### CONCLUSION

Taking into account all available evidence, fetal surgery in open spinal dysraphism cannot be considered as standard of care despite promising results regarding central nervous system protection due to early spinal canal closure and less severe hindbrain herniation. Many technical problems need to be solved in the future in order to make this option a safe and standard one.

The ethical conflicts related to prenatal counseling and different treatment modalities in open spina bifida potentially affecting two individuals, the mother and the (unborn) child, are complex and cannot be addressed here. But they should always be considered when decisions are to be made.

## REFERENCES

- Beuls E, Vanormelingen L, Van Aalst J, Vandersteen M, Adriaensen P, Cornips E, *et al.* The Arnold-Chiari type II malformation at midgestation. Pediatr Neurosurg 2003;39:149-58.
- Adzick NS, Harrison MR, Flake AW. Experimental studies on prenatal treatment of congenital anomalies. Br J Hosp Med 1985;34:154-9.
- Meuli-Simmen C, Meuli M, Hutchins GM, Harrison MR, Buncke HJ, Sullivan KM, et al. Fetal reconstructive surgery: Experimental use of the latissimus dorsi flap to correct myelomeningocele in utero. Plast Reconstr Surg 1995;96:1007-11.
- Meuli M, Meuli-Simmen C, Yingling CD, Hutchins GM, Hoffman KM, Harrison MR, et al. Creation of myelomeningocele in utero: A model of functional damage from spinal cord exposure in fetal sheep. J Pediatr Surg 1995;30:1028-32.
- Bouchard S, Davey MG, Rintoul NE, Walsh DS, Rorke LB, Adzick NS. Correction of hindbrain herniation and anatomy of the vermis after *in utero* repair of myelomeningocele in sheep. J Pediatr Surg 2003;38:451-8.
- Bruner JP, Richards WO, Tulipan NB, Arney TL. Endoscopic coverage of fetal myelomeningocele *in utero*. Am J Obstet Gynecol 1999;180:153-8.
- 7. Adzick NS. Fetal surgery for myelomeningocele: Trials and tribulations. Isabella Forshall Lecture. J Pediatr Surg 2012;47:273-81.

- Tulipan N, Bruner JP, Hernanz-Schulman M, Lowe LH, Walsh WF, Nickolaus D, et al. Effect of intrauterine myelomeningocele repair on central nervous system structure and function. Pediatr Neurosurg 1999;31:183-8.
- Danzer E, Finkel RS, Rintoul NE, Bebbington MW, Schwartz ES, Zarnow DM, et al. Reversal of hindbrain herniation after maternal-fetal surgery for myelomeningocele subsequently impacts on brain stem function. Neuropediatrics 2008;39:359-62.
- Zamłyńiski J, Olejek A, Bohosiewicz J, Bodzek P, Mańka G, Grettka K, et al. Perinatal results of intrauterine open fetal surgery of fetuses diagnosed with myelomeningocoele – The clinical report of ten cases. Ginekol Pol 2007;78:647-51.
- Kohl T, Tchatcheva K, Merz W, Wartenberg HC, Heep A, Müller A, et al. Percutaneous fetoscopic patch closure of human spina bifida aperta: Advances in fetal surgical techniques may obviate the need for early postnatal neurosurgical intervention. Surg Endosc 2009;23:890-5.
- 12. Kohl T. Minimally invasive fetoscopic interventions: An overview in 2010. Surg Endosc 2010;24:2056-67.
- Hisaba WJ, Cavalheiro S, Almodim CG, Borges CP, de Faria TC, Araujo Júnior E, *et al*. Intrauterine myelomeningocele repair postnatal results and follow-up at 3.5 years of age – Initial experience from a single reference service in Brazil. Childs Nerv Syst 2012;28:461-7.
- Adzick NS, Thom EA, Spong CY, Brock JW 3<sup>rd</sup>, Burrows PK, Johnson MP, et al. A randomized trial of prenatal versus postnatal repair of myelomeningocele. N Engl J Med 2011;364:993-1004.
- 15. Kohl T, Hartlage MG, Kiehitz D, Westphal M, Buller T, Achenbach S, *et al.* Percutaneous fetoscopic patch coverage of experimental lumbosacral full-thickness skin lesions in sheep. Surg Endosc 2003;17:1218-23.
- 16. Kohl T, Hering R, Heep A, Schaller C, Meyer B, Greive C, *et al.* Percutaneous fetoscopic patch coverage of spina bifida aperta in

the human–early clinical experience and potential. Fetal Diagn Ther 2006;21:185-93.

- Kohl T, Gembruch U. Current status and prospects of fetoscopic surgery for spina bifida in human fetuses. Response to Fichter *et al.*: Fetal spina bifida repair – Current trends and prospects of intrauterine neurosurgery (Fetal Diagn Ther 2008;23:271-286). Fetal Diagn Ther 2008;24:318-20.
- Verbeek RJ, Heep A, Maurits NM, Cremer R, Hoving EW, Brouwer OF, et al. Fetal endoscopic myelomeningocele closure preserves segmental neurological function. Dev Med Child Neurol 2012;54:15-22.
- Verbeek RJ, van der Hoeven JH, Sollie KM, Maurits NM, Bos AF, den Dunnen WF, *et al*. Muscle ultrasound density in human fetuses with spina bifida aperta. Early Hum Dev 2009;85:519-23.
- Verbeek RJ, van der Hoeven JH, Maurits NM, Brouwer OF, Hoving EW, Sival DA. In spina bifida aperta, muscle ultrasound can quantify the "second hit of damage". Childs Nerv Syst 2013; 29:469-74.
- 21. Van de Velde M, De Buck F. Fetal and maternal analgesia/anesthesia for fetal procedures. Fetal Diagn Ther 2012;31:201-9.
- 22. Dhaulakhandi DB, Rohilla S, Rattan KN. Neural tube defects: Review of experimental evidence on stem cell therapy and newer treatment options. Fetal Diagn Ther 2010;28:72-8.
- Li H, Gao F, Ma L, Jiang J, Miao J, Jiang M, *et al*. Therapeutic potential of *in utero* mesenchymal stem cell (MSCs) transplantation in rat foetuses with spina bifida aperta. J Cell Mol Med 2012;16:1606-17.

**How to cite this article:** Messing-Jünger AM. Role of fetal surgery in spinal dysraphism. Indian J Neurosurg 2013;2:4-8.

Source of Support: Nil, Conflict of Interest: None declared.

#### "Quick Response Code" link for full text articles

The journal issue has a unique new feature for reaching to the journal's website without typing a single letter. Each article on its first page has a "Quick Response Code". Using any mobile or other hand-held device with camera and GPRS/other internet source, one can reach to the full text of that particular article on the journal's website. Start a QR-code reading software (see list of free applications from http://tinyurl.com/ yzlh2tc) and point the camera to the QR-code printed in the journal. It will automatically take you to the HTML full text of that article. One can also use a desktop or laptop with web camera for similar functionality. See http://tinyurl.com/2bw7fn3 or http://tinyurl.com/3ysr3me for the free applications.