

Does nature of diet consumed has an association with the prevalence of dental caries?

Sir,

Polarization of caries is occurring on a worldwide basis, where the prevalence of caries is declining in developed countries and increasing in less-developed countries; also, this is epidemic in countries with emerging economies.^[1] Diet has been associated with the prevalence of dental caries for centuries. In the field of research about caries etiology, diet has probably received more attention than any other subjects.^[2]

Literature has shown that nature of diet consumed, whether vegetarian or mixed, shows association with prevalence of dental caries. To reveal this association, a study was conducted among 6-12-year-old school children in Shimla city, Himachal Pradesh, India. A cross-sectional study was conducted on 1,200 school children aged 6-12 years from 11 government elementary schools, which were selected randomly from all parts of the city. Data was collected through clinical examination of the children and questions were asked related to the nature of the consumed diet. Data was analyzed using SPSS (version 13), and Chi-square test was used for comparison.

Results of the study revealed the association. Children following a vegetarian diet had a higher prevalence of dental caries than those following a mixed diet (79.4% vs. 74.2%); this difference was highly statistically significant ($\chi^2=19.71$, $P<0.01$). Similar results were reported by Khan *et al.*^[3] and Bhardwaj *et al.*^[4] Individuals who consume mixed diet rich in protein develop less amount of acid in their mouth and are relatively protected from dental caries. This may be due to putrefaction of the protein diet rather than fermentation of carbohydrates, where alkalinity replaces acidity and no decalcification is usually observed.^[3]

Thus, the study revealed that vegetarian/mixed diet has an impact on the prevalence of dental caries.

However, to prove this point, further studies need to be conducted among different populations and age groups.

**Vinay Kumar Bhardwaj, Nishant Negi¹,
Shailee Fotedar, Jai Ram Kaundal²**

Department of Public Health Dentistry, ¹Orthodontics and Dentofacial Orthopaedics, H. P. Govt. Dental College and Hospital, Shimla, Himachal Pradesh, India

Address for correspondence:

Dr. Vinay Kumar Bhardwaj,
Department of Public Health Dentistry, H. P. Govt. Dental College and Hospital,
Shimla - 171 001, Himachal Pradesh, India.
E-mail: dr.viney@gmail.com

REFERENCES

1. Sudha P, Bhasin S, Anegundi RT. Prevalence of dental caries among 5-13 year old children of Mangalore city. *J Indian Soc Pedod Prev Dent* 2005;23:74-9.
2. Gustafson BE, Quensel CE, Lauke LS. The Vipeholm dental caries study; the effect of different levels of carbohydrate intake on caries activity in 436 individuals observed for five years. *Acta Odontol Scand* 1954;11:232-364.
3. Khan AA, Jain SK, Shrivastav A. Prevalence of dental caries among the population of Gwalior (India) in relation of different associated factors. *Eur J Dent* 2008;2:81-5.
4. Bhardwaj VK, Sharma A. An epidemiological study of dental caries prevalence in mixed dentition among school children and its association with different variables in Shimla city. *J Res-The Bede Athenaeum* 2011;2:1-5.

Access this article online

Quick Response Code:



Website:
www.ejgd.org

DOI:
10.4103/2278-9626.115990