Original Article

Evaluation of ⁶⁸Ga-DOTA-Ubiquicidin (29–41) for imaging *Staphylococcus aureus* (Staph A) infection and turpentine-induced inflammation in a preclinical setting

ABSTRACT

Synthetic antimicrobial peptide fragment, ^{99m}Tc-Ubiquicidin 29–41, is shown to be sensitive and also specific for imaging bacterial infections. We undertook this study to explore the advantage of using a positron emission agent, ⁶⁸Ga-DOTA-Ubiquicidin 29–41 (⁶⁸Ga-DOTA-UBI), for detecting *Staph-A* infection in an animal model, and also evaluated its ability to distinguish a turpentine-induced sterile inflammation in an animal model. Pure Ga-68 was freshly eluted from a ⁶⁸Ge/⁶⁸Ga generator (IGG-100). DOTA-UBI (50 µg) was radiolabeled with pure Ga-68 (500MBq) by incubating the reaction mixture at pH 4.5 for 10 min, 95 C. Rats were infected with *Staph-A* at the hind leg joint of rats to form bacterial abscess. Sterile inflammation was induced in the right thigh muscle by injecting 200 µl of 100% turpentine oil. Rats were injected intravenously with 10–15 MBq of tracer, and images were acquired at different time intervals with Siemens (Biograph mCT) positron emission tomography computed tomography scanner. The early images at 6 min postinjection clearly indicated mild uptake of the agent corresponding to the infection site, which increased dramatically at 20, 30, and 60 min postinjection. The target to background ratio (T/B) increased significantly over the same time period of study (1.6, 4.2, and 6.1, respectively). There was a mild uptake of ⁶⁸Ga-DOTA-UBI at the site corresponding to sterile inflammation at 6 min postinjection, which was rapidly washed off as seen at 25 and 45 min images. The images indicated fast clearance of the agent from liver and soft tissues within 6 min. Control rats showed similar biodistribution of activity. The mild uptake of ⁶⁸Ga-DOTA-UBI at the corresponding *Staph-A* infection lesion and very fast kinetics of clearance from the blood pool and soft tissues suggested a very high clinical potential for this agent. The absence of uptake of the agent at sterile inflammation site suggests that the agent may be useful in distinguishing infection from inflammation.

Keywords: ⁶⁸Ga-DOTA-Ubiquicidine (29–41), ⁶⁸Ga-positron emission tomography, infection imaging, inflammation imaging, *Staphylococcus aureus*

INTRODUCTION

Developing an ideal agent for imaging infection has been challenging over the past four decades. A number of single photon-emission computerized tomography (SPECT) agents have emerged in the past but with varying degree of success and limitations.^[1,2] ^{99m}Tc-labeled leukocytes have shown promise, but the procedure was complicated and required special skills. ⁶⁷Ga-citrate was used for the past four decades for imaging infection, but it required up to 72 h waiting time before the images could be obtained. Gold standard positron emission tomography (PET) agent ¹⁸F-fluorodeoxyglucose played a significant role in infection imaging due to its very high sensitivity, but it

Access this article online	
	Quick Response Code
Website: www.wjnm.org	
DOI: 10.4103/wjnm.WJNM_103_20	

DILIP KUMAR BODDETI^{1,2,3}, VIJAY KUMAR^{1,2,4}

¹Department of Nuclear Medicine and PET, Westmead Hospital, ²Department of Nuclear Medicine, The Children's Hospital at Westmead, Westmead, ³Westmead Clinical School, University of Sydney, ⁴Discipline of Child and Adolescent Health, Sydney Medical School, University of Sydney, Sydney NSW, Australia

Address for correspondence: Prof. Vijay Kumar, Department of Nuclear Medicine and PET, Westmead Hospital, Westmead 2145, NSW, Australia. E-mail: vijay.kumar@health.nsw.gov.au

Submitted: 21-Jul-2020, Revised: 09-Sep-2020, Accepted: 30-Sep-2020, Published: 10-Apr-2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Boddeti DK, Kumar V. Evaluation of ⁶⁸Ga-DOTA-Ubiquicidin (29–41) for imaging *Staphylococcus aureus* (Staph A) infection and turpentine-induced inflammation in a preclinical setting. World J Nucl Med 2021;20:266-72.

© 2021 World Journal of Nuclear Medicine | Published by Wolters Kluwer - Medknow

lacked specificity. ⁶⁸Ga tracer is now readily available from a commercial ⁶⁸Ge/⁶⁸Ga generator, which is cost-effective and we explored the ability of ⁶⁸Ga-apo-transferrin,^[3] ⁶⁸Ga-Citrate-PET for the diagnostic imaging of *Staph-A* infection in rats and for intra-abdominal infection in patients^[4] and ⁶⁸Ga-DOTA-Ubiquicidin 29–41 (⁶⁸Ga-DOTA-UBI) to detect *Staph-A* infection lesions in an animal model.^[5]

Antimicrobial peptides are low-molecular-weight proteins, which have the broad spectrum of antimicrobial activity against bacteria.^[5] Ubiquicidin 29-41 (UBI 29-41) is a 12 amino acid peptide (Thr-Gly-Arg-Alu-Lys-Arg-Met-Gln-Tyr-Asn-Arg-Arg) is a synthetic cationic antimicrobial peptide fragment with a weight of 1,693 Da. k^{[6,7] 99m}Tc-UBI 29–41 targets specifically bacterial and fungal cell wall but fails to target mammalian cells or cancer cells.^[8-12] Due to such excellent properties, several studies were undertaken to show the usefulness of UBI 29-41 for imaging infection. 99mTc-UBI was shown to be a useful agent in the diagnosis of orthopedic infection.^[13,14] Zijlstra et al.^[15] have demonstrated the utility of ¹⁸F-Fluorine labelled 4-fluoro ¹⁸F-UBI 29-41 for imaging infection. Recent studies by Ebenhan et al.^[16] shown the evaluation of⁶⁸Ga NOTA (1,4,7-triazacyclononane-triacetic acid)-UBI 29-41 PET for imaging infection. In the present study, we describe ⁶⁸Ga labeling of antibacterial peptide UBI 29-41, using DOTA as the bifunctional chelator (BFC), which has high thermodynamic and kinetic stability, for detecting Staph-A infection in an animal model. We have used DOTA as the BFC for the first time to study Ga-68 labeling with UBI.^[17] Subsequently, DOTA has been used as BFC for Ga-UBI by two other groups.^[18,19]

MATERIALS AND METHODS

An automated radiosynthesizer was used for the production of ⁶⁸Ga-DOTA-UBI. All chemicals were of pharmaceutical grade and of high purity were obtained from Merck and Sigma-Aldrich (Germany). Water for trace analysis (Trace SELECT) was purchased from Honeywell (Germany). DOTA-UBI was purchased from Auspep (Australia) with purity >95%. DOTA-UBI was supplied as freeze-dried powder in vials containing 100 μ g aliquots which were stored at-80°C (+5°C) freezer. DOTA-UBI was reconstituted 1.0 mg/mL with water prior to use. The 68Ge/68Ga generator (IGG-100) and automated radiosynthesizer (Modular-Lab PharmTracer) were obtained from Eckert and Ziegler, USA. Alumina backed chromatography plate Silica gel 60 F_{254} was supplied by Merck Millipore, USA. A clinical isolate of Staph A was obtained from the Department of Clinical Microbiology, Institute of Clinical Pathology and Medical Research Westmead Hospital, Westmead. A laminar flow cabinet was used during all microbiological work to prevent contamination.

Preparation of ⁶⁸Ga-DOTA-Ubiquicidin 29–41 (29–41)

Briefly, the 68 Ge/ 68 Ga generator was eluted with 7.0 mL of hydrochloric acid (0.1M) and the Ga-68 was bound onto a strata resin strong cation exchange and subsequently eluted with 5.0 M NaCl containing 5.5M HCl, which was then reacted with 50 µg of DOTA-UBI in 2.0 mL of water (TraceSELECT) and 400 µl of sodium acetate buffer pH 4.5 and by heating at 95°C for 10 min. After labeling, the reaction vial was cooled and passed through C-18 cartridge to remove any free Ga-68.

Determination of radiochemical purity

The radiochemical purity (RCP) was analyzed using radio-thin layer chromatography (TLC) and high-performance liquid chromatography (HPLC) methods. The percentage of RCP of ⁶⁸Ga-DOTA-UBI was determined by alumina backed chromatography plate and sodium citrate (0.1 M), pH 5.5 as the solvent. Bioscan Radio-TLC Imaging instrument was used to scan the plate. The %RCP is also estimated using a Shimadzu Radio-HPLC gradient method, Kinetex C18 column (Phenomenex, Australia) (3.0 micron × 150 mm), with (A) 0.1% TFA/H₂O and (B) Acetonitrile as solvents with a flow rate of 0.6 ml/min.

Stability studies

The stability of ⁶⁸Ga-DOTA-UBI was evaluated at four different postlabeling time intervals at 0, 30, 60, and 3 h. Human serum stability study was performed by mixing 0.3 ml of ⁶⁸Ga-DOTA-UBI solution with 1.5 ml of human serum (five times volume) and incubating at 37°C for 1 h. RCP was estimated at the end of 1 h incubation period.

In vitro binding of ⁶⁸Ga-DOTA-Ubiquicidin 29–41 with *Staph-A*

Binding of ⁶⁸Ga-DOTA-UBI to *Staph A* was assessed at different time intervals for up to 2 h, by incubating the labelled agent with *Staph A* at 37°C as follows. Briefly, in a sterile 10 mL reaction vial, 2.0 mL of ⁶⁸Ga-DOTA-UBI (25MBq activity) containing 25 µg/mL DOTA-UBI) was incubated with *Staph A* (4 × 10⁷ CFU) in 5.0 mL saline. 1.0 mL aliquots were taken from the reaction vial at different time intervals (5, 30, 60, 90, and 120 min). The aliquots were filtered by using Millex-GV Syringe Filter Unit, 0.22 µm, PVDF, 33 mm (gamma sterilized) followed by 3 mL saline wash. The filter was pretreated with 2 mL saline before use. Activities associated with the filter and the filtrate fractions were measured by using a dose calibrator, and the values were recorded. The percentage binding of ⁶⁸Ga-DOTA-UBI was calculated using the activity associated with the (filter unit × 100)/total activity (filter unit + filtrate).

The activities at different time intervals were decay corrected. A control group was set up containing radiolabeled peptide in saline (without *Staph A*) to determine nonspecific peptide binding.

Inducing infection and sterile inflammation in animal model

The experimental protocol was approved by the Western Sydney Local Health District Animal Ethics Committee (WSLHD AEC Protocol No: 9010.08.12), Westmead Hospital, Westmead NSW 2145. Male Wister rats (6–7) weeks old weighing 250–300 g were anesthetized by intra-peritoneal injection of the solution containing ketamine (100 mg/ml)/ xylazine (20 mg/ml) (2:1 v/v). Animals were randomly assigned into two groups. Group 1 (n = 12)-Infection was induced in the right thigh muscle by intramuscular injection of 5 × 10⁵ CFU Staph A in 0.1 ml volume of sterile saline for injection. Three to four days after inoculation, an abscess formed. Group 2 (n = 6) sterile inflammation was induced in the right thigh muscle by intramuscular injection with 200 µl of 100% turpentine oil (Sigma Aldrich, Australia). Turpentine oil caused visible redness and swelling within 2–3 h after injection.

Dosage and animal imaging

Anesthetised rats in both the groups were injected intravenously in the tail vain with 10–15 MBq/0.1 mL saline of ⁶⁸Ga-DOTA-UBI. The images were acquired at three different postinjection time points at 6, 30, and 60 min for Staph A infection induced rats and 6, 25, and 45 min intervals for sterile inflammation rats. Images were acquired for 5 min each with a matrix size of 200, reconstruction method: True X + TOF (Ultra HD-PET), 21 subsets using PET/ computed tomography scanner (Siemens Biograph mCT). Standardized uptake value (SUV) max was calculated over different organs in rats by drawing the regions of interest by a computer-generated program, which is used routinely in patient's studies at our department.

RESULTS

Quality control

The RCP of ⁶⁸Ga-DOTA-UBI was >99%. ⁶⁸Ga-DOTA-UBI stayed at the origin [Figure 1a], while free Ga-68 moved with the solvent front [Figure 1b]. The RCP of ⁶⁸Ga-DOTA-UBI was > 98% at 0, 30, 60 min, and 3 h after radiolabeling without any significant changes. The stability was not tested beyond 3 h, as the $T_{1/2}$ of ⁶⁸Ga is only 68 min. The retention time by HPLC for free Ga-68 was 2 min and for the labelled product 7.2 min [Figure 1c]. The radiolabeling yield was >99%. ⁶⁸Ga-DOTA-UBI complex was stable in human serum as shown by RCP >99% when studied for up to 1 h at 37°C.

Biodistribution of ⁶⁸Ga-DOTA-Ubiquicidin 29–41 in normal rats

Biodistribution of ⁶⁸Ga-DOTA-UBI was shown in normal rats at 6, 30, and 60 min postinjection period [Figure 2]. The images clearly showed that there was significant uptake of the agent



Figure 1: Instant thin-layer chromatography of ⁶⁸Ga-DOTA-Ubiquicidin 29– 41 (a) and free Ga-68 (b) Using Sodium citrate (0.1M, pH 5.5) as the solvent. Free 68Ga moved with the solvent front to the top (Rf = 1.0), while labeled product stayed at the origin (Rf = 0–0.4). (c) Gradient high-performance liquid chromatogram with 0.1% TFA/H2O and acetonitrile as solvents showed the retention time of 7.2 min for⁶⁸Ga-DOTA-Ubiquicidin 29–41

in the kidneys and the bladder. The activities associated with the other soft tissues were very low and were cleared rapidly as shown at 6 min postinjection image. There was very little background activity in the soft tissues at 30 and 60 min postinjection images.

Estimation of standardized uptake values at different organs in a normal rat

Using a normal rat, uptake of the agent at various organs was estimated at different time intervals using a computer program, and the values were expressed as SUV [Figure 3]. The activities associated with the heart, muscle, bone, and liver were shown to be very low over the time period studied (SUV < 3). On the other hand, the activities associated with the kidneys have increased to SUV = 15 at 30 min period, and then began to wash off. There was significant activity associated with the bladder up to 50 min postinjection (SUV = 30) which then declined, commensurate with the kidney activity as expected.

In vitro binding studies

Results of *in vitro* binding studies clearly showed that ⁶⁸Ga-DOTA-UBI was able to bind avidly with *Staph-A* up to

50% of the injected dose (ID) within 30 min. There was a mild decrease with time, but the percentage binding was retained at 48% of ID even at 120 min of incubation [Figure 4].

⁶⁸Ga-DOTA-Ubiquicidin 29–41 for imaging *Staph-A* infection in an animal model

The results clearly showed that there was a mild uptake of ⁶⁸Ga-DOTA-UBI activity at the area corresponding to



into healthy normal rats, and the images were acquired at 6, 30, and 60 min postinjection. The images clearly show an avid accumulation of the agent by the kidneys and bladder. There is a mild cardiac blood pool activity at but all the other organs show very little activity even at 6 min postinjection



Figure 4: *In vitro* binding of 68 Ga-DOTA-Ubiquicidin 29–41 with Staph A was performed as described in the method section. The percentage binding of the tracer by Staph A was plotted against time as shown

bacterial lesion within 6 min postinjection in rats [Figure 5]. The intensity of the uptake has increased significantly at 30 and 60 min postinjection period. Very high uptake of the agent in the kidneys and bladder was comparable to the biodistribution of the agent in normal rats. The activity associated with other soft tissues was very low indicating fast clearance of the agent from soft tissues. This is consistent with rapid accumulation activity in the kidneys and bladder.



Figure 3: Standardized uptake value max was calculated at each time point for different organs in healthy rats, and the values were plotted in the graph as shown



Figure 5: ⁶⁸Ga-DOTA-Ubiquicidin 29–41 (10 MBq/0.1mL saline) was injected into a Staph A infected rat, and images were acquired at different time intervals (6, 30, and 60 min postinjection). Infection lesions are shown by the arrows. Mild uptake of the agent was seen at the infected site within 6 min of post injection, which becomes very intense at 30 and 60 min postinjection

⁶⁸Ga-DOTA-Ubiquicidin 29–41 for imaging sterile inflammation in an animal model

There was a mild uptake of ⁶⁸Ga-DOTA-UBI seen at the site corresponding to sterile inflammation at 6 min post injection, which was rapidly washed off as seen at 25 and 45 min images [Figure 6]. The uptake of the agent in the kidneys and bladder was comparable to biodistribution of the agent in normal rats including fast clearance of the agent from other soft tissues. A measurable uptake of the agent was also observed in the liver at 6 min post injection, which decreased significantly over 45 min time.

DISCUSSION

In recent years, several preclinical studies were reported using ^{99m}Tc-UBI 29–41 to study bacterial and fungal infection in animal models.^[20] Clinical trials report by Akhtar *et al.*^[21] with ^{99m}Tc-UBI 29–41 also added further support that ^{99m}Tc-UBI is a highly sensitive and specific agent for detecting infective foci in bone and soft tissues of humans. However, its clinical utility is compromised due to poor image quality of ^{99m}Tc, which is intrinsic to SPECT agents. The present study was conducted to study the utility of ⁶⁸Ga-DOTA-UBI for bacterial infection imaging, instead of ^{99m}Tc, to take the advantage of high resolution and sensitivity associated with ⁶⁸Ga PET tracer. We have used DOTA as the BFC (bifunctional chelator) to label the UBI peptide fragment with ⁶⁸Ga, as it is believed to have higher biological, thermodynamic, and kinetic stability *in vivo*.

The other purpose of our study was to investigate if this agent was capable of differentiating sterile inflammation from bacterial infection. Studies by Ebenhan *et al.*^[22,23] described



Figure 6: ⁶⁸Ga-DOTA-Ubiquicidin 29–41 (10 MBq/0.1 mL saline) was injected into a rat having turpentine-induced inflammation, and images were acquired at different time intervals (6, 25, and 45 min postinjection). Inflammation areas are shown by the arrows. A very mild uptake of the agent was seen at the site at 6 min of postinjection, which disappears at 25 and 45 min postinjection

⁶⁸Ga-labelled-UBI with NOTA as the BFC, and they have demonstrated its utility in imaging infection and its ability to differentiate infection from inflammation. Our results showed that ⁶⁸Ga-DOTA-UBI could be prepared by a simple method with high yield (>95%) and high RCP (>99%) within 10 min. In our experimental design, an optimal pH of 4.5 was used for radiolabeling of the peptide with ⁸Ga, which is comparable to reports in the literature.

The overall labeling efficiency of ⁶⁸Ga-DOTA-UBI was >99%, and therefore, no further C-18 purification was required. Using sodium acetate (2.5 M) as buffer, NOTA-conjugated peptides were labeled at 50°C with the yields between 40 and 50% labeled product; however, optimal labeling efficiencies (>99%) were achieved at 85°C.^[16] In our studies, the stability of the purified, labeled compound was positively tested, without any significant change of RCP for up to 6 h, to warrant its prospective use in preclinical studies. The agent was stable in human serum when studied for up to 1 h at 37°C.

Our *in vitro* results clearly indicated very high binding, which is nearly 50% of the added activity of ⁶⁸Ga-DOTA-UBI to bacteria, which is consistent with reports in the literature as >45% binding of (^{nat}Ga) NOTA-UBI 29–41 (Lys [Abz]) to the *Staph A* cells was reported by Ebenhan *et al.*^[23] and 34.6 \pm 3.0%^[25] and 40%–50%^[25] binding found for ^{99m}Tc-UBI 29–41. Welling *et al.*^[24,21] studied a ^{99m}Tc-UBI 29–41 fragment with a scrambled amino acid sequence and reported significantly decreased bacterial binding, thus supporting our *in vitro* findings. Available literature on the *in vitro* studies suggested that binding of ⁶⁸Ga-DOTA-UBI to bacteria could be the result of its high thermodynamic stability, selectivity, and stereospecificity.^[26]

Our *in vivo* results clearly indicated an avid uptake of the agent at the infection lesions within 6 min postinjection, which is faster than any other studies reported in the literature. The avidity of uptake increased dramatically at 30 and 60 min postinjection. Bio-kinetics studies by Meléndez-Alafort *et al.*^[27] in pediatric patients with bone infection explained that the short uptake time of UBI 29–41 at sites of infection could be due to the antimicrobial peptide UBI 29–41 interacts electrostatically with the membrane lipids of the bacteria. After entering the cell, the radiopharmaceutical could be bound to a cytoplasmatic specific site on a target bacterial protein causing a fast cell death with the subsequent bacterial removal.

Our studies indicated fast clearance of the agent from liver and soft tissues within 6 min post-injection and the delayed images clearly showed low activity in the blood pool and soft-tissues. A very high activity was associated with the kidneys and urinary bladder, as they are the excretory mechanism for the tracer. Our findings are consistent with recent human dosimetry measures^[28,29] in four healthy volunteers (2 women and 2 men) using 68Ga-DOTA-UBI, which showed that the tracer was rapidly cleared from the body by urinary excretion (kidney and bladder). They also suggested that the dose to the urinary bladder wall can be reduced by frequent bladder voiding. Control rats showed similar biodistribution of activity. Sterile inflammation induced rats showed a significant uptake of ⁶⁸Ga-DOTA-UBI in the liver at 6 min post injection, which may be due to high injected activity. Liver activity was rapidly washed off as seen at 25 and 45 min images. The uptake of the agent in the kidneys and bladder was comparable to the biodistribution of the agent in normal rats.

CONCLUSIONS

⁶⁸Ga-DOTA-UBI can be prepared using an automated radio synthesizer within 15 min. *In vitro* studies showed that the tracer bound the bacteria, *Staph-A*, with very high affinity. The *in vivo* studies showed mild uptake of ⁶⁸Ga-DOTA-UBI at *Staph-A* infection lesion within 6 min, which increased significantly at 30 and 60 min post-injection periods. Clearance from the blood pool and soft tissues showed very fast kinetics. ⁶⁸Ga-DOTA-UBI was not accumulating in sites corresponding to sterile inflammation. Therefore, ⁶⁸Ga-DOTA-UBI PET has a very high clinical potential for imaging infection and probably differentiate infection from inflammation.

Acknowledgments

The authors acknowledge Thomas Olma for providing *Staphylococcus aureus* and assisting in infecting the rats and Scott Evans for assisting in acquiring PET-CT images.

Financial support and sponsorship Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Hnatowich DJ. A review of radiopharmaceutical development with short-lived generator-produced radionuclides other than ^{99m}Tc. Int J Appl Radiat Isot 1977;28:169-81.
- Green MA, Welch MJ. Gallium radiopharmaceutical chemistry. Int J Rad Appl Instrum B 1989;16:435-48.
- Kumar V, Boddeti DK, Evans SG, Roesch F, Howman-Giles R. Potential use of ⁶⁸Ga-apo-transferrin as a PET imaging agent for detecting *Staphylococcus aureus* infection. Nucl Med Biol 2011;38:393-8.
- Kumar V, Boddeti DK, Evans SG, Angelides S. ⁶⁸Ga-Citrate-PET for diagnostic imaging of infection in rats and for intra-abdominal infection

in a patient. Curr Radiopharm 2012;5:71-5.

- Martin E, Ganz T, Lehrer RI. Defensins and other endogenous peptide antibiotics of vertebrates. J Leukoc Biol 1995;58:128-36.
- Epand RM, Vogel HJ. Diversity of antimicrobial peptides and their mechanisms of action. Biochim Biophys Acta 1999;1462:11-28.
- Lupetti A, Nibbering PH, Welling MM, Pauwels EK. Radiopharmaceuticals: New antimicrobial agents. Trends Biotechnol 2003;21:70-3.
- Ebenhan T, Gheysens O, Kruger HG, Zeevaart JR, Sathekge MM. Antimicrobial peptides: Their role as infection-selective tracers for molecular imaging. Biomed Res Int 2014;2014:867381.
- Akhtar MS, Iqbal J, Khan MA, Irfanullah J, Jehangir M, Khan B, *et al.* ^{99m}Tc-labeled antimicrobial peptide ubiquicidin (29-41) accumulates less in Escherichia coli infection than in Staphlococcus aureus infection. J Nucl Med 2004;45:849-56.
- Lupetti A, Welling MM, Pauwels EK, Nibbering PH. Radiolabelled antimicrobial peptides for infection detection. Lancet Infect Dis 2003;3:223-9.
- Vallejo E, Martinez I, Tejero A, Hernandez S, Jimenez L, Bialostozky D, et al. Clinical utility of ^{99m}Tc-labeled ubiquicidin 29-41 antimicrobial peptide for the scintigraphic detection of mediastinitis after cardiac surgery. Arch Med Res 2008;39:768-74.
- Welling MM, Mongera S, Lupetti A, Balter HS, Bonetto V, Mazzi U, et al. Radiochemical and biological characteristics of 99mTc-UBI 29-41 for imaging of bacterial infections. Nucl Med Biol 2002;29:413-22.
- Saeed S, Zafar J, Khan B, Akhtar A, Qurieshi S, Fatima S, *et al.* Utility of ⁹⁹mTc-labelled antimicrobial peptide ubiquicidin (29-41) in the diagnosis of diabetic foot infection. Eur J Nucl Med Mol Imaging 2013;40:737-43.
- Nazari B, Azizmohammadi Z, Rajaei M, Karami M, Javadi H, Assadi M, et al. Role of 99mTc-ubiquicidin 29-41 scintigraphy to monitor antibiotic therapy in patients with orthopedic infection: A preliminary study. Nucl Med Commun 2011;32:745-51.
- Zijlstra S, Gunawan J, Freytag C, Burchert W. Synthesis and evaluation of fluorine-18 labelled compounds for imaging of bacterial infections with PET. Appl Radiat Isot 2006;64:802-7.
- Ebenhan T, Zeevaart JR, Venter JD, Govender T, Kruger GH, Jarvis NV, et al. Preclinical evaluation of ⁶⁸Ga-labeled 1,4,7-triazacyclononane-1,4,7-triacetic acid-ubiquicidin as a radioligand for PET infection imaging. J Nucl Med 2014;55:308-14.
- Boddeti D, Evans S, Kumar V. Potential of ⁶⁸Ga-DOTA-UBI to detect Staph-A infection lesions in an animal model. J Nucl Med May 2014;55:383.
- Sasikumar A, Joy A, Nanabala R, Pillai MR, T A H. ⁶⁸Ga-DOTA Ubiquicidin PET/CT in an infected implant. Clin Nucl Med 2017;42:e115-6.
- Sriwiang W, Rangsawai W, Pumkhem S. ⁶⁸Ga-labeled Ubiquicidin for monitoring of mouse infected with *Staphylococcus aureus*. Int J Phys Conf Ser 2019 1285;1:012028.
- Beiki D, Yousefi G, Fallahi B, Tahmasebi MN, Gholamrezanezhad A, Fard-Esfahani A, *et al.* (99m) Tc-Ubiquicidin [29-41], a promising radiopharmaceutical to differentiate orthopedic implant infections from sterile inflammation. Iran J Pharm Res 2013;12:347-53.
- Akhtar MS, Qaisar A, Irfanullah J, Iqbal J, Khan B, Jehangir M, et al. Antimicrobial peptide ^{99m}Tc-ubiquicidin 29-41 as human infection-imaging agent: Clinical trial. J Nucl Med 2005;46:567-73.
- Ebenhan T, Govender T, Kruger G, Pulker T, Zeevaart JR, Sathekge M. Synthesis of ⁶⁸Ga-NOTA-UBI30-41 and *in vivo* biodistribution in vervet monkeys towards potential PET/CT imaging of infection. J Nucl Med 2012;53:1520.
- 23. Ebenhan T, Chadwick N, Sathekge MM, Govender P, Govender T, Kruger HG, et al. Peptide synthesis, characterization and ⁶⁸Ga-radiolabeling of NOTA-conjugated ubiquicidin fragments for prospective infection imaging with PET/CT. Nucl Med Biol 2014;41:390-400.

- Welling MM, Paulusma-Annema A, Balter HS, Pauwels EK, Nibbering PH. Technetium-99m labelled antimicrobial peptides discriminate between bacterial infections and sterile inflammations. Eur J Nucl Med 2000;27:292-301.
- Ferro-Flores G, Arteaga de Murphy C, Pedraza-López M, Meléndez-Alafort L, Zhang YM, Rusckowski M, *et al. In vitro* and *in vivo* assessment of ^{99m}Tc-UBI specificity for bacteria. Nucl Med Biol 2003;30:597-603.
- Nibbering PH, Welling MM, Paulusma-Annema A, Brouwer CP, Lupetti A, Pauwels EK. 99mTc-Labeled UBI 29-41 peptide for monitoring the efficacy of antibacterial agents in mice infected with *Staphylococcus aureus*. J Nucl Med 2004;45:321-6.
- Meléndez-Alafort L, Rodríguez-Cortés J, Ferro-Flores G, Arteaga De Murphy C, Herrera-Rodríguez R, Mitsoura E, *et al.* Biokinetics of (99m) Tc-UBI 29-41 in humans. Nucl Med Biol 2004;31:373-9.
- Ferro-Flores G, de María Ramírez F, Meléndez-Alafort L, de Murphy CA, Pedraza-López M. Molecular recognition and stability of ⁹⁹mTc-UBI 29-41 based on experimental and semiempirical results. Appl Radiat Isot 2004;61:1261-8.
- Carrasco-Hernandez J, SolÃs-Laral H, Altamirano-Ley J, Avila-Rodriguez M. Measured human dosimetry of ⁶⁸Ga-DOTA-UBI 29-41, a potential tracer for imaging bacterial infection processes. J Nucl Med 2016;57:1020.