

Seasonality, morningness-eveningness, and sleep in common non - communicable medical conditions and chronic diseases in a population

Syaron Basnet^{1,2}
Ilona Merikanto^{1,3}
Tuuli Lahti^{1,3}
Satu Männistö¹
Tiina Laatikainen^{1,4,5}
Erkki Vartiainen¹
Timo Partonen¹

¹ National Institute for Health and Welfare (THL), Department of Public Health Solutions - Helsinki - Finland.

² University of Helsinki, Department of Public Health, - Helsinki - Finland.

³ University of Helsinki, Department of Psychology, - Helsinki - Finland.

⁴ University of Eastern Finland,, Institute of Public Health and Clinical Nutrition, - Kuopio - Finland.

⁵ Hospital District of North Karelia, - Joensuu - Finland.

ABSTRACT

Introduction: The seasonal pattern for mood and behaviour, the behavioural trait of morningness-eveningness, and sleep are interconnected features, that may serve as etiological factors in the development or exacerbation of medical conditions. **Methods:** The study was based on a random sample of inhabitants aged 25 to 74 years living in Finland. As part of the national FINRISK 2012 study participants were invited (n=9905) and asked whether the doctor had diagnosed or treated them during the past 12 months for chronic diseases. **Results:** A total of 6424 participants filled in the first set of questionnaires and 5826 attended the physical health status examination, after which the second set of questionnaires were filled. Regression models were built in which each condition was explained by the seasonal, diurnal and sleep features, after controlling for a range of background factors. Of the chronic diseases, depressive disorder was associated with longer total sleep duration ($p<.0001$) and poor sleep quality ($p<.0001$). Of the measurements for health status assessment, none associated with sleep features, but systolic blood pressure yielded significant ($p<.0001$) associations with both seasonal and diurnal features at large. **Conclusion:** Sleep quality was the most sensitive probe in yielding associations with chronic diseases in this population-based study. The seasonal variations in mood and social activity, and the ease in getting up and tiredness in the morning were the most sensitive probes in yielding associations with blood pressure and waist circumference. Assessment of sleep quality, seasonal and diurnal features provides thus added value for health surveys of the general population.

Keywords: Affective Symptoms; Chronobiology Disorders; Population; Sleep Wake Disorders.

Corresponding author: Syaron Basnet.
E-mail: syaronbasnet@gmail.com
Received: May 4, 2017; Accepted: March 25, 2018.

DOI: 10.5935/1984-0063.20180017

INTRODUCTION

Circadian-clocks create an intrinsic time-tracking system that measures the passage of time in our tissues, generates and maintains the endogenous circadian rhythms¹. These clocks, enable organisms to adapt their behavior (such as: feeding, sleeping, and mating) and physiological functions (such as: cardiovascular activity, endocrine functions, body temperature and hepatic metabolism) with the environmental changes caused by the rotation of the earth along with its trajectory around the sun²⁻⁴. Disruptions in these circadian alignments with environmental entrainment factors may manifest sleep disorders, hormonal imbalances, chronic diseases, and reduces life-span⁵.

Measures of these circadian rhythms, could be an important determinant of the health status. For example, sleep problems are linked with health complaints in shift workers, and among those with existing chronic illnesses⁶⁻¹⁰. Further, to certain extent of a problem, reoccurring seasonal variations in mood and behavior (seasonality) seems to impair well-being by causing atypical depressive symptoms, such as carbohydrate craving, overeating, weight gain, hypersomnia, lack of energy, and decreased libido that are common atypical symptoms in seasonal affective disorder^{11,12}. In a similar way, the extreme traits of morningness-eveningness (chronotype), which is based on the intrinsic tendency of individuals to wake-up and fall-asleep at particular times of the day: possibly affects sleep schedules, contributes relapses of an illness, disturbs physical activity patterns, delays cognitive performance, affects endocrine functions, and overall behavioral choices¹³⁻¹⁶.

Circadian alignment with environmental entrainment factors contribute to biological changes that may catalyse to be an etiological factor in the development and exacerbation of disease and conditions including cardiovascular and metabolic disorder¹⁷. Such factors include, sleep, seasonality and chronotype that are interconnected features, but if disrupted, may compromise the health status¹⁸. To elaborate, poor sleep is suggested to elevate inflammatory markers like cytokines, and increase oxidative stress^{10,19}. A shortage of sunlight during winter months may lead to inadequate resetting of the circadian clocks, and thereby links with the etiology of chronic diseases and pronounced seasonality²⁰.

Moreover, the duration and intensity variation in light exposures influences the relapse of an illness throughout the year based on individual chronotype¹⁸. It is also suggested that eveningness associates with or even predisposes a range of common chronic diseases which are not limited to bronchial asthma, cardiovascular diseases, depressive disorders, seasonal affective disorder, sleep disorders, spinal diseases, and type-2 diabetes, but also a range of common medical conditions such as: anxiety, burnout, obesity, and substance use²¹⁻³⁴.

Moreover, non-communicable chronic diseases impose the largest public health burden globally. This burden accounted for 68% (38 million) of the world's 56 million deaths in the year of 2012³⁵, however, it extends beyond mortality through its impact on health with larger financial consequences. One

possible etiological mechanism for these diseases is a possible misalignment between a time-givers (Zeitgebers) and the circadian rhythms like sleep-wake cycle. Hence, by understanding the relationship between chronic diseases and circadian systems biology for sleep characteristics, chronotype and seasonality, a new perspective on the course, treatment, and outcome of these diseases could be achieved.

Thus, in the present study, we have analyzed the association of three indicators of circadian alignment with individual (sleep characteristics and chronotype) and environmental factors (seasonality) with common chronic non-communicable diseases and medical conditions at population level. In addition, we studied how routine objective health examination measurements were related to these three indicators.

METHODS

Participants

The national FINRISK-study, is a Finnish population-based health-examination survey conducted in every five-years since 1972 in Finland. For the present study, in 2012, a random sample of 10,000 adults aged 25 to 74-years living in five geographical areas of Finland, stratified by sex and age, were derived from the population information system of the Finnish national population register center. At start of the study, there were 9905 individuals alive and living in Finland, and they were invited to participate.

A total of 6424 participants filled in the first-set of questionnaires, and they reported their total-sleep duration and sleep quality. A total of 5826 participants attended the health examination, after which they were asked to fill in and send back the second set of questionnaires where they reported their seasonal variations in mood and behavior, and their diurnal preference in behavior. The participant rate was 64%.

Assessment

The study included two-sets of self-administered questionnaires that included structured questions on socioeconomic factors, medical history, health behavior, psychosocial factors, and a physical examination of health status accompanied with laboratory test measurements. These were sent by regular-mail together with an invitation to attend a health status examination in a local health care center or other facility near to the participant's residence. The participants filled in the questionnaire at home and returned it at the health status examination where it was checked by the staff and, if needed, completed with the participant. At the health status examination, the participants were given another questionnaire which they returned via regular mail to the institute.

Health examination measurements in the current study included height (cm), weight (kg), body-mass index (BMI, kg per m²), systolic and diastolic blood pressures (mean of three measurements in mmHg), waist circumference (cm), and the daily energy consumption (basic metabolic rate as assessed with bioimpedance measurement [TBF300 MA, Tanita, Arlington

Heights, IL, USA] in kcal per day). These physical measurements were made at the health status examination in a local health care center or other facility near the participant's residence. They were made by nurses who were trained by the staff at the institute for two weeks before the start of the study.

Seasonal variations were assessed with a modified self-rating Global Seasonality Score (GSS), the key subscale of the Seasonal Pattern Assessment Questionnaire (SPAQ)³⁶. It consists of the answers given to the six items asking, to what degree the participant's sleep duration, social activity, mood, weight, appetite, and energy level change with the seasons? Each item was scored on a Likert-like scale as 0 (no variation) to 3 (marked variation). The behavioral trait of morningness-eveningness was assessed with the six-items (items 4, 7, 9, 15, 17 and 19) derived from the original nineteen-items Morningness-Eveningness questionnaire (MEQ)³⁷.

These six items explained 83% of the variation in the original MEQ sum score, with their Cronbach's alpha of .80³⁸. The item-4 asks the ease in getting up in the morning, and it was categorized into: not easy to get up (not at all or not very easy) *vs.* easy to get up (quite easy or very easy). The item-7 asks the feeling of tiredness in the morning, and it was categorized into: feeling tired during the first half-hour after having woken in the morning (very tired or quite tired) *vs.* feeling rested (quite rested or very rested). The item-9 asks the early morning performance in some physical exercise, and it was categorized into: feeling difficult performing in morning hours (would feel very difficult or would feel quite difficult) *vs.* feeling in good condition (would be in moderate condition or would be in good condition).

The item-15 asks the alternative time slots for hard physical work if free to plan the day, and it was categorized into: morning-hour choices (11 am to 1 pm or 8 to 10 am) *vs.* evening-hour choices (7 to 9 pm or 3 to 5 pm). The item-17 asks the choice for working hours as a 5-hour block, and it was scored on a Likert-like scale and coded as: 1 (five consecutive hours starting between 5 pm and 4 am), 2 (five-consecutive hours starting between 2 and 5 pm), 3 (five-consecutive hours starting between 9 am and 2 pm), 4 (five-consecutive hours starting between 8 and 9 am), or 5 (five-consecutive hours starting between 4 and 8 am). The item-19 asks the opinion about being a morning or evening type of person, and it was categorized into: evening types (definitely evening type or more evening than morning type) *vs.* morning types (more morning than evening type or definitely morning type).

Sleep variables were subjectively reported for total sleep duration and sleep quality. Total sleep duration was based on the response about the average sleep duration (in hours and minutes) in 24 hours. There was a single item asking about sleep quality: "Do you think that you sleep enough?": "Yes, almost always; Yes, often; Seldom or nearly never; I cannot say". Sleep duration was asked with three separate items: "How many hours do you sleep on average a) at night, b) per day (night sleep plus daytime naps as together)?", "What is your usual bedtime (when you are going to go to bed and have sleep) a) on workdays/

weekdays, b) on free days/in weekends?", and "What is your usual wake-up time (when you are not going to go back to bed) a) on workdays/weekdays, b) on free days/in weekends?".

Covariates

Background information covariates were age (in years), gender (male or female), civil status as living with somebody (either married, cohabitating or registered partnership) *vs.* alone (either single, separated or divorced, or widow), education as low (less than 4 years of high school), medium (either high school only or 1 to 3 years post high school) or high (4 or more years post high school) levels, region as living in North Karelia and Kuopio, North Savo, Turku and Loimaa, Helsinki and Vantaa, or in Oulu.

Lifestyle covariates were smoking as smokers (either smoked daily or occasionally) *vs.* non-smokers (smoked not at all), alcohol consumption as alcohol consumption (at least once or more than once a month) *vs.* no alcohol consumption (no alcohol consumption at all), and physical activity as regular exercise (either several times a week or at least 3-4 hours per week or) *vs.* no exercise.

The participants were asked if (yes or no) the doctor had diagnosed or treated them in the past 12 months for the following common non-communicable medical conditions and chronic diseases: hypertension, high cholesterol, cardiac insufficiency, angina pectoris, diabetes, cancer, bronchial asthma, chronic obstructive pulmonary disease (COPD), gallstones, rheumatoid arthritis, other joint diseases, degenerative arthritis of the back, depressive disorder, other mental disorders, renal failure, and proteinuria. These sixteen variables were the primary outcome measures for the study.

Statistical analysis

Group differences were calculated, and the statistical significance was tested. Univariate and binary logistic regression models with the health examination measurements and the medical conditions and chronic diseases as the dependent variable, and the GSS items, sleep variables and MEQ items as the independent explanatory variables were generated, after controlling for BMI and the background information and lifestyle covariates (age, gender, civil status, education level, region, smoking, alcohol consumption, and physical activity).

No seasonal variation in the respective GSS item, good sleep quality, easy getting up in the morning, well rested, good condition, morning hours, and morning type categories were used as the reference. The level of significance was adjusted for the number of statistical tests we calculated (294 tests) by using the conservative Bonferroni correction, and thus the *p*-values of less than .00017 were considered as significant. The data were analyzed with the IBM SPSS statistics 21 software.

Ethics

The data collection was conducted according to the guidelines of the Declaration of Helsinki and international ethical standards.

The Ethics Committee of the Hospital District of Helsinki and Uusimaa (HUS) approved the research protocols. All the participants gave a written informed consent.

RESULTS

The 6424 participants (3383 women, 3041 men) were aged 51 years on average. Of the participants, 19% slept for less than 7 hours, 63% slept for 7 to 8 hours, and 17% slept for more than 8 hours per night. Altogether, 14% of the participants (8% of women, 6% of men) reported poor sleep quality. About 70% of the participants reported seasonal variations in sleep duration (73%), social activity (71%), mood (72%) or energy level (75%), and about 40% those in weight (46%) or appetite (43%).

Of the 4689 participants (2562 women, 2127 men) with the complete assessment of seasonality, 168 (3.58%) had the variations to the extent equal to seasonal affective disorder, 429 (9.15%) equal to subsyndromal seasonal affective disorder, and 4092 (87.27%) had normal seasonality. Of the 4414 participants (2450 women, 1964 men) with the complete assessment of chronotype, 595 (3.5%) were evening types ("night owls"), 1884 (42.7%) were intermediate types, and 1935 (43.8%) were morning types ("morning larks"). Of the health examination measurements, systolic blood pressure was significantly ($p < .0001$) associated with all the six items of seasonality, except that of weight, and with all the six items of morningness-eveningness, except that of choice for working hours (see Table 1 & Table 2).

In addition, diastolic blood pressure yielded significant associations more with diurnal features, whereas the waist circumference more with seasonal ones, and the daily energy consumption per day was associated with the seasonal variations in weight and appetite.

Concerning the medical conditions and chronic diseases, depressive disorder was significantly associated with poor sleep quality ($\beta = .459$, $p < .001$) and longer total sleep durations ($\beta = 3.33$, $p < .001$) (see Table 3). Depressive disorder associated with the seasonal variations in mood ($OR = .46$, $p < .05$) and appetite ($OR = .51$, $p < .01$). Poor sleep quality was reported in almost all the chronic diseases with the highest significant odds for gallstones ($OR = 21.59$, $p < .01$) and COPD ($OR = 4.89$, $p < .01$).

Morningness-eveningness in relation to the outcome is reported in Table 4. Bronchial asthma was significantly associated with morning tiredness ($OR = 1.69$, $p < .05$) and being an evening type ($OR = .46$, $p < .01$) (see Table 4 for details).

DISCUSSION

In the current study, we analyzed, whether key features of seasonality, morningness-eveningness and sleep were associated with the presence of common non-communicable medical conditions or chronic diseases and assessed their independent contributions. This study is, to our best knowledge, the first one to report such associations on population level. This study reveals two major findings as follows:

First, of the 16 medical conditions and chronic diseases assessed, depressive disorder and bronchial asthma yielded the greatest number of associations with the explanatory variables.

Of these, the associations of sleep features with depressive disorder remained significant after adjustment for multiple testing. Thus, this current finding, which emerged from the single-item analysis corroborated the earlier reports on the association of depressive disorder with sleep³⁹.

This finding fails to reproduce earlier reports with the association of depressive disorder with global seasonality⁴⁰ or chronotype³² which both thus appear to be more complex constructs than the self-reported sleep quality or duration. In contrast, there are epidemiological and clinical studies, reporting strong correlation between sleep disorders and depression^{41,42}. For example, in a large community-based population study, the self-reported short sleep duration and increased sleep disturbances were independently associated with increased cortisol secretion suggesting chronic stress⁴³. In bronchial asthma, we found significant associations with eveningness, this result reciprocates with Merikanto et al.²⁹. Furthermore, there are others reporting associations of poor sleep quality with breathing abnormalities in respiratory diseases⁴⁴⁻⁴⁶, while not many exist to validate the current finding.

Second, of the five health examination measurements we analyzed, systolic blood pressure that has its approximate 24-hour (i.e., circadian) rhythm was significantly associated with a number of the explanatory variables, except those of sleep features. Interestingly, energy consumption per day, as assessed with bioimpedance measurement, was significantly associated with the seasonal variations in weight and appetite, but not with the remaining seasonal variations. Thus, this finding fits in the view that the global seasonality is a mixture of two components, as being evidenced by loadings on two factors^{47,48}, where the one includes weight and appetite and the other includes the remaining.

There are limitations that need to be addressed for the current study. First, the data on the medical conditions and chronic diseases, seasonality, morningness-eveningness, and sleep length and quality were based on self-reports, which were provided as part of the health examination study. Thus, some assessment noise and misclassification may have occurred. However, the health examination measurements were assessed with objective methods. Second, the study design was cross-sectional, leaving the causal relationships between the outcome and explanatory variables unanswered. Third, the associations should be interpreted with caution due to a small number of cases for some outcomes. Further, earlier research suggests that the direct genetic effect on the chronotype equals to 40% to 70%, while the rest is influenced by environmental factors such as age, physical activity, meal time and melatonin⁴⁹. These factors need to be addressed in further studies.

Despite limitations, there are also strengths in the current study. First, it was based on a large population-based data, covering large areas of the country, due to which the results are generalizable, and the potential risk of recruitment bias is reduced. Second, the medical conditions and chronic diseases which were based on the subjective report of the participants were also clinically verified by the diagnoses assessed or treatment provided by

Table 1. Correlation coefficient (r) values for the seasonality, sleep, and morningness-eveningness in relation to the five health measurements.

Measurement	Seasonality (GSS items)						Sleep	
	Sleep length	Social activity	Mood	Weight	Appetite	Energy level	Sleep duration	Sleep quality
Body-mass index	-.008	-.001	.000	-.003	-.004	-.004	-.053**	.012
Systolic blood pressure	.056****	.109****	.112****	-.013	.068****	.094****	-.004	-.008
Diastolic blood pressure	.032*	.055****	.062****	-.031*	.027	.040**	-.029	-.020
Waist circumference	.065****	.077****	.079****	-.148****	-.035*	.052***	-.025	-.010
Energy consumption per day	.006	.007	.019	-.224****	-.114****	-.011	-.016	.003

Abbreviations: GSS=Global Seasonality Score Questionnaire. Covariates of the partial correlations include age and gender. Reference group: GSS items=no seasonal variation in the respective item; Sleep quality=good sleep. Significance indicated as $p=*$ <0.05, $**$ <0.01, $***$ <0.001, $****$ <0.0001.

Table 2. Correlation coefficient (r) values for morningness-eveningness in relation to the five health measurements.

Measurements	Morningness-eveningness (MEQ items)					
	Not easy to get up	Feeling morning tiredness	Poor early morning performance	Evening hours choice	Evening choice of consecutive hours	Evening type
Body-mass index	-.026	-.037*	-.006	-.005	.013	-.010
Systolic blood pressure	.146****	.204****	.128****	.076****	.010	.104****
Diastolic blood pressure	.104****	.128****	.131****	.038*	-.010	.085****
Waist circumference	.086****	.116****	.059***	.061****	.021	.025
Energy consumption per day	.045**	.058***	.009	.055***	.003	.032

MEQ=Morningness-Eveningness Questionnaire
Significance indicated as $p=*$ <0.05, $**$ <0.01, $***$ <0.001, $****$ <0.0001

Table 3. Odds ratios with 95% confidence intervals, or beta values with standard errors, for the seasonality, sleep and morningness-eveningness in relation to the 16 outcome measures.

Condition	Seasonality (GSS items)						Sleep	
	Sleep length	Social activity	Mood	Weight	Appetite	Energy	Sleep quality	Sleep duration
	OR (95% CI)							Beta (s.e.)
Hypertension	.83 (.59-1.17)	1.19 (.82-1.71)	1.14 (.77-1.66)	.99 (.71-1.39)	.86 (.61-1.21)	.80 (.53-1.19)	1.01 (.62-1.65)	-.006(.075)
High cholesterol	.923 (.66-1.28)	.95 (.66-1.36)	1.05 (.72-1.53)	1.14 (.82-1.58)	1.02 (.73-1.43)	1.01 (.69-1.49)	1.22 (.77-1.93)	-.07(.073)
Cardiac insufficiency	.615 (.26-1.44)	1.18 (.50-2.79)	.99 (.38-2.53)	.78 (.36-1.68)	1.20 (.53-2.67)	.96 (.36-2.52)	3.55 (1.37-9.14)**	-.017(.173)
Angina pectoris	.82 (.34-1.97)	.41 (.14-1.15)	1.12 (.43-2.88)	1.18 (.53-2.64)	.77 (.33-1.76)	.66 (.22-1.93)	1.60 (.52-4.89)	-.085(.400)
Diabetes	1.23 (.73-2.06)	1.20 (.68-2.08)	1.01 (.55-1.83)	1.17 (.69-1.97)	.72 (.42-1.23)	.78 (.41-1.45)	1.11 (.47-2.56)	.131(.113)
Cancer	.97 (.39-2.35)	1.48 (.56-3.90)	.82 (.28-2.36)	2.47 (.92-6.58)	2.40 (.82-6.97)	.26 (.07-.84)*	2.00 (.54-7.38)	.168(.207)
Bronchial asthma	.81 (.46-1.40)	.77 (.41-1.43)	.81 (.43-1.52)	.86 (.50-1.44)	1.18 (.68-2.01)	1.51 (.80-2.83)	2.41 (1.30-4.47)**	.227(.113)*
COPD	.71 (.24-2.10)	1.81 (.59-5.48)	.84 (.25-2.75)	.61 (.22-1.64)	.99 (.35-2.75)	.43 (.11-1.63)	4.90 (1.26-18.93)*	.565(.207)**
Gallstones	1.81 (.22-14.86)	.48 (.03-6.00)	.45 (.02-7.77)	3.51 (.53-22.9)	1.57 (.23-10.64)	.43 (.023-8.01)	21.60 (2.76-168.8)**	-.014(.415)
Rheumatoid arthritis	.83 (.33-2.06)	1.30 (.49-3.40)	1.65 (.60-4.51)	.66 (.25-1.70)	3.18 (1.06-9.55)*	.82 (.29-2.28)	.79 (.18-3.38)	-.176(.199)
Other joint disease	.70 (.45-1.08)	1.26 (.79-1.98)	.82 (.50-1.32)	.66 (.44-.98)*	.86 (.57-1.28)	1.05 (.63-1.71)	1.16 (.65-2.06)	.066(.090)
Degenerative arthritis	.88 (.62-1.24)	1.07 (.73-1.57)	.58 (.38-.87)**	.82 (.58-1.13)	1.20 (.85-1.68)	.86 (.56-1.30)	1.69 (1.09-2.60)**	.081(.075)
Depressive disorder	1.09 (.62-1.90)	1.00 (.51-1.92)	.46 (.20-.99)*	1.09 (.65-1.83)	.51 (.29-.87)**	.68 (.30-1.50)	3.33 (1.82-6.08)****	.459(.115)****
Other mental disorder	1.80 (.76-4.22)	1.59 (.57-4.37)	.80 (.25-2.51)	1.09 (.44-2.63)	.60 (.24-1.49)	.45 (.12-1.65)	1.71 (.56-5.22)	.562(.184)**
Renal failure	.26 (.01-7.42)	1.43 (.12-15.90)	.80 (.04-13.80)	4.00 (.29-54.82)	3.89 (.19-77.59)	.55 (.02-12.92)	14.43 (.53-387.49)	.67(1.670)
Proteinuria	.25 (.03-2.16)	2.16 (.57-8.10)	.72 (.12-4.10)	1.11 (.32-3.79)	.48 (.13-1.69)	.34 (.03-3.46)	2.43 (.57-10.21)	.981(-.007)

Abbreviations: GSS=Global Seasonality Score; MEQ=Morningness-Eveningness Questionnaire; OR=odds ratio; CI=confidence interval; s.e.= standard error; COPD=chronic obstructive pulmonary disease. Reference groups: GSS items=no seasonal variation in the respective item; Sleep quality=good sleep; Covariates of the regression models include the body-mass index, age, gender, region, civil status, education level, alcohol intake, and smoking. Significance indicated as $p=*$ <0.05, $**$ <0.01, $***$ <0.001, $****$ <0.0001.

Table 4. Odds ratios with 95% confidence intervals, or beta values with standard errors, for morningness-eveningness in relation to the 16 outcome measures.

Condition	Morningness-eveningness (MEQ items)					
	Not easy to get up	Having morning tiredness	Poor early morning performance	Evening hours choice	Evening choice of consecutive hours	Evening type
	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	Beta (S.E)	OR (95%CI)
Hypertension	.93 (.59-1.46)	1.39 (.95-2.00)	1.01 (.73-1.37)	1.16 (.76-1.77)	.036(.197)	.80 (.57-1.10)
High cholesterol	1.29 (.82-2.03)	.84 (.57-1.22)	.89 (.65-1.20)	1.50 (.99-2.26) *	.027(.199)	.78 (.57-1.07)
Cardiac Insufficiency	1.15 (.38-3.44)	1.65 (.68-3.97)	.31 (.13-.70) **	1.65 (.62-4.38)	.056(.497)	1.58 (.72-3.45)
Angina Pectoris	1.32 (.36-4.75)	.43 (.14-1.30)	.92 (.41-2.01)	1.84 (.67-5.06)	-.110(.482)	.82 (.36-1.82)
Diabetes	1.17 (.58-2.37)	1.24 (.69-2.20)	.92 (.56-1.50)	1.65 (.88-3.07)	.399(.330)	.70 (.42-1.16)
Cancer	1.00 (.28-3.51)	1.37 (.48-3.85)	.57 (.24-1.33)	.41 (.08-1.89)	.768(.684)	1.54 (.66-3.56)
Bronchial asthma	.63 (.30-1.27)	1.69 (.98-2.92) *	1.79 (1.08-2.92) *	.82 (.41-1.61)	-.350(.288)	.46 (.27-.77) **
COPD	1.37 (.25-7.51)	.37 (.09-1.42)	1.60 (.61-4.17)	1.25 (.31-4.95)	-0.64(.579)	.85 (.31-2.32)
Gallstone	4.65 (.26-80.53)	.01 (0-0.47) *	3.02 (.43-21.08)	1.25 (.08-17.45)	-2.13(1.02) *	1.46 (.22-9.53)
Rheumatoid arthritis	.67 (.20-2.18)	1.66 (.62-4.40)	1.65 (.67-4.01)	2.12 (.73-6.11)	.155(.537)	1.19 (.47-3.00)
Other joint diseases	1.13 (.63-2.02)	.70 (.43-1.12)	1.13 (.77-1.65)	1.38 (.83-.29)	-.034(.245)	.96 (.64-1.41)
Degenerative arthritis	1.25 (.79-1.97)	.78 (.53-1.15)	1.34 (.97-1.83)	1.24 (.82-1.86)	.161(.195)	.99 (.71-1.37)
Depressive disorder	.90 (.43-1.86)	1.27 (.71-2.24)	1.35 (.82-2.20)	.79 (.37-1.65)	-.370(.305)	.94 (.55-1.58)
Other mental disorder	1.26 (.31-5.14)	.49 (.15-1.49)	.80 (.34-1.82)	2.93 (1.07-8.04) *	.111(.488)	1.08 (.44-2.63)
Renal failure	-	14.43 (.53-387.4)	.65 (.06-6.62)	-	1.089(1.500)	.70 (.06-7.22)
Proteinuria	.77 (.11-5.14)	.77 (.17-3.32)	.80 (.23-2.69)	1.04 (.19-5.44)	-.043(.721)	1.27 (.35-4.52)

Abbreviations: Reference group MEQ items=Morningness-Eveningness Questionnaire: easy to get up, feeling rested in the morning, feels good about early morning performance, morning hour choice, morning choice of consecutive hours, and morning type; OR=odds ratio; CI=confidence interval; s.e.=standard error; COPD=chronic obstructive pulmonary disease.

Covariates of the regression models include the body-mass index, age, gender, region, civil status, education level, alcohol intake, and smoking. Significance indicated as $p < .05$, $** < 0.01$, $*** < 0.001$, $**** < 0.0001$.

a medical doctor. Third, the present study compares the circadian alignment with environmental entrainment factors together, while the other studies assessed the same circadian alignment independently with the same population study^{39,40,50}.

CONCLUSION

To conclude, we herein analyzed the seasonality, morningness-eveningness, and sleep features simultaneously in the same statistical model which we controlled for a range of confounding factors and whose results we adjusted for multiple testing. We found that poor sleep quality contributed most to the outcome in case of depressive disorder but was not associated with any of the health examination measurements.

ACKNOWLEDGEMENT

The first author (S.B.) has been supported by research grants from Filha (Finnish Lung Health Association, 02042016), Respiratory Diseases Research Foundation (Hengityssairauksien tutkimussäätiö, 05052015), Juho Vainio Foundation (2016-2018), Orion Research Foundation (01112016-17), and Signe and Ane Gyllenberg Foundation (11052016).

REFERENCES

- Panda S, Hogenesch JB, Kay SA. Circadian rhythms from flies to human. *Nature*. 2002;417(6886):329-35.
- Magnusson A, Partonen T. The diagnosis, symptomatology, and epidemiology of seasonal affective disorder. *CNS Spectr*. 2005;10(8):625-34.
- Gerhart-Hines Z, Lazar MA. Rev-erb α and the circadian transcriptional regulation of metabolism. *Diabetes Obes Metab*. 2015;17 Suppl 1:12-6.
- Panda S. Circadian physiology of metabolism. *Science*. 2016;354(6315):1008-15.
- Bass J, Lazar MA. Circadian time signatures of fitness and disease. *Science*. 2016;354(6315):994-9.
- Healey ES, Kales A, Monroe LJ, Bixler EO, Chamberlin K, Soldatos CR. Onset of insomnia: role of life-stress events. *Psychosom Med*. 1981;43(5):439-51.
- Pailhoux E, Benoit O, Goldenberg F, Bouard G, Payant C. Psychological profile and sleep organization in young subjects with poor quality of sleep. *Psychiatry Res*. 1988;26(3):327-36.
- Hyypä MT, Kronholm E. Quality of sleep and chronic illnesses. *J Clin Epidemiol*. 1989;42(7):633-8.
- Abbott SM, Reid KJ, Zee PC. Circadian Rhythm Sleep-Wake Disorders. *Psychiatr Clin North Am*. 2015;38(4):805-23.
- Morris CJ, Purvis TE, Hu K, Scheer FA. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc Natl Acad Sci U S A*. 2016;113(10):E1402-11.
- Rosenthal NE, Sack DA, Gillin JC, Lewy AJ, Goodwin FK, Davenport Y, et al. Seasonal affective disorder. A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry*. 1984;41(1):72-80.
- Partonen T, Lönnqvist J. Seasonal affective disorder. *Lancet*. 1998;352(9137):1369-74.

13. Hagenauer MH, Lee TM. The neuroendocrine control of the circadian system: adolescent chronotype. *Front Neuroendocrinol.* 2012;33(3):211-29.
14. Roenneberg T, Allebrandt KV, Merrow M, Vetter C. Social jetlag and obesity. *Curr Biol.* 2012;22(10):939-43.
15. Goel N, Basner M, Rao H, Dinges DF. Circadian rhythms, sleep deprivation, and human performance. *Prog Mol Biol Transl Sci.* 2013;119:155-90.
16. Wennman H, Kronholm E, Partonen T, Peltonen M, Vasankari T, Borodulin K. Evening typology and morning tiredness associates with low leisure time physical activity and high sitting. *Chronobiol Int.* 2015;32(8):1090-100.
17. Luyster FS, Strollo PJ, Zee PC, Walsh JK; Boards of Directors of the American Academy of Sleep Medicine and the Sleep Research Society. Sleep: a health imperative. *Sleep.* 2012;35(6):727-34.
18. Brambilla C, Gavinelli C, Delmonte D, Fulgosi MC, Barbini B, Colombo C, et al. Seasonality and sleep: a clinical study on euthymic mood disorder patients. *Depress Res Treat.* 2012;2012:978962.
19. Motivala SJ. Sleep and inflammation: psychoneuroimmunology in the context of cardiovascular disease. *Ann Behav Med.* 2011;42(2):141-52.
20. Laposky AD, Bass J, Kohsaka A, Turek FW. Sleep and circadian rhythms: key components in the regulation of energy metabolism. *FEBS Lett.* 2008;582(1):142-51.
21. Drennan MD, Klauber MR, Kripke DF, Goyette LM. The effects of depression and age on the Horne-Ostberg morningness-eveningness score. *J Affect Disord.* 1991;23(2):93-8.
22. Adan A. Chronotype and personality factors in the daily consumption of alcohol and psychostimulants. *Addiction.* 1994;89(4):455-62.
23. Richardson GS. The human circadian system in normal and disordered sleep. *J Clin Psychiatry.* 2005;66 Suppl 9:3-9.
24. Levy AJ, Lefler BJ, Emens JS, Bauer VK. The circadian basis of winter depression. *Proc Natl Acad Sci U S A.* 2006;103(19):7414-9.
25. Lucassen EA, Zhao X, Rother KI, Mattingly MS, Courville AB, de Jonge L, et al.; Sleep Extension Study Group. Evening chronotype is associated with changes in eating behavior, more sleep apnea, and increased stress hormones in short sleeping obese individuals. *PLoS One.* 2013;8(3):e56519.
26. Merikanto I, Kronholm E, Peltonen M, Laatikainen T, Lahti T, Partonen T. Relation of chronotype to sleep complaints in the general Finnish population. *Chronobiol Int.* 2012;29(3):311-7.
27. Merikanto I, Lahti T, Kronholm E, Peltonen M, Laatikainen T, Vartiainen E, et al. Evening types are prone to depression. *Chronobiol Int.* 2013;30(5):719-25.
28. Merikanto I, Lahti T, Puolijoki H, Vanhala M, Peltonen M, Laatikainen T, et al. Associations of chronotype and sleep with cardiovascular diseases and type 2 diabetes. *Chronobiol Int.* 2013;30(4):470-7.
29. Merikanto I, Englund A, Kronholm E, Laatikainen T, Peltonen M, Vartiainen E, et al. Evening chronotypes have the increased odds for bronchial asthma and nocturnal asthma. *Chronobiol Int.* 2014;31(1):95-101.
30. Merikanto I, Lahti T, Seitsalo S, Kronholm E, Laatikainen T, Peltonen M, et al. Behavioral trait of morningness-eveningness in association with articular and spinal diseases in a population. *PLoS One.* 2014;9(12):e114635.
31. Merikanto I, Kronholm E, Peltonen M, Laatikainen T, Vartiainen E, Partonen T. Circadian preference links to depression in general adult population. *J Affect Disord.* 2015;188:143-8.
32. Antypa N, Vogelzangs N, Meesters Y, Schoevers R, Penninx BW. Chronotype associations with depression and anxiety disorders in a large cohort study. *Depress Anxiety.* 2016;33(1):75-83.
33. Merikanto I, Suvisaari J, Lahti T, Partonen T. Eveningness relates to burn-out and seasonal sleep and mood problems among young adults. *Nord J Psychiatry.* 2016;70(1):72-80.
34. Merikanto I, Lahti T, Seitsalo S, Kronholm E, Laatikainen T, Peltonen M, et al. Eveningness has the increased odds for spinal diseases but the decreased odds for articular diseases with prospective hospital treatments. *Biol Rhythm Res.* 2017;48(2):263-74.
35. World Health Organization (WHO). Global status report on noncommunicable diseases 2014. Geneva: WHO; 2014.
36. Rosenthal N, Bradt G, Wehr T. Seasonal Pattern Assessment Questionnaire (SPAQ). Bethesda: National Institute of Mental Health; 1984.
37. Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol.* 1976;4(2):97-110.
38. Hätönen T, Forsblom S, Kieseppä T, Lönnqvist J, Partonen T. Circadian phenotype in patients with the co-morbid alcohol use and bipolar disorders. *Alcohol Alcohol.* 2008;43(5):564-8.
39. Basnet S, Merikanto I, Lahti T, Männistö S, Laatikainen T, Vartiainen E, et al. Associations of common chronic non-communicable diseases and medical conditions with sleep-related problems in a population-based health examination study. *Sleep Sci.* 2016;9(3):249-54.
40. Basnet S, Merikanto I, Lahti T, Männistö S, Laatikainen T, Vartiainen E, et al. Seasonal variations in mood and behavior associate with common chronic diseases and symptoms in a population-based study. *Psychiatry Res.* 2016;238:181-8.
41. Nutt D, Wilson S, Paterson L. Sleep disorders as core symptoms of depression. *Dialogues Clin Neurosci.* 2008;10(3):329-36.
42. Wells RD, Day RC, Carney RM, Freedland KE, Duntley SP. Depression predicts self-reported sleep quality in patients with obstructive sleep apnea. *Psychosom Med.* 2004;66(5):692-7.
43. Kumari M, Badrick E, Ferrie J, Perski A, Marmot M, Chandola T. Self-reported sleep duration and sleep disturbance are independently associated with cortisol secretion in the Whitehall II study. *J Clin Endocrinol Metab.* 2009;94(12):4801-9.
44. Sundar IK, Yao H, Huang Y, Lyda E, Sime PJ, Sellix MT, et al. Serotonin and corticosterone rhythms in mice exposed to cigarette smoke and in patients with COPD: implication for COPD-associated neuropathogenesis. *PLoS One.* 2014;9(2):e87999.
45. Choudhary SS, Choudhary SR. Sleep effects on breathing and respiratory diseases. *Lung India.* 2009;26(4):117-22.
46. Koyanagi A, Garin N, Olaya B, Ayuso-Mateos JL, Chatterji S, Leonardi M, et al. Chronic conditions and sleep problems among adults aged 50 years or over in nine countries: a multi-country study. *PLoS One.* 2014;9(12):e114742.
47. Rintamäki R, Grimaldi S, Englund A, Haukka J, Partonen T, Reunanen A, et al. Seasonal changes in mood and behavior are linked to metabolic syndrome. *PLoS One.* 2008;3(1):e1482.
48. Grimaldi S, Englund A, Partonen T, Haukka J, Pirkola S, Reunanen A, et al. Experienced poor lighting contributes to the seasonal fluctuations in weight and appetite that relate to the metabolic syndrome. *J Environ Public Health.* 2009;2009:165013.
49. Baron KG, Reid KJ. Circadian misalignment and health. *Int Rev Psychiatry.* 2014;26(2):139-54.
50. Basnet S, Merikanto I, Lahti T, Männistö S, Laatikainen T, Vartiainen E, et al. Associations of common noncommunicable medical conditions and chronic diseases with chronotype in a population-based health examination study. *Chronobiol Int.* 2017; 34(4):462-70.