

Treatment of Advanced Prostate Cancer: Where Are We in 2019?

Tratamento do câncer avançado de próstata: Onde estamos em 2019?

Fernando Sabino Marques Monteiro^{1,2,3} , Andrey Soares^{1,4,5}, Fernando Nunes Galvão de Oliveira^{1,6}, Pedro Castilhos de Freitas Crivelaro⁷, Pablo Moura Barrios⁷, Andre Poisl Fay^{1,7}

ABSTRACT

In the last decade, important advances have been made in the treatment of metastatic prostate cancer, resulting in a better understanding of the biology underlying the disease, and in the approval of several therapeutic agents such as immunotherapy, new generation antiandrogens, cytotoxic chemotherapies, and radiopharmaceuticals. All these recent advances have been incorporated in clinical guidelines and a critical analysis of the data available should be important to help the decision-making process. In addition, the incorporation of well established therapies in early disease stages have demonstrated a robust overall survival gain for patients with castration-sensitive metastatic prostate cancer. However, no predictive biomarkers of response are available and the selection of the best therapeutic option is still challenging depending on clinical and pathological factors. Many questions related to the optimal sequencing of agents, or comparison of its efficacy remain unanswered.

Keywords: Prostate; Review/drug therapy; Receptors; Androgen; Androgen Antagonists; Advanced Treatment; Prostate-Specific Antigen

1. Latin American Cooperative Oncology Group - LACOG GU, Genito Urinary Session - Porto Alegre - Rio Grande do Sul - Brazil
2. Hospital Santa Lucia, Departamento de Oncologia - Brasília - Distrito Federal - Brazil
3. Hospital Universitário de Brasília, Unidade de Oncologia - Brasília - Distrito Federal - Brazil
4. Hospital Israelita Albert Einstein, Departamento de Oncologia - São Paulo - São Paulo - Brazil [5] - Centro Paulista de Oncologia - CPO, Oncologia Clínica - São Paulo - São Paulo - Brazil
6. Clínica de Oncologia - Clion, Oncologia Clínica - Salvador - Bahia - Brazil
7. Pontifícia Universidade Católica - PUC, Faculdade de Medicina - Porto Alegre - Rio Grande do Sul - Brazil

Financial support: none to declare.

Conflicts of interest: The authors declare no conflict of interest relevant to this manuscript.

Corresponding author: Fernando Sabino Marques Monteiro. Hospital Santa Lucia. SHLS 716 Bloco C. Zip-code:70390-700. fsabinocba@gmail.com

Received on: July 19, 2018 | **Accepted on:** March 13, 2019

DOI: 10.5935/2526-8732.20190001

RESUMO

Na última década, tem havido avanços importantes no tratamento de câncer de próstata metastático, o que resultou em uma melhor compreensão da biologia subjacente da doença e a aprovação de vários agentes terapêuticos, tais como imunoterapia, nova geração anti-andrógenos, quimioterapias citotóxicas e radiofármacos. Todos esses avanços recentes foram incorporados às diretrizes clínicas e uma análise crítica dos dados disponíveis seria importante para ajudar no processo de tomada de decisão. Além disso, a incorporação de terapias bem estabelecidas nos estágios iniciais da doença demonstrou um aumento robusto na sobrevida global em pacientes com câncer de próstata metastático sensível à castração. No entanto, não existem biomarcadores preditivos de resposta disponíveis e a seleção da melhor opção terapêutica permanece um desafio, dependendo dos fatores clínicos e patológicos. Muitas questões relacionadas ao sequenciamento ótimo de agentes ou à comparação de sua eficácia permanecem sem resposta.

Descritores: Próstata; Revisão/terapia medicamentosa; Receptores Androgênicos; Antagonistas de andrógenos; Tratamento Avançado; Antígeno específico da próstata

INTRODUCTION

Prostate cancer (PCa) is the most prevalent tumor among men except for non-melanoma skin cancers. According to Brazilian National Cancer Institute (INCA), 68,220 new cases of prostate cancer are estimated for 2018 being the second leading cause of cancer deaths in Brazil.⁽¹⁾

A portion of patients with early stage disease submitted to locoregional treatment will relapse of the disease and may require systemic treatment. There is another part of the patients presenting with metastatic disease at diagnosis (metastatic “di novo”) and will also need systemic treatment.

During the last decade, the treatment of metastatic prostate cancer has dramatically changed with the approval of seven new agents for patients with metastatic castration-resistant prostate cancer (mCRPC). These new therapeutic options have provided an overall survival (OS) increase from 9-18 months⁽²⁾ to an average of 30 months⁽³⁾ and also improved patients quality of life (QoL). In addition, recent studies have shown that the use of chemotherapy or new

generation antiandrogen in a selected population with metastatic castration-sensitive prostate cancer (mCSPC) is associated with a significant improvement of OS of approximately 13 months.⁽⁴⁾

Although several agents have resulted in improvement in OS in different scenarios of metastatic disease, head to head comparisons of efficacy and the best sequencing approach are questions that remain unanswered. Despite the advances in the tumor biology understanding, no predictive biomarkers of response are validated to be used in the clinical practice in order to choose the best therapeutic option. Unfortunately, clinical characteristics, such as PSA doubling time, exposure time to primary ADT, extent and location of metastatic disease, performance status (PS) and pathological features have been still used as tools to guide treatment selection.

In this manuscript, we review the latest evidence for treatment of mCSPC and mCRPC discussing the best options in each scenario and, in the absence of approved predictive biomarkers tests, how to choose subsequent treatments.

METASTATIC CASTRATION SENSITIVE PROSTATE CANCER (mCSPC)

When metastatic disease is diagnosed, initial treatment of prostate cancer is testosterone suppression at castration levels. It can be done through surgical orchiectomy or drug blockade with GnRH antagonist or GnRH agonist. This stage of disease is called sensitive- castration prostate cancer.^(5,6)

Despite the suppression of testosterone levels, after an average time of 18 to 24 months, the majority of patients invariably will present progression of disease (elevation of PSA and/or worsening of pre-existing metastatic lesions and/or the appearance of new metastatic lesions).⁽⁷⁾

The use of peripheral antiandrogens (bicalutamide, nilutamide, flutamide, etc.) associated with LHRH

agonists (complete hormonal blockade) can be used for a short period of 2 to 4 weeks to avoid the worsening of symptoms (pain, urinary retention, etc.) that may be caused by a transient increase in testosterone (flare phenomenon).⁽⁸⁾ Long-term complete hormonal blockade may be an initial option with some meta-analyses suggesting OS benefit of 3% to 5% after 10 years of follow-up, but with increase toxicity and worsening of QoL.^(9,10)

Recently three prospective randomized phase 3 studies (Table 1) evaluated the benefit of the combination of docetaxel with ADT in the setting of the mCSPC.

The French trial GETUG-AFU 15 randomized 385 patients with mCSPC to ADT alone or in combination with docetaxel 75mg/m² every 3 weeks for up to 9 cycles (ADT + D). This study demonstrated a benefit of median OS for the ADT plus docetaxel group but without statistical significance (46.5 vs 60.9 months, HR = 0.90, p = 0.44), independently of the extent of metastatic disease at diagnosis.^(11,12)

On the other hand, the American trial CHAARTED-ECOG 3805 that randomized 790 patients with mCSPC for ADT alone or ADT + D every 3 weeks for up to 6 cycles, demonstrated better OS (around 13 months) for the ADT + D group (44 vs 57.6 months, HR = 0.61, p < 0.001).

A pre-specified subgroup analysis was performed according to the stage of the disease at the diagnoses. In those defined as high-volume disease (visceral metastasis or ≥ 4 bone lesions with one lesion in the appendicular skeleton) the benefit of OS was even more expressive, approximately 17 months (32.2 vs 49.2 months, HR = 0.60, P < 0.001), but in those defined as low-volume disease there was no statistical benefit of OS.⁽⁴⁾ In addition, the QoL decreased initially (at 3 months; p = 0.02) but improve after 12 months and even it better statistically (p = 0.04).

In the same context as previous trials, the STAMPEDE trial randomized 2962 patients with high risk localized disease or mCSPC to ADT or ADT + D every 3 weeks for up to 6 cycles and showed an improvement in OS for ADT + D group (71 vs 81 months, HR = 0.78, p = 0.006). In the patients with metastatic disease (61%) the OS benefit was even greater, around 22 months (HR = 0.76).^(14,15)

A meta-analysis including several trials, including the three trials discussed above, evaluated the role of chemotherapy in men with high-risk localized or mCSPC and demonstrated an improvement in 4 year-survival of 9% (HR = 0.77; p < 0.0001) for ADT + D in those with metastatic disease.⁽¹⁶⁾

In, 2017, the Multicentric LATITUDE Trial randomized 1199 patients to abiraterone plus ADT or ADT alone. The patients were required to have at least two of the three following high-risk factors associated with poor prognosis: a Gleason score of 8 or more (on a scale of 2 to 10, with higher scores indicating more aggressive disease), at least three bone lesions and the presence of measurable visceral metastasis. Radiographic progression free survival (HR = 0.47, 95% CI:0.39-0.55, p < 0.001) (rPFS) and OS (56.3 vs 36.5 months, HR = 0.66, p < 0.0001), both primary endpoints, was significantly improved with the addition of abiraterone.⁽¹⁷⁾ A non pre-planned analysis regarding volume of disease (CHARTEED criteria) showed an improvement only in patients with high volume disease (high volume: HR = 0.62, p < 0.0001 and low volume: HR = 0.72, p = 0.1242).⁽¹⁸⁾

The STAMPEDE trial had an arm that evaluated the benefit of adding abiraterone to ADT in patients with localized high risk and mCSPC. This strategy compared to ADT alone showed an improvement in OS (HR = 0.63, p < 0.001) and in failure free survival (HR = 0.29, p < 0.001). The OS in the 52% of the metastatic population was even better (HR = 0.61).⁽¹⁹⁾ An exploratory analysis in patients with mCSPC from

Table 1. Comparative summary of phase 3 studies with in mCSPC.

Study	N	Follow-up (months)	High-volume disease (%)	Gleason ≥ 8 (%)	Age (years)	Patients who received the planned treatment (%)	Overall Survival (months)	HR and p
GETUG-AFU 15 ⁽¹¹⁾	385	84	52	55	64	48	46.5 (ADT) vs 60,9 (ADT+CHEMO)	0.9 and 0.44
CHAARTED ⁽⁴⁾	790	29	65	65	63	74	44 (ADT) vs 57.6 (ADT+CHEMO)	0.61 and < 0.001
STAMPEDE ⁽¹⁴⁾	2962	43	NI	74	65	77	71 (ADT) vs 81 (ADT+CHEMO)	0.77 and < 0.0001
LATITUDE ⁽¹⁶⁾	1199	30	NI	100	68	88	34.7 (ADT) vs NR (ADT + ABI)	0.62 and < 0.001
ARCHES ⁽²⁴⁾	1150	14.4	62	67	70	NI	NR (ADT) vs NR (ADT + ENZA)	0.81 and 0.3361

STAMPEDE evaluated efficacy of abiraterone in patients with low and high risk per LATITUDE criteria and low and high volume per CHARTEED criteria. This analysis showed a statistically improvement of adding abiraterone in all scenarios.⁽²⁰⁾

In absence of a head-to-head trial, several comparisons studies, meta-analysis and a directly comparison in the STAMPEDE trial try to answer if abiraterone is better than docetaxel in mCSPC setting. There is no evidence that one is better than the other in this scenario.^(15,21-23)

Recently another new generation anti-androgen, enzalutamide, showed efficacy in mCSPC. ARCHES trial showed that addition of enzalutamide to ADT improves outcomes over ADT alone. Primary endpoint of rPFS favored enzalutamide plus ADT (HR = 0.39, $p < 0.0001$). All subgroup analysis was statistically in favor of combination, including patients with low and high volume (CHARTEED criteria) and prior docetaxel use or not. OS data are still immature.⁽²⁴⁾

Local treatment was explored in mCSPC in two randomized trials comparing with standard of care (SOC). HORRAD trial compares the addition of

radiotherapy to the prostate in patients with bone metastasis CSPC and use of ADT. OS was similar between groups (45 vs 43 months, HR = 0.9, $p = 0.4$). Patients with less than 5 bone metastasis had the best outcome, but still not statistically significant (HR = 0.68, CI = 0.42-1.1).⁽²⁵⁾ The STAMPEDE RT trial failed to demonstrate OS advantage in patients with mCSPC that received local radiotherapy plus SOC *versus* SOC (HR = 0.92, $p = 0.266$). In a pre-planned analysis patients with low volume disease (CHARTEED criteria) had an OS improvement (49.1 vs 45.4 months, HR = 0.68, CI = 0.52-0.90).⁽²⁶⁾

Based on these two trials local treatment in mCSPC patients are not recommended routinely. It must be discussed in a tumor board session and with patients regarding efficacy and toxicities. There are a few trials evaluating the addition of local treatment in patients with mCSPC and new evidences will be available soon.

The choice of treatment should include a discussion with the patient about the potential toxicities associated with abiraterone and docetaxel, as well as the duration, oncologist's experience and access, patient's desire and cost of treatment.

METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (mCRPC)

Castration-resistant prostate cancer is defined as an elevation in PSA levels between two consecutive measurements (minimum interval of 3 weeks), with serum testosterone below 50 ng/dl, in the presence of hormonal blockade.⁽²⁷⁾ Even in the castration resistance setting, androgen receptors (AR) still plays a key role. Molecular analyses of tissue samples from patients with mCRPC have demonstrated that the overexpression of AR and the increased sensitivity of these receptors (through overexpression of their nuclear coactivators) make possible their action even in an environment with low testosterone concentration.⁽²⁸⁻³¹⁾ Understanding the importance of the AR and knowing the production and metabolism of androgens in the scenario of mCRPC was fundamental for the development of therapeutic agents.

There are currently seven therapeutic agents, with different mechanisms of action and outcomes for the treatment of mCRPC (Table 2).

Docetaxel

Until 2010, docetaxel (taxane) was the only agent with OS improvement and it was the standard first-line therapy for mCRPC. The phase III study TAX 327 randomized 1006 mCRPC patients to receive Docetaxel 75mg/m² every 3 weeks for up to 10 cycles *versus* Docetaxel 30mg/m² weekly *versus* Mitoxantrone 12mg/m² every 3 weeks and showed an improvement of OS for Docetaxel every 3 weeks (19.2 vs 17.8 vs 16.9 months, HR = 0.79, $p = 0.004$) in all

subgroups. (symptomatic vs asymptomatic, presence vs absence of visceral metastases, good KPS vs poor KPS and age < 68 years vs ≥ 69 years), The patients in the docetaxel every 3 weeks group also presented a higher PSA response rate of 50% and improvement in QoL.^(32,40) Adverse events (AE) such as neuropathy (30%), fatigue (53%), alopecia (65%), diarrhea (32%) and neutropenia grade 3 or 4 (30%) were more common with docetaxel, however the treatment discontinuation rate was similar between groups.

Another option for the use of docetaxel is the schedule every 2 weeks, based on a randomized phase 3 study of 177 patients which compared docetaxel 50mg/m² every 2 weeks *versus* docetaxel 75mg/m² every 3 weeks and demonstrated an increase in time to treatment failure (5.6 vs 4.9 months, $p = 0.01$), a lower incidence of grade 3 and 4 neutropenia (36 vs 53%) and lower incidence of febrile neutropenia (4 vs 14%) for the use every 2 weeks.⁽⁴¹⁾

Cabazitaxel

Cabazitaxel is a new generation taxane developed to overcome resistance to docetaxel. In addition, this drug has the capacity to cross the blood-brain barrier.⁽⁴²⁾ Its activity was demonstrated in the phase 3 TROPIC trial that randomized 755 mCRPC patients who had progression during or after therapy with docetaxel. These patients received cabazitaxel 25 mg/m² every 3 weeks associated with prednisone 10 mg daily for up to 10 cycles (C+P) or mitoxantrone 12 mg/m² every 3 weeks also associated with prednisone.

Table 2. Summary of therapeutic options for mCRPC.

	Scenario/Stage	Administration	Corticoid	PSA response	OS
Docetaxel	CRPC - M1	IV each 3 weeks	Yes	Yes	2.9 months ⁽³²⁾
Cabazitaxel	CRPC - M1 After docetaxel	IV each 3 weeks	Yes	Yes	2.4 months ⁽³³⁾
Abiraterone acetate	CRPC - M1 before or after Docetaxel	Oral Daily	Yes	Yes	Before Docetaxel: 5.2 months ⁽³⁴⁾ After Docetaxel: 4.6 months ⁽³⁵⁾
Enzalutamide	CRPC - M0/M1 before or after Docetaxel	Oral Daily	No	Yes	M0: immature ⁽⁸⁷⁾ Before Docetaxel: 4.0 months ⁽³⁶⁾ After Docetaxel: 4.8 months ⁽³⁷⁾
Sipuleucel-T*	CRPC - M1 before or after Docetaxel	IV each 2 weeks	No	No	4.1 months ⁽³⁸⁾
Radium-223	CRPC - M1 before or after Docetaxel	IV each 4 weeks	No	Uninformed	3.6 months ⁽³⁹⁾
Apalutamide	CRPC - M0	Oral Daily	No	Yes	M0 immature ⁽⁹⁸⁾
Darolutamide*	CPEC - M0	Oral twice Daily	No	Yes	M0 immature ⁽¹¹²⁾

* Not available in Brazil.

With a median follow-up of 13.7 months, this study showed an improvement in OS (15.1 vs 12.7 months, HR = 0.70, 95% CI 0.59-0.83, $p < 0.0001$) and in progression-free survival (PFS) (2.8 vs 1.4 months, HR = 0.74, 95% CI 0.64-0.86, $p = 0.0002$) for those treated with C+P.⁽³³⁾ Treatment with C+P was associated with higher toxicity, such as diarrhea (47%), fatigue (37%), neutropenia grade 3 and 4 (82%), and febrile neutropenia (8%).

Due mainly to hematological toxicity, the non-inferiority phase 3 PROSELICA trial evaluated the efficacy of cabazitaxel in a lower dose. In this study, 1200 patients with mCRPC who failed docetaxel were randomized to cabazitaxel 25mg/m² (C25) every 3 weeks or to cabazitaxel 20mg/m² (C20) every 3 weeks and no inferiority was demonstrated in OS or PFS at lower dose. In addition, grade 3 and 4 AE (39.5% C20 vs 54.5% C25), grade 4 neutropenia (21.3% C20 vs 48.6% C25) and neutropenic infection (2.2% C20 vs 6.1% C25) were less frequent at the lower dose.⁽⁴³⁾

Therefore, cabazitaxel at a dose of 20mg/m² every 3 weeks is one of the options for second-line mCRPC (after docetaxel failure) treatment.

Abiraterone acetate

Abiraterone acetate is a hormonal agent that inhibits androgen biosynthesis through inactivation of the CYP-17 enzyme, acting in the three sources of androgen production: testis, adrenal gland and intra-tumor. Its activity was demonstrated in mCRPC as first line treatment and as second line (after docetaxel failure).

The phase 3 COU-AA 302 trial randomized 1088 chemotherapy-naïve mCRPC patients, asymptomatic or oligosymptomatic, to abiraterone 1g/day and prednisone 10mg/day or to placebo and prednisone 10mg/day. A follow-up of 49 months showed improvement of OS (34.7 vs 30.3 months, HR = 0.81, $p = 0.003$) and rPFS (16.5 vs 8.2 months, HR = 0.52, $p < 0.0001$) for the abiraterone group. However, 44% of patients in the placebo group received abiraterone later, which may have influenced the magnitude of the results.⁽⁴⁴⁾ All secondary outcomes (time to initiation of chemotherapy, skeletal events, progression of PSA, deterioration of KPS, etc.) and QoL were significantly favorable for abiraterone.⁽⁴⁵⁾

The COU-AA 301 phase 3 trial randomized 1195 mCRPC patients who failed to docetaxel to abiraterone 1g/day and prednisone 10g/day or placebo and prednisone 10mg/day. With a 20 months median follow-up an improvement in OS was demonstrated (15.8 vs 11.2 months, HR = 0.74, $p < 0.001$) for the abiraterone group. This group had also benefit in all secondary outcomes (time to PSA progression, rPFS, PSA response).⁽⁴⁶⁾ An exploratory analysis of this study also showed benefit of abiraterone for pain relief, time for pain worsening and reduction of skeletal-related events.⁽⁴⁷⁾

In general, toxicity of abiraterone is related to the excess of mineralocorticoids, so the concomitant use of prednisone is necessary. Water retention (33%), hypertension (11%) and hypokalemia (18%) are among the most frequent AE. Other AE observed are changes in liver function tests (11%) and

miscellaneous cardiac disorders (16%). Despite the toxicity mentioned above, the rate of discontinuation of treatment related to AE was similar to that of the placebo group.⁽³⁵⁾

Thus, abiraterone 1g/day and prednisone 10mg/day is an option for mCRPC either in the first or second line treatment (after docetaxel).

Enzalutamide

Enzalutamide is a strong AR blocker, thus, it acts directly by blocking the AR, preventing nuclear translocation of the AR and inhibiting the binding of the receptor to the nucleus DNA. Its efficacy in mCRPC was demonstrated in M0 and M1 disease setting, in both chemotherapy-naïve patients and in those who failed to docetaxel. Three important phase 3 trials evaluated enzalutamide in mCRPC setting. Two trials evaluated enzalutamide in metastatic castration-resistant prostate cancer patients (M1CRPC) and one trial included only non-metastatic castration-resistant prostate cancer patients (M0CRPC) with high-risk disease (rapidly rising PSA).

The PREVAIL trial randomized 1717 chemotherapy-naïve mCRPC patients, asymptomatic or oligo-symptomatic, to enzalutamide 160mg/day or placebo. With a median follow-up of 22 months it was shown improvement in OS (34.7 vs 30.2 months, HR = 0.71, $p < 0.001$) and in secondary outcomes (rPFS and reduction of skeletal events) for the enzalutamide group. In addition, there was also a significant improvement in QoL for the enzalutamide group.^(36,48) Another study, the AFFIRM trial randomized 1199 mCRPC patients previously treated with docetaxel to enzalutamide 160mg/day or placebo. The enzalutamide group showed improvement in OS (18.4 vs 13.6, HR = 0.63, $p < 0.001$) and in all secondary outcomes (PSA response rate, time to PSA progression, rPFS). In a secondary analysis of the trial, those treated with enzalutamide had a significant improvement in QoL.^(37,49)

The most common AE related to enzalutamide are fatigue (34%), hot flashes (20%) and diarrhea (21%). In the AFFIRM trial some patients that received enzalutamide presented seizures and, due to this, those with a past of seizures and/or recent brain event were excluded from the PREVAIL trial.⁽³⁶⁾

Thus, enzalutamide 160 mg/day is an option for M0CRPC high-risk disease treatment and for M1CRPC first and second line treatment.

Sipuleucel-T

Sipuleucel-T is an autologous vaccine of dendritic cells developed to increase T cell-mediated immune response against prostatic acid phosphatase. Its activity in mCRPC was demonstrated in the phase 3 IMPACT trial. This trial randomized 512 mCRPC patients (asymptomatic or oligosymptomatic, previously treated with docetaxel or not) to sipuleucel-T every 2 weeks for 3 cycles or placebo. With a median follow-up of 34 months it was showed an OS improvement (25.8 vs 21.7 months, HR = 0.78, $p = 0.03$) for the vaccine group. However, neither improvement in PFS nor in PSA response rate was observed.⁽³⁸⁾

This therapy is well tolerated and, as in other immunotherapies, the most common vaccine-related AE were chills (53%), fatigue (41%), and fever (35%).⁽⁵⁰⁾

Therefore, sipuleucel-T is an option of for mCRPC, but it is not available outside the USA.

Radium-223

Radium-223 is an α -particle emitter therapeutic agent which has tropism for bone tissue. It is able to achieve bone metastases with high energy in a small radius of action, minimizing toxicity in the bone marrow.

The phase 3 ALSYMPCA trial randomized 921 mCRPC patients previously treated or not with docetaxel, symptomatic and with bone metastases only (no known visceral metastases) to receive radium-223 every 4 weeks up to six cycles or placebo. This trial demonstrated an improvement in OS for those treated with radium-223 as first or second line of treatment (14.9 vs 11.3 months, HR = 0.70, $p < 0.001$). Treatment was well tolerated and there was no significant difference in grade 3 or 4 toxicity when compared with the control group.^(39,51) However, the reduction to the first bone event was only observed in the subgroup of patients who received radium-223 as second line of treatment.⁽⁹⁵⁾ Regardless of the line of treatment, the use of radium-223 was associated with a significant improvement in QoL.⁽⁹⁶⁾

Thus, radium-223 is another option of treatment - as first or second line (after docetaxel) - for mCRPC with bone metastases.

NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (M0CRPC)

The systemic treatment of patients who are castrate resistant and have not still developed metastatic disease have been recently studied. In the era of Gallium-68 prostate-specific membrane antigen (PSMA) positron emission tomography (PET), which has a higher sensitivity and specificity in identifying metastatic disease, this scenario in progressively less

common in clinical practice.⁽¹¹¹⁾ However, different agents have shown to be associated with clinical benefit in this population.

The use of enzalutamide was compared to placebo in the PROSPER trial. This trial enrolled 1401 men with a PSA doubling time of ≤ 10 months and a serum PSA ≥ 2 ng/mL. Metastasis-free survival (MFS) - the primary endpoint- was longer with enzalutamide (39.6 vs 14.7 months, HR = 0.29, 95% CI: 0.24-0.35,

$p < 0.001$). Time to first use of a new antineoplastic agent and time to PSA progression (39.6 vs 17.7 months, HR = 0.21, 95% CI: 0.17-0.26, $p < 0.001$ and 37.2 vs 3.9 months, HR = 0.07, 95% CI: 0.05-0.08, $p < 0.001$, respectively) have also favored the use of enzalutamide.⁽⁹⁶⁾

Apalutamide is a nonsteroidal antiandrogen agent that binds directly to the ligand-binding domain of the AR and prevents AR translocation, DNA binding, and AR mediated transcription.⁽⁹⁷⁾ The Spartan trial is a double blind phase 3 trial involving men with M0CRPC and a PSA doubling time of 10 months or less. In this trial, 1207 men were randomized to apalutamide ($n = 806$) or placebo ($n = 401$). The primary endpoint (MFS) was longer in the apalutamide group (40.5 vs 16.2 months, HR = 0.28, 95% CI: 0.23-0.35, $P < 0.001$). All secondary endpoints, time to metastasis,

PFS and time to symptomatic progression were significantly longer with apalutamide than with placebo ($P < 0.001$ for all comparisons).⁽⁹⁸⁾

Recently, darolutamide, another antiandrogen, has been tested in the same clinical scenario (ARAMIS study). This double blind, placebo-controlled phase III trial randomized 1,509 patients in a 2:1 ratio. Patients were stratified by PSA doubling time (≤ 6 months or > 6 months) and use of osteoclast-targeted therapy. Median MFS was 40.4 months with darolutamide vs 18.4 months with placebo (HR 0.41; 95% CI 0.34-0.50; $p < 0.0001$). In addition, there was a trend favoring darolutamide in terms of OS (Median not reached for either arm, HR 0.71, 95% CI 0.50-0.99, $p = 0.045$).⁽¹¹²⁾ Importantly, robust data on OS are lacking to define the real benefit of these three new agents.

SEQUENCING

Advances in treatment of mCRPC and availability of new therapeutic agents are notorious, but many questions related to optimal sequencing or the comparison of efficacy between these new agents remain unanswered. Until then, there are no randomized studies comparing directly the agents of different classes.

Recent data suggest that regardless of the line of treatment and its sequence, patients who are exposed to as many agents as possible have improvement in OS.⁽⁵²⁾

Thus, the search and identification of predictive biomarkers of response would be one of the tools to select the best agent and/or the best time to use it. In this scenario, several studies have evaluated the identification of androgen receptor variants (AR-V) as a possible biomarker. It is known that the presence of AR-V is one of the mechanisms of resistance to castration, since it gives the AR constant activity without requiring stimulation for circulating androgens.

Several variants of the AR have been identified, but the most common and best characterized is variant 7 (AR-V7).^(53,54) The potential role of AR-V7 as a biomarker in CRPC has been demonstrated in a retrospective study of 31 patients treated with enzalutamide and 31 patients treated with abiraterone. Those with the presence of AR-V7 in RNA from circulating tumor cells (CTC) did not show a PSA response rate for either enzalutamide or abiraterone (unlike those in whom the presence of AR-V7 was not identified). Positivity for AR-V7 was also associated with lower PSA progression-free survival (pPFS) ($p < 0.001$), lower radiological and clinical progression free survival ($p < 0.001$), and lower OS ($p = 0.006$).⁽⁵⁵⁾

In another retrospective study, 37 patients with CRPC were treated with taxanes and the presence of AR-V7 in CTC was identified in 46% of them. Unlike

the study with enzalutamide and abiraterone, the presence of AR-V7 did not interfere with the rate of PSA response with chemotherapy.⁽⁵⁶⁾

Despite these results, it is worth mentioning that these are retrospective studies with small number of patients, and it is necessary to await the results of prospective studies in progress that can define AR-V7 as a predictive or only prognostic factor.

A recent prospective study involved 202 patients that received either enzalutamide or abiraterone. The patients were submitted to an analysis of the RNA of CTC to verify quantification of androgen AR and to examine the prognostic value of this data. Patients with high amounts of receptors correlated with the presence of variant 7 (AR-V7). In addition, the number of copies of the AR showed a prognostic value in patients treated with abiraterone or enzalutamide: the higher the number of copies, the lower PFS (laboratory, radiological and clinical) and the lower OS.⁽⁹⁹⁾

In the absence of randomized studies comparing the agents and the lack of predictive biomarkers, the tools for choosing the first or second line of treatment are the clinical features of each patient. It is important to consider the performance status of the patient, comorbidities and symptoms, clinical-pathological characteristics of the disease as extension of metastatic disease, presence of undifferentiated or small cell components and toxicity profile of each agent. In addition, patient's preference (intravenous or oral), cost and availability of the agents should be taken into account.

Retrospective studies suggest a relationship between the time of exposure to the initial hormone therapy and the activity of the hormonal agents in the castration-resistant setting. In this context, lower exposure (< 16 months vs ≥ 16 months) or the short-lived response (< 12 months vs ≥ 12 months) to the initial hormone therapy is associated with a greater probability of not responding to subsequent

treatment with abiraterone or enzalutamide.^(57,58) On the other hand, in studies that evaluated time of exposure or time of response to initial hormone therapy and subsequent treatment with taxanes, despite the lower biochemical PFS and the lower OS, taxanes responded regardless of the time of treatment of initial hormone therapy.^(59,60)

Gleason score is a prognostic factor established in localized disease.⁽⁶¹⁾ In metastatic disease, an exploratory analysis of the study TAX 327 demonstrated that the improvement of OS with docetaxel is independent of Gleason score ($p = 0.009$).⁽⁶²⁾ Likewise, a retrospective analysis including all patients of the pivotal studies involving abiraterone (COU-AA 301 and COU-AA 302) showed that patients with Gleason score ≥ 8 also had improvement in OS ($p < 0.0001$).⁽⁶³⁾

Undifferentiated tumors or tumors with small cell components are rare, aggressive, more common in young patients, and usually with low PSA regardless of the volume of metastatic disease.⁽⁶⁴⁾ In this situation, hormone therapy has limited action and chemotherapy is preferentially used.⁽⁶⁵⁾

First Line Treatment

The options for first-line mCRPC treatment available in Brazil, the inclusion criteria, as well as the most common adverse effects are summarized in Table 3.

Unlike TAX 327 and ALSYMPCA trials, in which 45% and 100% of the patients, respectively, were symptomatic, in PREVAIL and COU-AA 302 trials the symptomatic patients were excluded. Patients with visceral metastasis were excluded of COU-AA 302 and ALSYMPCA trials, but represented 11% of PREVAIL trial patients and 22% of TAX 327 trial patients.^(32,34,36,39)

Regarding to lymph node disease, those with lymph node enlargement ($> 3\text{cm}$) were excluded from the ALSYMPCA trial.

Recently, a prospective multicenter phase 3b trial of 839 mCRPC patients evaluated Radium-223 in some different scenarios of the ALSYMPCA trial. In this trial, 20% of the patients were asymptomatic and Abiraterone (18%) or Enzalutamide (5%) were allowed to be used associated with Radium-223. Subgroup analysis according to the symptomatology and level of alkaline phosphatase demonstrated a greater OS for asymptomatic patients and for those with alkaline phosphatase levels within normality.⁽⁹⁵⁾

The improvement of OS in these subgroups raises the question whether this benefit occurs from the early onset of Radium-223 (at the earliest stage of the metastatic disease) or from the more indolent and less aggressive disease.

As previously discussed, due to seizures in some patients that received enzalutamide after prior therapy with docetaxel (AFFIRM trial), those with a historic of seizures or some recent brain event (e.g. stroke) were also excluded from the PREVAIL trial.^(36,37)

Recent data from patients in a phase 2 trial evaluating the sequence of first-line treatment with abiraterone and enzalutamide in mCRPC showed a worsening of depressive symptoms and an increase of cognitive impairment in those treated with enzalutamide.⁽⁶⁶⁾

Unlike docetaxel and abiraterone, the use of enzalutamide and radium-223 does not require association with corticosteroids. Some retrospective series suggest that there is no negative impact on the efficacy and safety of abiraterone when used with a lower dose of corticosteroid (prednisone 5 mg/day), as well as the lower incidence of side effects related to prolonged corticoid use.^(67,68)

Recently docetaxel and cabazitaxel were compared in first-line treatment. The FIRSTANA trial randomized 1168 mCRPC patients to receive cabazitaxel 20mg/m²

Table 3. Options for first-line treatment of mCRPC available in Brazil.

	Docetaxel (TAX 327) ⁽³²⁾	Abiraterone (COU-AA 302) ⁽³⁴⁾	Enzalutamide (PREVAIL) ⁽³⁶⁾	Radium-223 (ALSYMPCA) ⁽³⁹⁾
Asymptomatic or oligosymptomatic	Yes	Yes	Yes	No
Symptomatic	Yes	No	No	Yes
Visceral Metastasis	Yes	No	Yes	No
≥ 2 bone metastases	Yes	Yes	Yes	Yes
Performance Status	KPS ≥ 60	ECOG 0/1	ECOG 0/1	ECOG 0/2
Seizure	Yes	Yes	No	No
Significant cardiac dysfunction	No	No (EF ≥ 50)	No (EF ≥ 45)	No
Resistant hypertension	Yes	No	No	No
Specific adverse effects	Neutropenia Neuropathy Fatigue Diarrhea	Edema Hypertension Hypokalemia Elevation of hepatic enzymes Cardiac events	Fatigue Hot flashes Hypertension Cardiac events Seizure	Thrombocytopenia Diarrhea

or cabazitaxel 25mg/m² or docetaxel 75mg/m² as first line treatment. Despite the higher response rate with Cabazitaxel 25mg/m², there was no difference in OS (24.5 vs 25.2 vs 24.3 months, $p = 0.99$). The AE and the discontinuation rate of the treatments were similar.⁽⁶⁹⁾

Currently there is another possible setting for the second-line treatment: those patients who received chemohormonal as first-line treatment in the mCSPC setting. There was no standard treatment for those patients. Table 4 summarizes the alternatives that are offered after progression with initial chemohormonal treatment.

A retrospective analysis of the GETUG-AFU 15 trial demonstrated a PSA response rate of 13-70% and symptom improvement in 16-33% of patients when exposed to other agents following progression.⁽⁷⁰⁾ Exposing these patients to docetaxel again does not seem to be a good option, considering that docetaxel re-treatment in mCRPC, despite the PSA response rate, did not show improvement in OS.⁽⁷¹⁾ In this setting, the most appropriate would be the agents that improve OS in the second-line treatment after docetaxel (Abiraterone, Enzalutamide or Cabazitaxel - see details below).

Second-line treatment after docetaxel

The second line treatment options available in Brazil, some details of each trials and the most common AE associated with these therapies are specified in Table 5.

To this moment there are no studies comparing these second line therapies or evaluating their use in sequence. Therefore, the data and results of the main phase 3 trials and some retrospective analysis of the action of these drugs when used in sequence should be of great help when choosing the most appropriate treatment.

Considering the control arms of each clinical trial in second line treatment after failure to docetaxel, placebo plus prednisone and placebo alone where used as controls arms in the COU-AA-301 and AFFIRM trials respectively. In the other hand, the TROPIC trial control group received mitoxantrone, a chemotherapy with known palliative action in pain control and PSA response rate, however with no improvement in OS.⁽³³⁾ That could explain the results found in the update of the trial that showed no difference in pain palliation and QoL in comparison with cabazitaxel.⁽⁷²⁾

Table 4. Subsequent therapies after chemo-hormone therapy.

	GETUG-AFU 15 ⁽⁷⁰⁾		CHAARTED ⁽⁴⁾		STAMPEDE ⁽¹⁴⁾	
	HT (193)	CT+HT (192)	HT (393)	CT+HT (397)	HT (1184)	CT+HT (592)
Progression		287	238	761	315	
Treatment	42%	68%	55%	88%	83%	
Docetaxel	40%	15%	51%	23%	41%	14%
Cabazitaxel	1%	2%	13%	26%	3%	7%
Abiraterone/Enzalutamide	11%	12%	38%	47%	23%/9%	28%/8%

CT: chemotherapy, HT: hormone therapy.

Table 5. Second-line trials: characteristics and adverse events.

	Cabazitaxel ⁽³³⁾ (TROPIC)	Abiraterone ⁽³⁵⁾ (COU-AA 301)	Enzalutamide ⁽³⁷⁾ (AFFIRM)	Radium-223 ⁽³⁹⁾ (ALSYMPCA)
Symptomatic	Yes	Yes	Yes	Yes
Metastasis	Bone: 84% Viscera: 24%	Bone: 89% Lymph node: 44% Liver: 10%	Bone: 92% Lymph node: 55% Lung: 15% Liver: 11%	Bone: 100%
Performance Status	ECOG 0/2	ECOG 0/2	ECOG 0/2	ECOG 0/2
Neuropathy Grade ≥ 2	No	No	No	No
Significant cardiac dysfunction	No (EF ≥ 50)	No (EF ≥ 50)	No (EF ≥ 45)	-----
Corticosteroids	Yes	Yes	No	No
Control Arm	Mitoxantrone	Placebo + Prednisone	Placebo	Placebo
Specific adverse events	Neutropenia Fatigue Diarrhea	Edema Hypertension Hypokalemia AST/ALT rising Cardiac alterations	Fatigue Hot flushes Hypertension Cardiac alterations Seizure	Thrombocytopenia Diarrhea Vomiting

The secondary endpoints of the trials were different. While COU-AA-301 and AFFIRM trials considered disease progression as evidence of tumor progression radiologically, the TROPIC trial defined disease progression as rise in PSA, evidence of tumor progression radiologically or worsening of the pain. Thus, treatment in the TROPIC trial may have been discontinued sooner in comparison to treatment in the COU-AA-301 and AFFIRM trial, therefore resulting in a shorter duration of treatment.

The TROPIC, COU-AA-301 and AFFIRM trial enrolled patients that progressed after treatment with docetaxel. In this context, 63% of the patients enrolled in the TROPIC trial and 45% of the patients in the COU-AA-301 had discontinued the treatment with docetaxel due to progression.^(73,74) That may indicate that patients enrolled in the TROPIC trial, probably, presented a higher resistance to docetaxel.

At the same time that in the COU-AA-301 and AFFIRM trials the OS benefit was seen in patients regardless of age (≥ 65 vs < 65 yrs), in the TROPIC trial there was no difference in OS in patients under 65 yrs. old.⁽³³⁾

All secondary endpoints were positive in the three trials. Time to the first skeletal event was longer in the COU-AA-301 and AFFIRM trial, however this endpoint was not evaluated in the TROPIC trial.^(33,35,37)

The median duration of treatment was 8 and 8.3 months for the COU-AA-301 and AFFIRM trials, respectively. Patients in the TROPIC trial received a median of six treatment cycles. The cross-over was allowed in the COU-AA-301 and AFFIRM trials and not allowed in the TROPIC trial.^(33,35,37)

The toxicity related to each treatment has been discussed previously. It is important to emphasize that, being chemotherapy, the hematological toxicity observed in the TROPIC trial may be a limiting factor for treatment with cabazitaxel. Regarding enzalutamide, as previously discussed, historic of seizures could be a contraindication.^(36,37) In the case of abiraterone, there is the possibility of hepatotoxicity with elevation of transaminases, which may be a limiting factor to its use in those with some type of underlying liver disease.

Second-line treatment after abiraterone or enzalutamide

There is no good level evidence for the best sequence treatment to mCRPC in this setting. Retrospective data suggest lower response rates when abiraterone and enzalutamide are used in sequence, indicating perhaps the existence of a cross-resistance mechanism between them^(81,82,85-86) (Table 6). Evidence also suggests that taxane chemotherapy may reverse or downregulate some of these resistance pathways, such as elimination of AR splice variants.⁽⁵⁵⁾ However, a retrospective series with 80 patients who were treated with abiraterone followed by enzalutamide versus abiraterone followed by docetaxel and then enzalutamide or enzalutamide followed by docetaxel

and then abiraterone, demonstrated worse progression-free survival of PSA (4.4 vs 1.6 months, $p < 0.01$) and worse rPFS (5.7 vs 2.2 months) for the use of docetaxel intercalated with the new hormonal agents.⁽⁹⁴⁾

Recently, exploring the sequencing of these two agents, a phase 2 trial with 202 M1CRPC patients evaluated abiraterone plus prednisone followed by enzalutamide (arm A) versus enzalutamide followed by abiraterone plus prednisone (arm B), both at PSA progression. The primary endpoints, PSA decline $> 50\%$ on second line therapy and time to second PSA progression, were better with arm A sequence (34 vs 4% and 2.7 vs 1.3 months, respectively).⁽⁸³⁾ In addition to this data, there are only retrospective series evaluating the activity of sequencing these agents in this setting (Table 6).

Although docetaxel was the main second line treatment in patients after disease progression on COU-AA-302 (44% of the abiraterone and 58% of the placebo group receive some second line treatment)⁽³⁴⁾ and PREVAIL trial (40% of enzalutamide and 70% of placebo group receive some second line treatment),⁽³⁶⁾ the activity of chemotherapy (docetaxel or cabazitaxel) in this setting is uncertain.

Retrospective data from patients treated with docetaxel after abiraterone demonstrated an OS of 12.5 months, lower than the 19.2 months showed by the TAX 327 trial, which might suggest the hypothesis of some degree of cross-resistance between the agents.⁽⁷⁷⁾ On the other hand, in a post-hoc analysis of COU-AA-302 evaluating docetaxel activity in patients treated previously with abiraterone, a similar PSA response rate to the TAX 327 study was demonstrated.⁽⁸⁴⁾ While some series demonstrate no benefit with docetaxel in patients that failed abiraterone (primary resistance), others show responses in these patients and do not establish a direct relation between responders and non-responders to abiraterone and responders and non-responders to docetaxel.⁽⁷⁷⁻⁸⁰⁾

Given the limitations of these retrospective data, it can also be inferred that docetaxel maintains its activity after previous use of abiraterone.

However, the response or non-response to previous treatment with abiraterone or enzalutamide does not seem to be a predictive factor for response to cabazitaxel.⁽⁸⁷⁾ As well as cabazitaxel also has activity in those patients that failed (progression ≤ 3 months) docetaxel.⁽⁸⁴⁾

Third-line treatment after docetaxel

Although a considerable number of patients have good clinical conditions to receive some treatment in this setting, there is no standard recommendation. In the absence of randomized clinical trials, some small retrospective series summarized in Table 7 have demonstrated a PSA response rate ranging from 3 to 39% and a median OS of 7 to 16 months.⁽⁸⁵⁻⁹³⁾

Table 6. Response rate and OS data of second-line treatments after failing abiraterone.

	N	Response PSA > 50%	OS (mo)
Abiraterone→Docetaxel			
De Bono et al. ⁽⁷⁵⁾	265	47%	NI
Mezynski et al. ⁽⁷⁶⁾	35	26%	12.5
Schweizer et al. ⁽⁷⁷⁾	24	38%	NI
Aggarwal et al. ⁽⁷⁸⁾	23	48%	12.4
Azad et al. ⁽⁷⁹⁾	86	35%	11.4
Abiraterone→Enzalutamide			
Azad et al. ⁽⁸⁰⁾	47	25.5%	8.6
Suzman et al. ⁽⁸¹⁾	30	34%	NI
Cheng et al. ⁽⁸²⁾	28	36%	NI

Table 7. Response rate and OS data of third-line treatments after failing docetaxel.

Docetaxel→Abiraterone→Enzalutamide			
Authors	N	PSA response	OS
Schrader et al. ⁽⁸⁵⁾	35	29%	7.1
Bianchini et al. ⁽⁸⁶⁾	39	13%	NI
Badrising et al. ⁽⁸⁷⁾	61	21%	7.3
Azad et al. ⁽⁸⁰⁾	68	22%	10.6
Docetaxel→Enzalutamide→Abiraterone			
Authors	N	PSA response	OS
Loriot et al. ⁽⁸⁸⁾	38	8%	7.2
Noonan et al. ⁽⁸⁹⁾	30	3%	11.6
Docetaxel→Abiraterone or Enzalutamide→Cabazitaxel			
Authors	N	PSA response	OS
Pezaro et al. ⁽⁹⁰⁾	41	39%	15.8
Sella et al. ⁽⁹¹⁾	24	32%	8.2
Wissing et al. ⁽⁹²⁾	69	32%	NI
Al Nakouzi et al. ⁽⁹³⁾	79	35%	10.9

New Therapeutic Agents

Given the proven efficacy of immunotherapy in several solid tumors (lung and kidney cancer, melanoma, etc.), this treatment strategy has also been evaluated in mCRPC. Ipilimumab, a humanized anti-CTLA-4 monoclonal antibody, was evaluated in two phase 3 trials. In the first trial 799 patients with mCRPC and at least one bone metastasis who had failed to docetaxel were randomized to receive bone-directed radiotherapy followed by either ipilimumab 10mg/kg or placebo every 3 weeks up to four doses. After median follow-up of 9.9 months there was no difference in OS (11.2 vs 10 months, HR = 0.85, 95% CI: 0.72-1.00, $p = 0.053$).⁽¹⁰⁰⁾ The other trial 602 asymptomatic or minimally symptomatic patients with chemotherapy-naïve mCRPC were randomized to ipilimumab 10mg/kg or placebo every 3 weeks up to four doses. No significant difference was observed in primary

endpoint OS (28.7 vs 29.7 months, HR = 1.11, 95.87% CI: 0.88-1.39, $p = 0.36$), but the secondary end point PFS was longer with ipilimumab (5.6 vs 3.8 months, HR = 0.67, 95.87% CI: 0.55-0.81).⁽¹⁰¹⁾

Pembrolizumab, a highly selective anti PD-1 humanized monoclonal antibody, demonstrated durable responses in 23 patients with mCRPC whose tumors have PD-L1 expression ≥ 1 percent in a non-randomized phase 1b trial.⁽¹⁰²⁾ The KEYNOTE-199, a recent phase 2 trial, with 258 mCRPC who had failed at least a one novel endocrine therapy (abiraterone or enzalutamide) and chemotherapy evaluated pembrolizumab 200mg every 3 weeks and showed an objective response rate (primary endpoint) of 6 percent in those with PD-L1 expression positive. The secondary endpoint disease control rate at six months was 15 percent. Interestingly, patients with BRCA 1/2 or ATM mutations had better overall response rate.⁽¹⁰³⁾

The activity combining immunotherapy was evaluated in a recent phase 2 trial with 78 asymptomatic or minimally symptomatic mCRPC patients who progressed after new generation hormone therapy and not received chemotherapy (cohort 1) or had received chemotherapy (cohort). The combination was nivolumab 1mg/Kg + ipilimumab 3mg/Kg every 3 weeks for 4 doses then nivolumab 480mg every 4 weeks. The primary endpoint ORR was 26 and 10% in cohort 1 e 2, respectively. In both cohorts the ORR was higher in patients with PD-L1 \geq 1% and DNA damage repair (DDR).⁽¹⁰⁴⁾

It has been know that mCRPC can harbor some DNA repair gene mutation like *BRCA1* or 2, *ATM*, *CHEK2*, *NBN* and *PALB2* that may be associated with more aggressive disease and poorer OS.^(105,106) Moreover, there are some trials showing durable antitumor activity in these mCRPC patients with treatment with poly-adenosine diphosphate ribose polymerase (PARP) inhibitors.⁽¹⁰⁷⁾ In this context, a phase 2 single-arm trial with 49 patients evaluated olaparib 400mg twice a day until disease progression, unacceptable toxicity or death. It should be noted that these mCRPC patients had failed to at least two prior

treatments and all of them failed to chemotherapy with docetaxel. All patients had tissue evaluated for DNA gene repair mutations and this was detected in 33 percent of patients. Olaparib was associated with a higher response rate (primary endpoint) in those patients with DNA gene repair mutations when compared to those without mutations (88 vs 6%). Also rPFS and OS were longer in those with mutations (9.8 vs 2.7 months and 13.8 vs 7.5 months, respectively).⁽¹⁰⁸⁾

Another phase 2 trial randomized 142 mCRPC patients who have failed to prior chemotherapy to olaparib and abiraterone or abiraterone. With a median follow-up of 15.9 months the rPFS was higher for olaparibe and abiraterone (13.8 vs 8.2 months, HR = 0.65, 95% CI:0.44-0.97, p = 0.034). In contrast, more AE were observed in combination treatment arm.⁽¹⁰⁹⁾

Recently the activity combining Pembrolizumab and Olaparib in heavily treated mCRPC was evaluated in one cohort of KEYNOTE-365 phase 1b/2 trial with 40 patients that showed PSA response rate and ORR of 13 and 7%, respectively.⁽¹¹⁰⁾

DISCUSSION AND FUTURE DIRECTIONS

Currently there are several options for the treatment of metastatic prostate cancer, with different mechanisms of action, toxicities, routes of administration, and costs. Importantly, all approved agents have shown improvement in OS. However, no head-to-head comparisons are available. Several efforts have been done in order to answer that question. An international retrospective epidemiologic trial of men with advanced prostate cancer (IRONMAN Trial) is ongoing collecting clinical information about this patients around the world. In addition, biological specimens have been collected to correlate clinical findings with biological characteristics. Hopefully, this registry which will include more than 5.000 patients may help us to answer those open questions.

Advances in the understanding of tumor biology have also influencing in the development of new therapeutic strategies and biomarkers of prognosis or response to available therapies. Unfortunately, no biomarkers have been validated to be used in the clinical practice. Therefore, treatment should be individualized taking into account clinical factors such as age, performance status and comorbidities; disease clinical course PSA doubling time, symptoms or not; and pathological features: Gleason score at diagnosis, neuroendocrine component. Another important factor in the decision making process is the toxicity profile of each agent that may vary according to each agent. In addition, especially in developing countries where resources and access to medications are restricted, cost should be taken in consideration.


In spite of the provocative and promising results with AR-V7, there is still no validation of it as a predictive factor and results from prospective studies are awaited, thus, testing for AR-V7 should not yet be used to guide decisions on treatment.

Considering that all agents are equally effective, practical issues such as availability, access and cost of the agents, besides the preference of patients may be decisive when choosing the agent. Flowcharts 1 and 2 suggest treatment sequences.


Based on all the prospective and retrospective data already mentioned, the suggestion is that for the majority of PCRC patients, hormonal treatment with abiraterone or enzalutamide should be initially offered, reserving chemotherapy in the case of clinical conditions, for symptomatic patients, with visceral disease, with an undifferentiated component/small cells in the pathology and/or those with progression after a short time of hormonal blockade (signal suggestive of resistance to hormonal therapy).

Retrospective data and post-hoc analyzes suggest that, if the patient has the clinical conditions, chemotherapy is the preferred option after failure to abiraterone or enzalutamide.


New discoveries are accompanied by new possibilities, however, the questions and doubts are increasing. In this manner, we await for the results of the prospective studies in progress evaluating these new agents directly and in earlier stages of the disease, as well as for the evaluation of biomarkers, so that we can understand in which patient and which moment of the disease to better use them.



Flowchart 1.



Flowchart 2.



Flowchart 3.
* Not available in Brazil.

AUTHOR'S CONTRIBUTION

Fernando Sabino Marques Monteiro: Collection and assembly of data, Conception and design, Data analysis and interpretation, Final approval of manuscript, Manuscript writing, Provision of study materials or patient

Andrey Soares: Data analysis and interpretation, Final approval of manuscript, Manuscript writing

Fernando Nunes Galvão de-Oliveira: Final approval of manuscript, Manuscript writing

Pedro Castilhos de Freitas Crivelaro: Manuscript writing

Pablo Moura Barrios: Manuscript writing

Andre Poisl Fay: Conception and design, Data analysis and interpretation, Final approval of manuscript, Manuscript writing

REFERENCES

1. Brazil. Ministério da Saúde. Instituto Nacional de Câncer José Alencar Gomes da Silva (INCA). Estimativa de incidência e mortalidade do Câncer no Brasil 2018. Rio de Janeiro: INCA; 2018.
2. Heidenreich A, Bastian PJ, Bellmunt J, Bolla M, Joniau S, van der Kwast T, et al.; European Association of Urology. EAU guidelines on prostate cancer. Part II: Treatment of advanced, relapsing, and castration-resistant prostate cancer. *Eur Urol.* 2014;65(2):467-79.
3. Omlin A, Pezaro C, Mukherji D, Mulick Cassidy A, Sandhu S, Bianchini D, et al. Improved survival in a cohort of trial participants with metastatic castration-resistant prostate cancer demonstrates the need for updated prognostic nomograms. *Eur Urol.* 2013;64(2):300-6.
4. Sweeney CJ, Chen YH, Carducci M, Liu G, Jarrard DF, Eisenberger M, et al. Chemohormonal Therapy in Metastatic Hormone-Sensitive Prostate Cancer. *N Engl J Med.* 2015;373(8):737-46.
5. Soloway MS, Chodak G, Vogelzang NJ, Block NL, Schellhammer PF, Smith JA Jr, et al. Zoladex versus orchiectomy in treatment of advanced prostate cancer: a randomized trial. Zoladex Prostate Study Group. *Urology.* 1991;37(1):46-51.
6. Seidenfeld J, Samson DJ, Hasselblad V, Aronson N, Albertsen PC, Bennett CL, et al. Single-therapy androgen suppression in men with advanced prostate cancer: a systematic review and meta-analysis. *Ann Intern Med.* 2000;132(7):566-77.
7. Damber JE, Aus G. Prostate cancer. *Lancet.* 2008;371(9625):1710-21.
8. Loblaw DA, Mendelson DS, Talcott JA, Virgo KS, Somerfield MR, Ben-Josef E, et al.; American Society of Clinical Oncology. American Society of Clinical Oncology recommendations for the initial hormonal management of androgen-sensitive metastatic, recurrent, or progressive prostate cancer. *J Clin Oncol.* 2004;22(14):2927-41.
9. Maximum androgen blockade in advanced prostate cancer: an overview of the randomised trials. Prostate Cancer Trialists' Collaborative Group. *Lancet.* 2000;355(9214):1491-8.
10. Schmitt B, Bennett C, Seidenfeld J, Samson D, Wilt T. Maximal androgen blockade for advanced prostate cancer. *Cochrane Database Syst Rev.* 2000;(2):CD001526.
11. Gravis G, Fizazi K, Joly F, Oudard S, Priou F, Esterni B, et al. Androgen-deprivation therapy alone or with docetaxel in non-castrate metastatic prostate cancer (GETUG-AFU 15): a randomised, open-label, phase 3 trial. *Lancet Oncol.* 2013;14(2):149-58.
12. Gravis G, Boher JM, Joly F, Soulié M, Albiges L, Priou F, et al.; GETUG. Androgen Deprivation Therapy (ADT) Plus Docetaxel versus ADT Alone in Metastatic Non castrate Prostate Cancer: Impact of Metastatic Burden and Long-term Survival Analysis of the Randomized Phase 3 GETUG-AFU15 Trial. *Eur Urol.* 2016;70(2):256-62.
13. Patrick-Miller L, Chen Y, Carducci MA, Cella D, DiPaola RS, Gartrell BA, et al. Quality of life (QOL) analysis from CHARTED: Chemohormonal androgen ablation randomized trial in prostate cancer (E3805). *J Clin Oncol.* 2016;34(15 Suppl):5004.
14. James ND, Sydes MR, Clarke NW, Mason MD, Dearnaley DP, Spears MR, et al.; STAMPEDE investigators. Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet.* 2016;387(10024):1163-77.
15. Feyerabend S, Saad F, Li T, Ito T, Diels J, Van Sanden S, et al. Survival benefit, disease progression and quality-of-life outcomes of abiraterone acetate plus prednisone versus docetaxel in metastatic hormone-sensitive prostate cancer: A network meta-analysis. *Eur J Cancer.* 2018;103:78-87.
16. Vale CL, Burdett S, Rydzewska LHM, Albiges L, Clarke NW, Fisher D, et al. Addition of docetaxel or bisphosphonates to standard of care in men with localized or metastatic hormone-sensitive prostate cancer: a systematic review and meta-analyses of aggregate data. *Lancet Oncol.* 2016;17(2):243-56.
17. Fizazi K, Tran N, Fein L, Matsubara N, Rodriguez-Antolin A, Alekseev BY, et al. Abiraterone plus Prednisone in Metastatic, Castration-Sensitive Prostate Cancer. *N Engl J Med.* 2017;377(4):352-60.

18. Fizazi K, Tran N, Fein LE, Matsubara N, Antolín AR, Ozguroglu M, et al. Final analysis of phase III LATITUDE study in patients (pts) with newly diagnosed high-risk metastatic castration-naïve prostate cancer (NDx-HR mCNPC) treated with abiraterone acetate + prednisone (AA + P) added to androgen deprivation therapy (ADT). *J Clin Oncol*. 2019;37(7 Suppl):141.
19. James ND, de Bono JS, Spears MR, Clarke NW, Mason MD, Dearnaley DP, et al.; STAMPEDE Investigators. Abiraterone for Prostate Cancer Not Previously Treated with Hormone Therapy. *N Engl J Med*. 2017;377(4):338-51.
20. Hoyle AP, Ali SA, James ND, Parker CC, Cook AD, Attard G, et al. Effects of abiraterone acetate plus prednisone/prednisolone in high and low risk metastatic hormone sensitive prostate cancer. *Ann Oncol*. 2018;29(Suppl 8):mdy424.033.
21. Wallis CJD, Klaassen Z, Bhindi B, Goldberg H, Chandrasekar T, Farrell AM, et al. Comparison of Abiraterone Acetate and Docetaxel with Androgen Deprivation Therapy in High-risk and Metastatic Hormone-naïve Prostate Cancer: A Systematic Review and Network Meta-analysis. *Eur Urol*. 2018;73(6):834-44.
22. Sydes MR, Spears MR, Mason MD, Clarke NW, Dearnaley DP, de Bono JS, et al. Adding abiraterone or docetaxel to long-term hormone therapy for prostate cancer: directly randomised data from the STAMPEDE multi-arm, multi-stage platform protocol. *Ann Oncol*. 2018;29(5):1235-48.
23. Kassem L, Shohdy KS, Abdel-Rahman O. Abiraterone acetate/androgen deprivation therapy combination versus docetaxel/androgen deprivation therapy combination in advanced hormone-sensitive prostate cancer: a network meta-analysis on safety and efficacy. *Curr Med Res Opin*. 2018;34(5):903-10.
24. Armstrong AJ, Szmulewitz RZ, Petrylak DP, Petrylak DP, Villers A, Azad A, Alcaraz A, et al. Phase 3 study of androgen deprivation therapy (ADT) with enzalutamide (ENZA) or placebo (PBO) in metastatic hormone-sensitive prostate cancer (mHSPC): The ARCHES trial. *J Clin Oncol*. 2019;37(Suppl 7):687.
25. Boevé LMS, Hulshof MCCM, Vis AN, Zwinderman AH, Twisk JWR, Witjes WPJ, et al. Effect on Survival of Androgen Deprivation Therapy Alone Compared to Androgen Deprivation Therapy Combined with Concurrent Radiation Therapy to the Prostate in Patients with Primary Bone Metastatic Prostate Cancer in a Prospective Randomised Clinical Trial: Data from the HORRAD Trial. *Eur Urol*. 2019;75(3):410-8.
26. Parker CC, James ND, Brawley CD, Clarke NW, Hoyle AP, Ali A, et al.; Systemic Therapy for Advanced or Metastatic Prostate cancer: Evaluation of Drug Efficacy (STAMPEDE) investigators. Radiotherapy to the primary tumour for newly diagnosed, metastatic prostate cancer (STAMPEDE): a randomised controlled phase 3 trial. *Lancet*. 2018;392(10162):2353-66.
27. Gillessen S, Omlin A, Attard G, de Bono JS, Efsthathiou E, Fizazi K, et al. Management of patients with advanced prostate cancer: recommendations of the St Gallen Advanced Prostate Cancer Consensus Conference (APCCC) 2015. *Ann of Oncol*. 2015;26(8):1589-604.
28. Chen CD, Welsbie DS, Tran C, Baek SH, Chen R, Vessella R, et al. Molecular determinants of resistance to antiandrogen therapy. *Nat Med*. 2004;10(1):33-9.
29. Visakorpi T, Hyytinen E, Koivisto P, Tanner M, Keinänen R, Palmberg C, et al. In vivo amplification of the androgen receptor gene and progression of human prostate cancer. *Nat Genet*. 1995;9(4):401-6.
30. Edwards J, Krishna NS, Grigor KM, Bartlett JM. Androgen receptor gene amplification and protein expression in hormone refractory prostate cancer. *Br J Cancer*. 2003;89(3):552-6.
31. Gregory CW, Johnson RT Jr, Mohler JL, French FS, Wilson EM. Androgen receptor stabilization in recurrent prostate cancer is associated with hypersensitivity to low androgen. *Cancer Res*. 2001;61(7):2892-8.
32. Tannock IF, de Wit R, Berry WR, Horti J, Pluzanska A, Chi KN, et al.; TAX 327 Investigators. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer. *N Engl J Med*. 2004;351(15):1502-12.
33. de Bono JS, Oudard S, Ozguroglu M, Hansen S, Machiels JP, Kocak I, et al.; TROPIC Investigators. Prednisone plus cabazitaxel or mitoxantrone for metastatic castration-resistant prostate cancer progressing after docetaxel treatment: a randomised open-label trial. *Lancet*. 2010;376(9747):1147-54.
34. Rathkopf DE, Smith MR, de Bono JS, Logothetis CJ, Shore ND, de Souza P, et al. Updated interim efficacy analysis and long-term safety of abiraterone acetate in metastatic castration-resistant prostate cancer patients without prior chemotherapy (COU-AA-302). *Eur Urol*. 2014;66(5):815-25.
35. Fizazi K, Scher HI, Molina A, Logothetis CJ, Chi KN, Jones RJ, et al.; COU-AA-301 Investigators. Abiraterone acetate for treatment of metastatic castration-resistant prostate cancer: final overall survival analysis of the COU-AA-301 randomised, double-blind, placebo-controlled phase 3 study. *Lancet Oncol*. 2012;13(10):983-92.
36. Beer TM, Armstrong AJ, Rathkopf DE, Loriot Y, Sternberg CN, Higano CS, et al.; PREVAIL Investigators. Enzalutamide in metastatic prostate cancer before chemotherapy. *N Engl J Med*. 2014;371(5):424-33.

37. Scher HI, Fizazi K, Saad F, Taplin ME, Sternberg CN, Miller K, et al.; AFFIRM Investigators. Increased survival with enzalutamide in prostate cancer after chemotherapy. *N Engl J Med*. 2012;367(13):1187-97.
38. Kantoff PW, Higano CS, Shore ND, Berger ER, Small EJ, Penson DF, et al.; IMPACT Study Investigators. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med*. 2010;363(5):411-22.
39. Parker C, Nilsson S, Heinrich D, O'Sullivan JM, Fossa SD, Wiechno ACPJ, et al. Updated analysis of the phase III, double-blind, randomized, multinational study of radium-223 chloride in castration-resistant prostate cancer (CRPC) patients with bone metastases (ALSYMPCA). *J Clin Oncol*. 2012;30:(18 Suppl):lba4512.
40. Petrylak DP, Tangen CM, Hussain MH, Lara PN Jr, Jones JA, Taplin ME, et al. Docetaxel and estramustine compared with mitoxantrone and prednisone for advanced refractory prostate cancer. *N Engl J Med*. 2004;351(15):1513-20.
41. Kellokumpu-Lehtinen PL, Harmenberg U, Joensuu T, McDermott R, Hervonen P, Ginman C, et al. 2-Weekly versus 3-weekly docetaxel to treat castration-resistant advanced prostate cancer: a randomised, phase 3 trial. *Lancet Oncol*. 2013;14(2):117-24.
42. Bissery MC, Vrignaud P, Riou JF, Lavelle F. In vivo isolation and characterization of a docetaxel resistant B16 melanoma. *Proc Am Assoc Cancer Res*. 1995;36:316.
43. De Bono J, Hardy-Bessard A, Ford D, Mourey L, Carles J, Parente P, et al. Phase III non-inferiority study of cabazitaxel (C) 20 mg/m² (C20) versus 25 mg/m² (C25) in patients (pts) with metastatic castration-resistant prostate cancer (mCRPC) previously treated with docetaxel (D). *J Clin Oncol*. 2016;15(Suppl):5008.
44. Ryan CJ, Smith MR, Fizazi K, Saad F, Mulders PF, Sternberg CN, et al.; COU-AA-302 Investigators. Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naive men with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind, placebo-controlled phase 3 study. *Lancet Oncol*. 2015;16(2):152-60.
45. Basch E, Autio K, Ryan CJ, Mulders P, Shore N, Kheoh T, et al. Abiraterone acetate plus prednisone versus prednisone alone in chemotherapy-naive men with metastatic castration-resistant prostate cancer: patient-reported outcome results of a randomised phase 3 trial. *Lancet Oncol*. 2013;14(12):1193-9.
46. Fizazi K, Scher HI, Molina A, Logothetis CJ, Chi KN, Jones RJ, et al.; COU-AA-301 Investigators. Abiraterone acetate for treatment of metastatic castration-resistant prostate cancer: final overall survival analysis of the COU-AA-301 randomised, double-blind, placebo-controlled phase 3 study. *Lancet Oncol*. 2012;13(10):983-92.
47. Logothetis CJ, Basch E, Molina A, Fizazi K, North SA, Chi KN, et al. Effect of abiraterone acetate and prednisone compared with placebo and prednisone on pain control and skeletal-related events in patients with metastatic castration-resistant prostate cancer: exploratory analysis of data from the COU-AA-301 randomised trial. *Lancet Oncol*. 2012;13(12):1210-7.
48. Loriot Y, Miller K, Sternberg CN, Fizazi K, De Bono JS, Chowdhury S, et al. Effect of enzalutamide on health-related quality of life, pain, and skeletal-related events in asymptomatic and minimally symptomatic, chemotherapy-naive patients with metastatic castration-resistant prostate cancer (PREVAIL): results from a randomised, phase 3 trial. *Lancet Oncol*. 2015;16(5):509-21.
49. Fizazi K, Scher HI, Miller K, Basch E, Sternberg CN, Cella D, et al. Effect of enzalutamide on time to first skeletal-related event, pain, and quality of life in men with castration-resistant prostate cancer: results from the randomised, phase 3 AFFIRM trial. *Lancet Oncol*. 2014;15(10):1147-56.
50. Hall SJ, Klotz L, Pantuck AJ, George DJ, Whitmore JB, Frohlich MW, et al. Integrated safety data from 4 randomized, double-blind, controlled trials of autologous cellular immunotherapy with sipuleucel-T in patients with prostate cancer. *J Urol*. 2011;186(3):877-81.
51. Sartor O, Coleman R, Nilsson S, Heinrich D, Helle SI, O'Sullivan JM, et al. Effect of radium-223 dichloride on symptomatic skeletal events in patients with castration-resistant prostate cancer and bone metastases: results from a phase 3, double-blind, randomized trial. *Lancet Oncol*. 2014;15(7):738-46.
52. Oudard S, Angelergues A, Efstathiou E, Gonzalez I, Gyftaki R, Delanoy N, et al. Updated results of the FLAC European database of metastatic castration resistant prostate cancer (mCRPC) patients (pts) treated with life extending therapies in post-docetaxel (D) setting. *Eur J Cancer*. 2015;51(Suppl 3):S489.
53. Dehm SM, Schmidt LJ, Heemers HV, Vessella RL, Tindall DJ. Splicing of a novel androgen receptor exon generates a constitutively active androgen receptor that mediates prostate cancer therapy resistance. *Cancer Res*. 2008;68(13):5469-77.

54. Hu R, Dunn TA, Wei S, Isharwal S, Veltri RW, Humphreys E, et al. Ligand-independent androgen receptor variants derived from splicing of cryptic exons signify hormone-refractory prostate cancer. *Cancer Res.* 2009;69(1):16-22.
55. Antonarakis ES, Lu C, Wang H, Luber B, Nakazawa M, Roeser JC, et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N Engl J Med.* 2014;371(11):1028-38.
56. Antonarakis ES, Lu C, Luber B, Wang H, Chen Y, Nakazawa M, et al. Androgen Receptor Splice Variant 7 and Efficacy of Taxane Chemotherapy in Patients with Metastatic Castration-Resistant Prostate Cancer. *JAMA Oncol.* 2015;1(5):582-91.
57. Loriot Y, Massard C, Albiges L, Di Palma M, Blanchard P, Chauchereau AB, et al. Personalizing treatment in patients with castrate-resistant prostate cancer: A study of predictive factors for secondary endocrine therapies activity. *J Clin Oncol.* 2012;30(5 Suppl):213.
58. Angelergues A, Mailliet D, Flechon A, Ozguroglu M, Mercier F, Le Moulec AGS, et al. Duration of response to androgen deprivation therapy (ADT) and efficacy of secondary hormone therapy, docetaxel (D), and cabazitaxel (C) in metastatic castration-resistant prostate cancer (mCRPC). *J Clin Oncol.* 2014;32(4 Suppl):282.
59. Huillard O, Albiges L, Eymard JC, Massard C, Di Palma M, Fizazi E, et al. Efficacy of docetaxel chemotherapy in metastatic prostate cancer (mPC) patients (pts) experiencing early castration resistance (CR). *J Clin Oncol.* 2013;31(15 Suppl):5075.
60. Oudard S, De Bono JS, Ozguroglu M, Hansen S, Machiels JP, Kocak I, et al. Impact of cabazitaxel (Cbz) + prednisone (P; CbzP) on overall survival (OS) at 2 yrs and in patients (pts) with aggressive disease: post-hoc analyses of TROPIC trial. In: *European Society for Medical Oncology Meeting; 2012 Sep 28 - Oct 2; Vienna, Austria.*
61. D'Amico AV, Chen MH, Roehl KA, Catalona WJ. Identifying patients at risk for significant versus clinically insignificant postoperative prostate-specific antigen failure. *J Clin Oncol.* 2005;23(22):4975-9.
62. van Soest RJ, de Morrée ES, Shen L, Tannock IF, Eisenberger MA, de Wit R. The initial biopsy Gleason score as a predictive marker for docetaxel survival benefit in patients with prostate cancer: data from the TAX 327 study. *J Clin Oncol.* 2013;31(Suppl 6):44.
63. Fizazi K, Flaig TW, Stöckle M, Scher HI, de Bono JS, Rathkopf DE, et al. Does Gleason score at initial diagnosis predict efficacy of abiraterone acetate therapy in patients with metastatic castration-resistant prostate cancer? An analysis of abiraterone acetate phase III trials. *Ann Oncol.* 2016;27(4):699-705.
64. Leibovici D, Spiess PE, Agarwal PK, Tu SM, Pettaway CA, Hitzhusen K, et al. Prostate cancer progression in the presence of undetectable or low serum prostate-specific antigen level. *Cancer.* 2007;109(2):198-204.
65. Moore SR, Reinberg Y, Zhang G. Small cell carcinoma of prostate: effectiveness of hormonal versus chemotherapy. *Urology.* 1992;39(5):411-6.
66. Parimi S, Eigl BJ, Sunderland K, Zulfiqar MT, Finch DL, Oja CD, et al. Effects of abiraterone (ABI) and enzalutamide (ENZA) on cognitive impairment and depressive symptoms in patients (pts) with metastatic castration-resistant prostate cancer (mCRPC). *J Clin Oncol.* 2016;34(15 Suppl):5059.
67. Ryan CJ, Crawford ED, Shore ND, Underwood W, Taplin ME, Londhe A, et al. IMAAGEN trial safety and efficacy update: Effect of abiraterone acetate and low-dose prednisone on prostate-specific antigen and radiographic disease progression in patients with non-metastatic castration-resistant prostate cancer. *J Clin Oncol.* 2016;34(15 Suppl):5061.
68. Fizazi K, Chi KN, de Bono JS, Gomella LG, Miller K, Rathkopf DE, et al. Low Incidence of Corticosteroid-associated Adverse Events on Long-term Exposure to Low-dose Prednisone Given with Abiraterone Acetate to Patients with Metastatic Castration-resistant Prostate Cancer. *Eur Urol.* 2016;70(3):438-44.
69. Sartor AO, Oudard S, Sengelov L, Daugaard G, Saad F, Hansen S, et al. Cabazitaxel vs docetaxel in chemotherapy-naïve (CN) patients with metastatic castration-resistant prostate cancer (mCRPC): A three-arm phase III study (FIRSTANA). *J Clin Oncol.* 2016;34(15 Suppl):5006.
70. Lavaud P, Gravis G, Legoupil C, Joly F, Oudard S, Priou F, et al. Efficacy and tolerance of treatments received beyond progression in men with metastatic hormone-naïve prostate cancer treated with androgen deprivation therapy (ADT) with or without docetaxel in the GETUG-AFU 15 phase III trial. *J Clin Oncol.* 2016;34(15 Suppl):5080.
71. Oudard S, Kramer G, Caffo O, Creppy L, Loriot Y, Hansen S, et al. Docetaxel rechallenge after an initial good response in patients with metastatic castration-resistant prostate cancer. *BJU Int.* 2015;115(5):744-52.
72. Bahl A, Oudard S, Tombal B, Ozguroglu M, Hansen S, Kocak I, et al.; TROPIC Investigators. Impact of cabazitaxel on 2-year survival and palliation of tumor-related pain in men with metastatic castration-resistant prostate cancer treated in the TROPIC trial. *Ann Oncol.* 2013;24(9):2402-8.
73. De Bono JS, Oudard S, Ozguroglu M, Hansen L, Machiels JH, Sartor LSAO, et al. A subgroup analysis of the TROPIC trial exploring reason for discontinuation of prior docetaxel and survival outcome of cabazitaxel in metastatic castration-resistant prostate cancer (MCRPC). *J Clin Oncol.* 2011;29(15 Suppl):4526.

74. Chi KN, Scher HI, Molina A, Logothetis C, Jones RJ, North JSA, et al. Exploratory analysis of survival benefit and prior docetaxel (D) treatment in COU-AA-301 a phase III study of abiraterone acetate (AA) plus prednisone (P) in metastatic castration-resistant prostate cancer (mCRPC). *J Clin Oncol*. 2012;30(5 Suppl):15.
75. De Bono J, Smith MR, Saad F, Rathkopf DE, Mulders PFA, Shore ESD, et al. Response to taxane chemotherapy as first subsequent therapy after abiraterone acetate (AA) in patients (pts) with metastatic castration-resistant prostate cancer (mCRPC): post-hoc analysis COU-AA-302. *J Clin Oncol*. 2015;33(7 Suppl):184.
76. Mezynski J, Pezaro C, Bianchini D, Zivi A, Sandhu S, Thompson E, et al. Antitumor activity of docetaxel following treatment with the CYP17A1 inhibitor abiraterone: clinical evidence for cross-resistance? *Ann Oncol*. 2012;23(11):2943-7.
77. Schweizer MT, Zhou XC, Wang H, Bassi S, Carducci MA, Eisenberger MA, et al. The influence of prior abiraterone treatment on the clinical activity of docetaxel in men with metastatic castration-resistant prostate cancer. *Eur Urol*. 2014;66(4):646-52.
78. Aggarwal R, Harris A, Formaker C, Small EJ, Molina A, Griffin TW, et al. Response to subsequent docetaxel in a patient cohort with metastatic castration-resistant prostate cancer after abiraterone acetate treatment. *Clin Genitourin Cancer*. 2014;12(5):e167-72.
79. Azad AA, Leibowitz-Amit R, Eigl BJ, Lester R, Wells JC, Murray RN, et al. A retrospective, Canadian multi-center study examining the impact of prior response to abiraterone acetate on efficacy of docetaxel in metastatic castration-resistant prostate cancer. *Prostate*. 2014;74(15):1544-50.
80. Azad AA, Eigl BJ, Murray RN, Kollmannsberger C, Chi KN. Efficacy of enzalutamide following abiraterone acetate in chemotherapy-naive metastatic castration-resistant prostate cancer patients. *Eur Urol*. 2015;67(1):23-9.
81. Suzman DL, Lubner B, Schweizer MT, Nadal R, Antonarakis ES. Clinical activity of enzalutamide versus docetaxel in men with castration-resistant prostate cancer progressing after abiraterone. *Prostate*. 2014;74(13):1278-85.
82. Cheng HH, Nadal R, Gulati R, Azad A, Twardowski P, Agarwal UNVN, et al. The effect of prior abiraterone (Abi) use on the activity of enzalutamide (Enza) in men with mCRPC. *J Clin Oncol*. 2014;32(4 Suppl):18.
83. Khalaf D, Annala M, Finch DL, Oja CD, Vergidis J, Zulfiqar M, et al. Phase 2 randomized crossover trial of abiraterone + prednisone (ABI+P) vs enzalutamide (ENZ) for patients (pts) with metastatic castration resistant prostate cancer (mCRPC): Results for 2nd-line therapy. *J Clin Oncol*. 2018;36:(15 Suppl):5015.
84. De Bono JS, Smith MR, Saad F, Rathkopf DE, Mulders PFA, Shore ESD, et al. Response to taxane chemotherapy as first subsequent therapy after abiraterone acetate (AA) in patients (pts) with metastatic castration-resistant prostate cancer (mCRPC): post-hoc analysis of COU-AA-302. *J Clin Oncol*. 2015;33(7 Suppl):184.
85. Schrader AJ, Boegemann M, Ohlmann CH, Schnoeller TJ, Krabbe LM, Hajili T, et al. Enzalutamide in castration-resistant prostate cancer patients progressing after docetaxel and abiraterone. *Eur Urol*. 2014;65(1):30-6.
86. Bianchini D, Lorente D, Rodriguez-Vida A, Omlin A, Pezaro C, Ferraldeschi R, et al. Antitumor activity of enzalutamide (MDV3100) in patients with metastatic castration-resistant prostate cancer (CRPC) pre-treated with docetaxel and abiraterone. *Eur J Cancer*. 2014;50(1):78-84.
87. Badrising S, van der Noort V, van Oort IM, van den Berg HP, Los M, Hamberg P, et al. Clinical activity and tolerability of enzalutamide (MDV3100) in patients with metastatic, castration-resistant prostate cancer who progress after docetaxel and abiraterone treatment. *Cancer*. 2014;120(7):968-75.
88. Lorient Y, Bianchini D, Ileana E, Sandhu S, Patrikidou A, Pezaro C, et al. Antitumor activity of abiraterone acetate against metastatic castration-resistant prostate cancer progressing after docetaxel and enzalutamide (MDV3100). *Ann Oncol*. 2013;24(7):1807-12.
89. Noonan KL, North S, Bitting RL, Armstrong AJ, Ellard SL, Chi KN. Clinical activity of abiraterone acetate in patients with metastatic castration-resistant prostate cancer progressing after enzalutamide. *Ann Oncol*. 2013;24(7):1802-7.
90. Pezaro CJ, Omlin AG, Altavilla A, Lorente D, Ferraldeschi R, Bianchini D, et al. Activity of cabazitaxel in castration resistant prostate cancer progressing after docetaxel and next-generation endocrine agents. *Eur Urol*. 2014;66(3):459-65.
91. Sella A, Sella T, Peer A, Berger R, Frank SJ, Gez E, et al. Activity of cabazitaxel after docetaxel and abiraterone acetate therapy in patients with castration-resistant prostate cancer. *Clin Genitourin Cancer*. 2014;12(6):428-32.
92. Wissing MD, Coenen JL, van den Berg P, Westgeest HM, van den Eertwegh AJ, van Oort IM, et al. CAST: a retrospective analysis of metastatic castrate-resistant prostate cancer previously treated with cabazitaxel and abiraterone acetate sequential treatment. *Int J Cancer*. 2015;136(6):E760-72.
93. Al Nakouzi N, Le Moulec S, Albighès L, Wang C, Beuzeboc P, Gross-Goupil M, et al. Cabazitaxel Remains Active in Patients Progressing After Docetaxel Followed by Novel Androgen Receptor Pathway Targeted Therapies. *Eur Urol*. 2015;68(2):228-35.

94. Maughan BL, Boudadi K, Nadal RM, Antonarakis ES. Intercalating docetaxel between novel hormone therapies (NHT) abiraterone and enzalutamide in metastatic castration resistant prostate cancer (mCRPC): Does it resensitize patients to the second NHT agent? *J Clin Oncol*. 2016;34:(15 Suppl):e16550.
95. Saad F, Carles J, Gillessen S, Heidenreich A, Heinrich D, Gratt J, et al.; Radium-223 International Early Access Program Investigators. Radium-223 and concomitant therapies in patients with metastatic castration-resistant prostate cancer: an international, early access, open-label, single-arm phase 3b trial. *Lancet Oncol*. 2016;17(9):1306-16.
96. Hussain M, Fizazi K, Saad F, Rathenborg P, Shore N, Ferreira U, et al. Enzalutamide in men with non-metastatic, castration-resistant prostate cancer. *N Engl J Med*. 2018;378(26):2465-74.
97. Clegg NJ, Wongvipat J, Joseph JD, Tran C, Ouk S, Dilhas A, et al. ARN-509: a novel antiandrogen for prostate cancer treatment. *Cancer Res*. 2012;72(6):1494-503.
98. Smith MR, Saad F, Chowdhury S, Oudard S, Hadaschik BA, Graff JN, et al. Apalutamide Treatment and Metastasis-free Survival in Prostate Cancer. *N Engl J Med*. 2018;378(15):1408-18.
99. Antonarakis ES, Lu C, Luber B, Wang H, Chen Y, Zhu Y, et al. Clinical Significance of Androgen Receptor Splice Variant-7 mRNA Detection in Circulating Tumor Cells of Men With Metastatic Castration-Resistant Prostate Cancer Treated With First- and Second-Line Abiraterone and Enzalutamide. *J Clin Oncol*. 2017;35(19):2149-56.
100. Kwon ED, Drake CG, Scher HI, Fizazi K, Bossi A, van den Eertwegh AJ, et al.; CA184-043 Investigators. Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicenter, randomized, double blind, phase 3 trial. *Lancet Oncol*. 2014;15(7):700-12.
101. Beer TM, Kwon ED, Drake CG, Fizazi K, Logothetis C, Gravis G, et al. Randomized, Double-Blind, Phase III Trial of Ipilimumab Versus Placebo in Asymptomatic or Minimally Symptomatic Patients With Metastatic Chemotherapy-Naive Castration-Resistant Prostate Cancer. *J Clin Oncol*. 2017;35(1):40-7.
102. Hansen AR, Massard C, Ott PA, Haas NB, Lopez JS, Ejadi S, et al. Pembrolizumab for advanced prostate adenocarcinoma: findings of the KEYNOTE-028 study. *Ann Oncol*. 2018;29(8):1807-13.
103. De Bono JS, Goh JCH, Ojamaa K, Rodriguez JMP, Drake CG, Hoimes CJ, et al. KEYNOTE-199: Pembrolizumab (Pembro) for docetaxel-refractory metastatic castration-resistant prostate cancer (mCRPC). *J Clin Oncol*. 2018;36(15 Suppl):5007.
104. Sharma P, Pachynski RK, Narayan V, Flechon A, Gravis G, Galsky MD, et al. Initial results from a phase II study of nivolumab (NIVO) plus ipilimumab (IPI) for the treatment of metastatic castration-resistant prostate cancer (mCRPC; CheckMate 650). *J Clin Oncol*. 2019;37(7 Suppl):142.
105. Grasso CS, Wu YM, Robinson DR, Cao X, Dhanasekaran SM, Khan AP, et al. The mutational landscape of lethal castration-resistant prostate cancer. *Nature*. 2012;487(7406):239-43.
106. Castro E, Goh C, Olmos D, Saunders E, Leongamornlert D, Tymrakiewicz M, et al. Germline BRCA mutations are associated with higher risk of nodal involvement, distant metastasis, and poor survival outcomes in prostate cancer. *J Clin Oncol*. 2013;31(14):1748-57.
107. Sandhu SK, Omlin A, Hylands L, Miranda S, Barber LJ, Riisnaes R, et al. Poly (ADP-ribose) polymerase (PARP) inhibitors for the treatment of advanced germline BRCA2 mutant prostate cancer. *Ann Oncol*. 2013;24(5):1416-8.
108. Mateo J, Carreira S, Sandhu S, Miranda S, Mossop H, Perez-Lopez R, et al. DNA-Repair defects and Olaparib in Metastatic Prostate Cancer. *N Engl J Med*. 2015;373(18):1697-708.
109. Clarke N, Wiechno P, Alekseev B, Sala N, Jones R, Kocak I, et al. Olaparib combined with abiraterone in patients with metastatic castration-resistant prostate cancer: a randomized, double blind, placebo-controlled, phase 2 trial. *Lancet Oncol*. 2018;19(7):975-86.
110. Yu EY, Massard C, Retz M, Tafreshi A, Galceran JC, Hammerer P, et al. Keynote-365 cohort a: Pembrolizumab (pembro) plus olaparib in docetaxel-pretreated patients (pts) with metastatic castrate-resistant prostate cancer (mCRPC). *J Clin Oncol*. 2019;37(7 Suppl):145.
111. Perera M, Papa N, Roberts M, Williams M, Udovicich C, Vela I, et al. Gallium-68 Prostate-specific Membrane Antigen Positron Emission Tomography in Advanced Prostate Cancer-Updated Diagnostic Utility, Sensitivity, Specificity, and Distribution of Prostate-specific Membrane Antigen-avid Lesions: A Systematic Review and Meta-analysis. *Eur Urol*. 2019;pii:S0302-2838(19)30095-8.
112. Fizazi K, Shore ND, Tammela T, Ulys A, Vjaters E, Polyakov S, et al. ARAMIS: Efficacy and safety of darolutamide in non-metastatic castration-resistant prostate cancer (nmCRPC). *J Clin Oncol*. 2019;37(7 Suppl):140.